

MATHEMATICS FOR 9TH CLASS (UNIT # 1)

Pilot Superone Mathematics 7 Class 9th

Exercise 1.1

1. Find the order of the following matrices.

$$A = \begin{bmatrix} 2 & 3 \\ -5 & 6 \end{bmatrix}, \quad B = \begin{bmatrix} 2 & 0 \\ 3 & 5 \end{bmatrix}, \quad C = [2 \quad 4]$$

$$D = \begin{bmatrix} 4 \\ 0 \\ 6 \end{bmatrix}, \quad E = \begin{bmatrix} a & d \\ b & e \\ c & f \end{bmatrix}, \quad F = [2]$$

$$G = \begin{bmatrix} 2 & 3 & 0 \\ 1 & 2 & 3 \\ 2 & 4 & 5 \end{bmatrix}, \quad H = \begin{bmatrix} 2 & 3 & 4 \\ 1 & 0 & 6 \end{bmatrix}$$

Solution: $A = \begin{bmatrix} 2 & 3 \\ -5 & 6 \end{bmatrix}$

Number of Rows (R) = 2

Number of Columns = 2

Order of the matrix = R by C

Order of the matrix = 2 by 2

$$B = \begin{bmatrix} 2 & 0 \\ 3 & 5 \end{bmatrix}$$

Order = 2 by 2

$$C = [2 \quad 4]$$

Order = 1 by 2

$$D = \begin{bmatrix} 4 \\ 0 \\ 6 \end{bmatrix}$$

Order = 3 by 1

$$E = \begin{bmatrix} a & d \\ b & e \\ c & f \end{bmatrix}$$

Order = 3 by 2

MATHEMATICS FOR 9TH CLASS (UNIT # 1)

Pilot Superone Mathematics 8

Class 9th

$$F = [2]$$

Order = 1 by 1

$$G = \begin{bmatrix} 2 & 3 & 0 \\ 1 & 2 & 3 \\ 2 & 4 & 5 \end{bmatrix}$$

Order = 3 by 3

$$H = \begin{bmatrix} 2 & 3 & 4 \\ 1 & 0 & 6 \end{bmatrix}$$

Order = 2 by 3

2. Which of the following matrices are equal?

$$A = [3], \quad B = [3 \ 5], \quad C = [5 \ -2]$$

$$D = [5 \ 3], \quad E = \begin{bmatrix} 4 & 0 \\ 6 & 2 \end{bmatrix}, \quad F = \begin{bmatrix} 2 \\ 6 \end{bmatrix}$$

$$G = \begin{bmatrix} 3-1 \\ 3+3 \end{bmatrix}, \quad H = \begin{bmatrix} 4 & 0 \\ 6 & 2 \end{bmatrix}, \quad I = [3 \ 3 + 2]$$

$$J = \begin{bmatrix} 2+2 & 2-2 \\ 2+4 & 2+0 \end{bmatrix}$$

Solution:

$$(i) \quad A = C \quad (ii) \quad B = I \quad (iii) \quad E = H = J$$

$$(iv) \quad F = G$$

Order of one is equal to the order of the other and their corresponding elements are equal.

Q.3. Find the values of a, b, c and d which satisfy the matrix equation.

$$\begin{bmatrix} a+c & a+2b \\ c-1 & 4d-6 \end{bmatrix} = \begin{bmatrix} 0 & -7 \\ 3 & 2d \end{bmatrix}$$

Solution: $a + c = 0 \quad (i)$

MATHEMATICS FOR 9TH CLASS (UNIT # 1)

Pilot Supreme Mathematics 9 Class 9th

$$c - 1 = 3 \quad (\text{ii})$$

$$a + 2b = -7 \quad (\text{iii})$$

$$4d - 6 = 2d \quad (\text{iv})$$

$$c - 1 = 3 \quad \text{From (ii)}$$

$$c = 3 + 1 = 4$$

$$a + c = 0 \quad (\text{i})$$

$$a + 4 = 0 \quad \text{From (ii)}$$

$$a = -4$$

$$4d - 6 = 2d \quad \text{From (iv)}$$

$$4d - 2d = 6$$

$$2d = 6$$

$$\therefore d = 3$$

$$a + 2b = -7 \quad \text{From (iii)}$$

Putting value of 'a'.

$$-4 + 2b = -7$$

$$2b = -7 + 4$$

$$2b = -3$$

$$b = -\frac{3}{2}$$

$$\text{Thus } a = -4, \quad b = -\frac{3}{2}, \quad c = 4, \quad d = 3$$

Types of Matrices:

Row Matrix: A row matrix has only ONE row. e.g.

$A = [3 \ -2 \ 0]$ is a row matrix.

Column Matrix: A column matrix has only ONE column e.g.

$B = \begin{bmatrix} 1 \\ 2 \\ -2 \end{bmatrix}$ is a column matrix.

Rectangular Matrix: If the number of rows is not equal to the number of columns in a matrix then it is a rectangular matrix.

MATHEMATICS FOR 9TH CLASS (UNIT # 1)

=====

Pilot Superone Mathematics 10 Class 9th

Square Matrix: A matrix is called a square matrix if its number of rows is equal to its number of columns.

Null or Zero Matrix: A matrix is called a null or zero matrix if each of its entries is zero.

Transpose of a Matrix: A matrix obtained by interchanging rows into columns or columns into rows is called a transpose of the given matrix.

Negative of a Matrix: A matrix obtained by changing the signs of all the entries of a matrix is called negative of the given matrix.

Symmetric Matrix: A square matrix is symmetric if it is equal to its transpose. If A is a square matrix then its symmetrix matric is written as A^T and $A^T = A$.

Skew-Symmetric Matrix: A square matrix A is said to be skew-symmetric if $A^T = -A$

Diagonal Matrix: A square matrix is called a diagonal matrix if atleast any one of the entries of its diagonal is not zero and non-diagonal entries must all be zero. e.g.

Scalar Matix. A diagonal matrix is called a scalar matrix, if all the diagonal entries are same and non-zero.

for example $\begin{bmatrix} k & 0 & 0 \\ 0 & k & 0 \\ 0 & 0 & k \end{bmatrix}$ where k is a constant

$$A = \begin{bmatrix} 2 & 0 & 0 \\ 0 & 3 & 0 \\ 0 & 0 & 0 \end{bmatrix}, B = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 4 & 0 \\ 0 & 0 & 5 \end{bmatrix}$$

Identity Matrix: A diagonal matrix is called identity (unit) matrix if all diagonal entries are '1'. It is denoted by I.

MATHEMATICS FOR 9TH CLASS (UNIT # 1)

=====

Pilot Superone Mathematics

11

Class 9th

Exercise 1.2

1. Find the following matrices, identify unit matrices, row matrices, column matrices and null matrices.

$$A = \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix}, B = [2 \ 3 \ 4], C = \begin{bmatrix} 4 \\ 0 \\ 6 \end{bmatrix}$$

$$D = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}, E = [0], F = \begin{bmatrix} 5 \\ 6 \\ 7 \end{bmatrix}$$

Solution: $A = \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix}$

A is a zero matrix.

$$B = [2 \ 3 \ 4]$$

B is a row matrix.

$$C = \begin{bmatrix} 4 \\ 0 \\ 6 \end{bmatrix}$$

C is a column matrix.

$$D = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$$

D is an identity (unit) matrix.

$$E = [0]$$

E is a zero matrix.

$$F = \begin{bmatrix} 5 \\ 6 \\ 7 \end{bmatrix}$$

F is a column matrix.

- Q.2. From the following matrices, identify.

- (a) Square matrices (b) Rectangular matrices
(c) Row matrices (d) Column matrices

MATHEMATICS FOR 9TH CLASS (UNIT # 1)

Pilot Superone Mathematics 12 Class 9th

(e) Identity matrices (f) Null matrices

$$\begin{array}{lll}
 (i) \begin{bmatrix} -8 & 2 & 7 \\ 12 & 0 & 4 \end{bmatrix} & (ii) \begin{bmatrix} 3 \\ 0 \\ 1 \end{bmatrix} & (iii) \begin{bmatrix} 6 & -4 \\ 3 & -2 \end{bmatrix} \\
 (iv) \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} & (v) \begin{bmatrix} 1 & 2 \\ 3 & 4 \\ 5 & 6 \end{bmatrix} & (vi) [3 \ 10 \ -1] \\
 (vii) \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix} & (viii) \begin{bmatrix} 1 & 2 & 3 \\ -1 & 2 & 0 \\ 0 & 0 & 1 \end{bmatrix} & (ix) \begin{bmatrix} 0 & 0 \\ 0 & 0 \\ 0 & 0 \end{bmatrix}
 \end{array}$$

Solution:

$$\begin{bmatrix} 6 & -4 \\ 3 & -2 \end{bmatrix}, \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}, \begin{bmatrix} 1 & 2 & 3 \\ -1 & 2 & 0 \\ 0 & 0 & 1 \end{bmatrix} \dots \text{Square matrices}$$

$$\begin{bmatrix} -8 & 2 & 7 \\ 12 & 0 & 4 \end{bmatrix}, \begin{bmatrix} 3 \\ 0 \\ 1 \end{bmatrix}, \begin{bmatrix} 1 & 2 \\ 3 & 4 \\ 5 & 6 \end{bmatrix} \dots \text{Rectangular matrices}$$

$$[3 \ 10 \ -1], \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}, \begin{bmatrix} 0 & 0 \\ 0 & 0 \\ 0 & 0 \end{bmatrix}$$

$$[3 \ 10 \ -1] \dots \text{Row matrix.}$$

$$\begin{bmatrix} 3 \\ 0 \\ 1 \end{bmatrix}, \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix} \dots \text{Column matrix}$$

$$\begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \dots \text{Identity matrix}$$

$$\begin{bmatrix} 0 & 0 \\ 0 & 0 \\ 0 & 0 \end{bmatrix} \dots \text{Zero matrix or null}$$

MATHEMATICS FOR 9TH CLASS (UNIT # 1)

Pilot Superone Mathematics 13 Class 9th

Q.3. From the following matrices, identify diagonal, scalar and unit (identity).

$$A = \begin{bmatrix} 4 & 0 \\ 0 & 4 \end{bmatrix}, B = \begin{bmatrix} 2 & 0 \\ 0 & -1 \end{bmatrix}, C = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$$

$$D = \begin{bmatrix} 3 & 0 \\ 0 & 0 \end{bmatrix}, \text{ and } E = \begin{bmatrix} 5-3 & 0 \\ 0 & 1+1 \end{bmatrix}$$

Solution:

$$A = \begin{bmatrix} 4 & 0 \\ 0 & 4 \end{bmatrix}, B = \begin{bmatrix} 2 & 0 \\ 0 & -1 \end{bmatrix} \dots \text{Diagonal matrices}$$

$$C = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}, D = \begin{bmatrix} 3 & 0 \\ 0 & 0 \end{bmatrix}, E = \begin{bmatrix} 5-3 & 0 \\ 0 & 1+1 \end{bmatrix}$$

$$A = \begin{bmatrix} 4 & 0 \\ 0 & 4 \end{bmatrix}, E = \begin{bmatrix} 5-3 & 0 \\ 0 & 1+1 \end{bmatrix} \dots \text{Scalar matrices}$$

$$C = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \dots \text{Identity matrix}$$

Q.4. Find negative of matrices A, B, C, D and F when:

$$A = \begin{bmatrix} 1 \\ 0 \\ -1 \end{bmatrix}, B = \begin{bmatrix} 3 & -1 \\ 2 & 1 \end{bmatrix}, C = \begin{bmatrix} 2 & 6 \\ 3 & 2 \end{bmatrix}$$

$$D = \begin{bmatrix} -3 & 2 \\ -4 & 5 \end{bmatrix}, E = \begin{bmatrix} 1 & -5 \\ 2 & 3 \end{bmatrix}$$

Solution:

$$A = \begin{bmatrix} 1 \\ 0 \\ -1 \end{bmatrix} \text{ sign of every element is changed.}$$

MATHEMATICS FOR 9TH CLASS (UNIT # 1)

Pilot Superone Mathematics 14 Class 9th

$$-A = -\begin{bmatrix} 1 \\ 0 \\ -1 \end{bmatrix} = \begin{bmatrix} -1 \\ 0 \\ 1 \end{bmatrix}$$

$$B = \begin{bmatrix} 3 & -1 \\ 2 & 1 \end{bmatrix}$$

$$-B = -\begin{bmatrix} 3 & -1 \\ 2 & 1 \end{bmatrix} = \begin{bmatrix} -3 & 1 \\ -2 & -1 \end{bmatrix}$$

$$C = \begin{bmatrix} 2 & 6 \\ 3 & -2 \end{bmatrix}$$

$$-C = -\begin{bmatrix} 2 & 6 \\ 3 & -2 \end{bmatrix} = \begin{bmatrix} -2 & -6 \\ -3 & 2 \end{bmatrix}$$

$$D = \begin{bmatrix} -3 & 2 \\ -4 & 5 \end{bmatrix}$$

$$-D = -\begin{bmatrix} -3 & 2 \\ -4 & 5 \end{bmatrix} = \begin{bmatrix} 3 & -2 \\ 4 & -5 \end{bmatrix}$$

$$E = \begin{bmatrix} 1 & -5 \\ 2 & 3 \end{bmatrix}$$

$$-E = -\begin{bmatrix} 1 & -5 \\ 2 & 3 \end{bmatrix} = \begin{bmatrix} -1 & 5 \\ -2 & -3 \end{bmatrix}$$

Q.5. Find the transpose of each of the following matrices.

$$A = \begin{bmatrix} 0 \\ 1 \\ -2 \end{bmatrix}, B = [5 \quad 1 \quad -6], C = \begin{bmatrix} 1 & 2 \\ 2 & -1 \\ 3 & 0 \end{bmatrix}$$

$$D = \begin{bmatrix} 2 & 3 \\ 0 & 5 \end{bmatrix}, E = \begin{bmatrix} 2 & 3 \\ -4 & 5 \end{bmatrix}, F = \begin{bmatrix} 1 & 2 \\ 3 & 4 \end{bmatrix}$$

MATHEMATICS FOR 9TH CLASS (UNIT # 1)

Pilot Superone Mathematics

15

Class 9th

$$A = \begin{bmatrix} 0 \\ 1 \\ -2 \end{bmatrix} \text{ columns are changed into rows}$$

$$A^t = [0 \quad 1 \quad -2]$$

$$B = [5 \quad 1 \quad -6] ; B^t = \begin{bmatrix} 5 \\ 1 \\ -6 \end{bmatrix}$$

$$C = \begin{bmatrix} 1 & 2 \\ 2 & -1 \\ 3 & 0 \end{bmatrix} ; C^t = \begin{bmatrix} 1 & 2 & 3 \\ 2 & -1 & 0 \end{bmatrix}$$

$$D = \begin{bmatrix} 2 & 3 \\ 0 & 5 \end{bmatrix} ; D^t = \begin{bmatrix} 2 & 0 \\ 3 & 5 \end{bmatrix}$$

$$E = \begin{bmatrix} 2 & 3 \\ -4 & 5 \end{bmatrix} ; E^t = \begin{bmatrix} 2 & -4 \\ 3 & 5 \end{bmatrix}$$

$$F = \begin{bmatrix} 1 & 2 \\ 3 & 4 \end{bmatrix} ; F^t = \begin{bmatrix} 1 & 3 \\ 2 & 4 \end{bmatrix}$$

Q.6. Verify that if $A = \begin{bmatrix} 1 & 2 \\ 0 & 1 \end{bmatrix}$, $B = \begin{bmatrix} 1 & 1 \\ 2 & 0 \end{bmatrix}$ then

(i) $(A^t)^t = A$ (ii) $(B^t)^t = B$

Solution:

$$6(i) \quad A = \begin{bmatrix} 1 & 2 \\ 0 & 1 \end{bmatrix}$$

$$A^t = \begin{bmatrix} 1 & 0 \\ 2 & 1 \end{bmatrix}$$

www

MATHEMATICS FOR 9TH CLASS (UNIT # 1)

Pilot Success Mathematics 16 Class 9th

$$(A^t)^t = \begin{bmatrix} 1 & 2 \\ 0 & 1 \end{bmatrix} = A$$

$$6(ii) \quad B = \begin{bmatrix} 1 & 1 \\ 2 & 0 \end{bmatrix}$$

$$B^t = \begin{bmatrix} 1 & 2 \\ 1 & 0 \end{bmatrix}$$

$$(B^t)^t = \begin{bmatrix} 1 & 1 \\ 2 & 0 \end{bmatrix} = B$$

Addition and subtraction of Matrices

Remember:

- (i) Two matrices are conformable for addition if they have the same order.
- (ii) In case of addition, corresponding elements are added.
- (iii) In case of subtraction, corresponding elements are subtracted in an order.

Commutative and Associative Laws of Addition of Matrices:

- (a) Commutative law under Addition.

$$A + B = B + A$$

- (b) Associative law under addition.

$$(A + B) + C = A + (B + C)$$

Additive identity of a matrix

For any matrix A and zero matrix O of the same order. O is called additive identity of A as

$$A + O = O + A$$

Additive Inverse of a matrix

If A and B are two matrices of the same order such that

$$A + B = O = B + A$$

then A and B are called additive inverse of each other.

We change the sign of all the elements of a matrix to get its

additive inverse

MATHEMATICS FOR 9TH CLASS (UNIT # 1)

Pilot Superone Mathematics

17

Class 9th

Exercise 1.3

1. Which of the following matrices are conformable for addition?

$$A = \begin{bmatrix} 2 & 1 \\ 1 & 3 \end{bmatrix}, B = \begin{bmatrix} 3 \\ 1 \end{bmatrix}, C = \begin{bmatrix} 1 & 0 \\ 2 & -1 \\ 1 & 2 \end{bmatrix}$$

$$D = \begin{bmatrix} 2+1 \\ 3 \end{bmatrix}, E = \begin{bmatrix} 1 & 0 \\ 1 & 2 \end{bmatrix}, F = \begin{bmatrix} 3 & 2 \\ 1+1 & 4 \\ 3+2 & 2+1 \end{bmatrix}$$

Solution

order of A = 2 by 2 order of B = 2 by 1

order of C = 3 by 2 order of D = 2 by 1

order of E = 2 by 2 order of F = 3 by 2

Matrices of the same order are conformable for addition. A and E, B and D, C and F are conformable for addition.

2. Find the additive inverse of the following matrices.

$$A = \begin{bmatrix} 2 & 4 \\ -2 & 1 \end{bmatrix}, B = \begin{bmatrix} 1 & 0 & -1 \\ 2 & -1 & 3 \\ 3 & -2 & 1 \end{bmatrix}, C = \begin{bmatrix} 4 \\ -2 \end{bmatrix}$$

$$D = \begin{bmatrix} 1 & 0 \\ 3 & -2 \\ 2 & 1 \end{bmatrix}, E = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}, F = \begin{bmatrix} \sqrt{3} & 1 \\ -1 & \sqrt{2} \end{bmatrix}$$

Solution:

$$A = \begin{bmatrix} 2 & 4 \\ -2 & 1 \end{bmatrix}$$

$$-A = \begin{bmatrix} 2 & 4 \\ 2 & 1 \end{bmatrix} = \begin{bmatrix} -2 & -4 \\ 2 & -1 \end{bmatrix} \text{ (Additive inverse of A)}$$

$$B = \begin{bmatrix} 1 & 0 & -1 \\ 2 & -1 & 3 \\ 3 & 2 & 1 \end{bmatrix}$$

MATHEMATICS FOR 9TH CLASS (UNIT # 1)

Pilot Superone Mathematics

18

Class 9th

$$B = - \begin{bmatrix} 1 & 0 & -1 \\ 2 & -1 & 3 \\ 3 & -2 & 1 \end{bmatrix} = \begin{bmatrix} -1 & 0 & 1 \\ -2 & 1 & -3 \\ 3 & 2 & -1 \end{bmatrix} \text{ (Additive inverse of B)}$$

$$C = \begin{bmatrix} 4 \\ -2 \end{bmatrix}$$

$$-C = - \begin{bmatrix} 4 \\ -2 \end{bmatrix} = \begin{bmatrix} -4 \\ 2 \end{bmatrix} \text{ (additive inverse of C)}$$

$$D = \begin{bmatrix} 1 & 0 \\ -3 & -2 \\ 2 & 1 \end{bmatrix}$$

$$-D = - \begin{bmatrix} 1 & 0 \\ -3 & -2 \\ 2 & 1 \end{bmatrix} = \begin{bmatrix} -1 & 0 \\ 3 & 2 \\ -2 & -1 \end{bmatrix} \text{ (additive inverse of D)}$$

$$E = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$$

$$-E = - \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} = \begin{bmatrix} -1 & 0 \\ 0 & -1 \end{bmatrix} \text{ (additive inverse of E)}$$

$$F = \begin{bmatrix} \sqrt{3} & 1 \\ -1 & \sqrt{2} \end{bmatrix}$$

$$-F = - \begin{bmatrix} \sqrt{3} & 1 \\ -1 & \sqrt{2} \end{bmatrix} = \begin{bmatrix} -\sqrt{3} & -1 \\ 1 & -\sqrt{2} \end{bmatrix} \text{ (additive inverse of F)}$$

MATHEMATICS FOR 9TH CLASS (UNIT # 1)

Pilot Superone Mathematics 19 Class 9th

3. If

$$A = \begin{bmatrix} 1 & 2 \\ 2 & 1 \end{bmatrix}, B = \begin{bmatrix} 1 \\ -1 \end{bmatrix}, C = [1 \ 1 \ 2],$$

$$D = \begin{bmatrix} 1 & 2 & 3 \\ -1 & 0 & 2 \end{bmatrix} \text{ then find}$$

$$(i) \quad A + \begin{bmatrix} 1 & 1 \\ 1 & 1 \end{bmatrix} \quad (ii) \quad B + \begin{bmatrix} 2 \\ 3 \end{bmatrix}$$

$$(iii) \quad C + [-2 \ 1 \ 3] \quad (iv) \quad D + \begin{bmatrix} 0 & 1 & 0 \\ 2 & 0 & 1 \end{bmatrix}$$

$$(v) \quad 2A \quad (vi) \quad (-1)B$$

$$(vii) \quad (-2)C \quad (viii) \quad 3D \quad (ix) \quad 3C$$

Solution:

$$(i) \quad A + \begin{bmatrix} 1 & 1 \\ 1 & 1 \end{bmatrix} = \begin{bmatrix} -1 & 2 \\ 2 & 1 \end{bmatrix} + \begin{bmatrix} 1 & 1 \\ 1 & 1 \end{bmatrix} \quad \text{Putting } A = \begin{bmatrix} 1 & 2 \\ 2 & 1 \end{bmatrix}$$

$$= \begin{bmatrix} 1+1 & 2+1 \\ 2+1 & 1+1 \end{bmatrix} = \begin{bmatrix} 0 & 3 \\ 3 & 2 \end{bmatrix}$$

$$(ii) \quad B + \begin{bmatrix} -2 \\ 3 \end{bmatrix} = \begin{bmatrix} 1 \\ -1 \end{bmatrix} + \begin{bmatrix} -2 \\ 3 \end{bmatrix} \quad \text{Putting } B = \begin{bmatrix} 1 \\ -1 \end{bmatrix}$$

MATHEMATICS FOR 9TH CLASS (UNIT # 1)

Pilot Superone Mathematics 20 Class 9th

$$= \begin{bmatrix} 1 & -2 \\ -1 & +3 \end{bmatrix} = \begin{bmatrix} -1 \\ 2 \end{bmatrix}$$

(iii) $C + \begin{bmatrix} 2 & 1 & 3 \end{bmatrix}$
 $= \begin{bmatrix} 1 & -1 & 2 \end{bmatrix} + \begin{bmatrix} -2 & 1 & 3 \end{bmatrix}$ Putting $C = \begin{bmatrix} 1 & -1 & 2 \end{bmatrix}$
 $= \begin{bmatrix} 1-2 & -1+1 & 2+3 \end{bmatrix} = \begin{bmatrix} -1 & 0 & 5 \end{bmatrix}$

(iv) $D + \begin{bmatrix} 0 & 1 & 0 \\ 2 & 0 & 1 \end{bmatrix}$
 $= \begin{bmatrix} 1 & 2 & 3 \\ 1 & 0 & 2 \end{bmatrix} + \begin{bmatrix} 0 & 1 & 0 \\ 2 & 0 & 1 \end{bmatrix}$ Putting $D = \begin{bmatrix} 1 & 2 & 3 \\ -1 & 0 & 2 \end{bmatrix}$
 $= \begin{bmatrix} 1+0 & 2+1 & 3+0 \\ 1+2 & 0+0 & 2+1 \end{bmatrix} = \begin{bmatrix} 1 & 3 & 3 \\ 3 & 0 & 3 \end{bmatrix}$

(v) $2A$
 $= 2 \begin{bmatrix} -1 & 2 \\ 2 & 1 \end{bmatrix}$ Putting $A = \begin{bmatrix} -1 & 2 \\ 2 & 1 \end{bmatrix}$
 $= \begin{bmatrix} 2(-1) & 2(2) \\ 2(2) & 2(1) \end{bmatrix} = \begin{bmatrix} -2 & 4 \\ 4 & 2 \end{bmatrix}$

(vi) $(-1)B$
 $= (-1) \begin{bmatrix} 1 \\ 1 \end{bmatrix}$ Putting $B = \begin{bmatrix} 1 \\ -1 \end{bmatrix}$
 $= \begin{bmatrix} (-1)(1) \\ (-1)(-1) \end{bmatrix} = \begin{bmatrix} -1 \\ 1 \end{bmatrix}$

(vii) $(-2)C$
 $= (-2) \begin{bmatrix} 1 & 1 & 2 \end{bmatrix}$ Putting $C = \begin{bmatrix} 1 & -1 & 2 \end{bmatrix}$

MATHEMATICS FOR 9TH CLASS (UNIT # 1)

Pt. of Superone Mathematics 21 Class 9th

$$= [(-2)(1) \quad (-2)(-1) \quad (-2)(2)] = [2 \quad 2 \quad -4]$$

(viii) 3 D

$$= 3 \begin{bmatrix} 1 & 2 & 3 \\ -1 & 0 & 2 \end{bmatrix} \quad \text{Putting } D = \begin{bmatrix} 1 & 2 & 3 \\ 1 & 0 & 2 \end{bmatrix}$$

$$= \begin{bmatrix} 3(1) & 3(2) & 3(3) \\ 3(-1) & 3(0) & 3(2) \end{bmatrix} = \begin{bmatrix} 3 & 6 & 9 \\ 3 & 0 & 6 \end{bmatrix}$$

(ix) 3 C

$$= 3 \begin{bmatrix} 1 & -1 & 2 \end{bmatrix} \quad \text{Putting } C = \begin{bmatrix} 1 & -1 & 2 \end{bmatrix}$$

$$= [3(1) \quad 3(-1) \quad 3(2)] = [3 \quad -3 \quad 6]$$

4. Perform the indicated operations and simplify the following.

$$(i) \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} + \begin{bmatrix} 0 & 2 \\ 3 & 0 \end{bmatrix} + \begin{bmatrix} 1 & 1 \\ 1 & 0 \end{bmatrix}$$

$$(ii) \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} + \begin{bmatrix} 0 & 2 \\ 3 & 0 \end{bmatrix} - \begin{bmatrix} 1 & 1 \\ 1 & 0 \end{bmatrix}$$

$$(iii) [2 \ 3 \ .] + ([1 \ 0 \ 2]) - [2 \ 2 \ 2])$$

$$(iv) \begin{bmatrix} 1 & 2 & 3 \\ -1 & -1 & -1 \\ 0 & 1 & 2 \end{bmatrix} + \begin{bmatrix} 1 & 1 & 1 \\ 2 & 2 & 2 \\ 3 & 3 & 3 \end{bmatrix}$$

$$(v) \begin{bmatrix} 1 & 2 & 3 \\ 2 & 3 & 1 \\ 3 & 1 & 2 \end{bmatrix} + \begin{bmatrix} 1 & 0 & 2 \\ -2 & 1 & 0 \\ 0 & 2 & -1 \end{bmatrix}$$

MATHEMATICS FOR 9TH CLASS (UNIT # 1)

Pilot Superone Mathematics

22

Class 9th

$$(vi) \begin{bmatrix} 1 & 2 \\ 0 & 1 \end{bmatrix} + \begin{bmatrix} 2 & 1 \\ 1 & 0 \end{bmatrix} + \begin{bmatrix} 1 & 1 \\ 1 & 1 \end{bmatrix}$$

Solution:

$$4 (i) \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} + \begin{bmatrix} 0 & 2 \\ 3 & 0 \end{bmatrix} + \begin{bmatrix} 1 & 1 \\ 1 & 0 \end{bmatrix}$$

$$= \left(\begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} + \begin{bmatrix} 0 & 2 \\ 3 & 0 \end{bmatrix} \right) + \begin{bmatrix} 1 & 1 \\ 1 & 0 \end{bmatrix}$$

$$= \begin{bmatrix} 1+0 & 0+2 \\ 0+3 & 1+0 \end{bmatrix} + \begin{bmatrix} 1 & 1 \\ 1 & 0 \end{bmatrix}$$

$$= \begin{bmatrix} 1 & 2 \\ 3 & 1 \end{bmatrix} + \begin{bmatrix} 1 & 1 \\ 1 & 0 \end{bmatrix} = \begin{bmatrix} 1+1 & 2+1 \\ 3+1 & 1+0 \end{bmatrix} = \begin{bmatrix} 2 & 3 \\ 4 & 1 \end{bmatrix}$$

$$4 (ii) \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} + \begin{bmatrix} 0 & 2 \\ 3 & 0 \end{bmatrix} - \begin{bmatrix} 1 & 1 \\ 1 & 0 \end{bmatrix}$$

$$= \left(\begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} + \begin{bmatrix} 0 & 2 \\ 3 & 0 \end{bmatrix} \right) - \begin{bmatrix} 1 & 1 \\ 1 & 0 \end{bmatrix}$$

$$= \begin{bmatrix} 1+0 & 0+2 \\ 0+3 & 1+0 \end{bmatrix} - \begin{bmatrix} 1 & 1 \\ 1 & 0 \end{bmatrix}$$

$$= \begin{bmatrix} 1 & 2 \\ 3 & 1 \end{bmatrix} - \begin{bmatrix} 1 & 1 \\ 1 & 0 \end{bmatrix} = \begin{bmatrix} 1-1 & 2-1 \\ 3-1 & 1-0 \end{bmatrix} = \begin{bmatrix} 0 & 1 \\ 2 & 1 \end{bmatrix}$$

$$4 (iii) \begin{bmatrix} 2 & 3 & 1 \end{bmatrix} + ([1 \ 0 \ 2] \begin{bmatrix} 2 & 2 & 2 \end{bmatrix})$$

$$= \begin{bmatrix} 2 & 3 & 1 \end{bmatrix} + ([1-2 \ 0 \ 2 \ 2 \ 2])$$

$$\begin{bmatrix} 2 & 3 & 1 \end{bmatrix} + [-1 \ -2 \ 0]$$

$$\begin{bmatrix} 2 & 1 & 3 & 2 & 1+0 \end{bmatrix} = \begin{bmatrix} 1 & 1 & 1 \end{bmatrix}$$

MATHEMATICS FOR 9TH CLASS (UNIT # 1)

Pot Superone Mathematics

23

Class 9th

$$4 \text{ (iv)} \quad \begin{bmatrix} 1 & 2 & 3 \\ -1 & -1 & -1 \\ 0 & 1 & 2 \end{bmatrix} + \begin{bmatrix} 1 & 1 & 1 \\ 2 & 2 & 2 \\ 3 & 3 & 3 \end{bmatrix}$$

$$= \begin{bmatrix} 1+1 & 2+1 & 3+1 \\ -1+2 & -1+2 & -1+2 \\ 0+3 & 1+3 & 2+3 \end{bmatrix} = \begin{bmatrix} 2 & 3 & 4 \\ 1 & 1 & 1 \\ 3 & 4 & 5 \end{bmatrix}$$

$$4 \text{ (v)} \quad \begin{bmatrix} 1 & 2 & 3 \\ 2 & 3 & 1 \\ 3 & 1 & 2 \end{bmatrix} + \begin{bmatrix} 1 & 0 & -2 \\ 2 & -1 & 0 \\ 0 & 2 & -1 \end{bmatrix}$$

$$= \begin{bmatrix} 1+1 & 2+0 & 3-2 \\ 2+2 & 3-1 & 1+0 \\ 3+0 & 1+2 & 2-1 \end{bmatrix} = \begin{bmatrix} 2 & 2 & 1 \\ 4 & 2 & 1 \\ 3 & 3 & 1 \end{bmatrix}$$

$$4 \text{ (vi)} \quad \begin{bmatrix} 1 & 2 \\ 0 & 1 \end{bmatrix} + \begin{bmatrix} 2 & 1 \\ 1 & 0 \end{bmatrix} + \begin{bmatrix} 1 & 1 \\ 1 & 1 \end{bmatrix}$$

$$= \left(\begin{bmatrix} 1 & 2 \\ 0 & 1 \end{bmatrix} + \begin{bmatrix} 2 & 1 \\ 1 & 0 \end{bmatrix} \right) + \begin{bmatrix} 1 & 1 \\ 1 & 1 \end{bmatrix}$$

$$= \begin{bmatrix} 1+2 & 2+1 \\ 0+1 & 1+0 \end{bmatrix} + \begin{bmatrix} 1 & 1 \\ 1 & 1 \end{bmatrix}$$

$$= \begin{bmatrix} 3 & 3 \\ 1 & 1 \end{bmatrix} + \begin{bmatrix} 1 & 1 \\ 1 & 1 \end{bmatrix}$$

$$= \begin{bmatrix} 3+1 & 3+1 \\ 1+1 & 1+1 \end{bmatrix} = \begin{bmatrix} 4 & 4 \\ 2 & 2 \end{bmatrix}$$

5. For the matrices

$$A = \begin{bmatrix} 1 & 2 & 3 \\ 2 & 3 & 1 \\ 1 & 1 & 0 \end{bmatrix}, B = \begin{bmatrix} 1 & -1 & 1 \\ 2 & -2 & 2 \\ 3 & 1 & 3 \end{bmatrix} \text{ and } C = \begin{bmatrix} -1 & 0 & 0 \\ 0 & 2 & 3 \\ 1 & 1 & 2 \end{bmatrix}$$

verify the following rules:

(i) $A + C = C + A$

(ii) $A + B = B + A$

MATHEMATICS FOR 9TH CLASS (UNIT # 1)

Plot Superone Mathematics 24 Class 9th

(iii) $B + C = C + B$ (iv) $A + (B + A) = 2A + B$

(v) $(C - B) + A = C + (A - B)$

(vi) $2A + B = A + (A + B)$

(vii) $(C - B) - A = (C - A) - B$

(viii) $(A + B) + C = A + (B + C)$

(ix) $A + (B - C) = (A - C) + B$

(x) $2A + 2B = 2(A + B)$

Solution:

5 (i) $A + C = C + A$

L.H.S = $A + C$

$$= \begin{bmatrix} 1 & 2 & 3 \\ 2 & 3 & 1 \\ 1 & -1 & 0 \end{bmatrix} + \begin{bmatrix} -1 & 0 & 0 \\ 0 & -2 & 3 \\ 1 & 1 & 2 \end{bmatrix}$$

$$\begin{bmatrix} 1-1 & 2+0 & 3+0 \\ 2+0 & 3-2 & 1+3 \\ 1+1 & -1+1 & 0+2 \end{bmatrix} = \begin{bmatrix} 0 & 2 & 3 \\ 2 & 1 & 4 \\ 2 & 0 & 2 \end{bmatrix} \quad (i)$$

R.H.S = $C + A$

$$= \begin{bmatrix} 1 & 0 & 0 \\ 0 & -2 & 3 \\ 1 & 1 & 2 \end{bmatrix} + \begin{bmatrix} 1 & 2 & 3 \\ 2 & 3 & 1 \\ 1 & -1 & 0 \end{bmatrix}$$

$$= \begin{bmatrix} -1+1 & 0+2 & 0+3 \\ 0+2 & -2+3 & 3+1 \\ 1+1 & 1-1 & 2+0 \end{bmatrix} = \begin{bmatrix} 0 & 2 & 3 \\ 2 & 1 & 4 \\ 2 & 0 & 2 \end{bmatrix} \quad (ii)$$

$A + C = C + A$ From (i) and (ii)

5 (ii) $A + B = B + A$

L.H.S = $A + B$

$$= \begin{bmatrix} 1 & 2 & 3 \\ 2 & 3 & 1 \\ 1 & -1 & 0 \end{bmatrix} + \begin{bmatrix} 1 & 1 & 1 \\ 2 & 2 & 2 \\ 3 & 1 & 3 \end{bmatrix}$$

$$= \begin{bmatrix} 1+1 & 2+1 & 3+1 \\ 2+2 & 3+2 & 1+2 \\ 1+3 & -1+1 & 0+3 \end{bmatrix} = \begin{bmatrix} 2 & 3 & 4 \\ 4 & 5 & 3 \\ 4 & 0 & 3 \end{bmatrix} \quad (i)$$

MATHEMATICS FOR 9TH CLASS (UNIT # 1)

Pilot Superone Mathematics

25

Class 9th

$$\begin{aligned} \text{R.H.S} &= \begin{bmatrix} 1 & 1 & 1 \\ 2 & -2 & 2 \\ 3 & 1 & 3 \end{bmatrix} + \begin{bmatrix} 1 & 2 & 3 \\ 2 & 3 & 1 \\ 1 & -1 & 0 \end{bmatrix} \\ &= \begin{bmatrix} 1+1 & 1+2 & 1+3 \\ 2+2 & -2+3 & 2+1 \\ 3+1 & 1-1 & 3+0 \end{bmatrix} = \begin{bmatrix} 2 & 1 & 4 \\ 4 & 1 & 3 \\ 4 & 0 & 3 \end{bmatrix} \quad (ii) \end{aligned}$$

$$A + B = B + A \quad \text{From (i) and (ii),}$$

$$\text{S(iii)} \quad B + C = C + B$$

$$\text{L.H.S} = B + C$$

$$\begin{aligned} &= \begin{bmatrix} 1 & 1 & 1 \\ 2 & -2 & 2 \\ 3 & 1 & 3 \end{bmatrix} + \begin{bmatrix} 1 & 0 & 1 \\ 0 & 2 & 3 \\ 1 & 1 & 2 \end{bmatrix} \\ &= \begin{bmatrix} 1+1 & -1+0 & 1+0 \\ 2+0 & 2+2 & 2+3 \\ 3+1 & 1+1 & 3+2 \end{bmatrix} = \begin{bmatrix} 0 & -1 & 1 \\ 2 & 4 & 5 \\ 4 & 2 & 5 \end{bmatrix} \quad (i) \end{aligned}$$

$$\text{R.H.S} = C + B$$

$$\begin{aligned} &= \begin{bmatrix} -1 & 0 & 0 \\ 0 & -2 & 3 \\ 1 & 1 & 2 \end{bmatrix} + \begin{bmatrix} 1 & -1 & 1 \\ 2 & -2 & 2 \\ 3 & 1 & 3 \end{bmatrix} \\ &= \begin{bmatrix} -1+1 & 0-1 & 0+1 \\ 0+2 & -2-2 & 3+2 \\ 1+3 & 1+1 & 2+3 \end{bmatrix} = \begin{bmatrix} 0 & -1 & 1 \\ 2 & -4 & 5 \\ 4 & 2 & 5 \end{bmatrix} \quad (ii) \end{aligned}$$

$$B + C = C + B \quad \text{From (i) and (ii)}$$

$$\text{S(iv)} \quad A + (B + A) = 2A + B$$

$$\text{L.H.S} = A + (B + A)$$

$$\begin{aligned} &= \begin{bmatrix} 1 & 2 & 3 \\ 2 & 3 & 1 \\ 1 & 1 & 0 \end{bmatrix} + \left(\begin{bmatrix} 1 & 1 & 1 \\ 2 & -2 & 2 \\ 3 & 1 & 3 \end{bmatrix} + \begin{bmatrix} 1 & 2 & 3 \\ 2 & 3 & 1 \\ 1 & 1 & 0 \end{bmatrix} \right) \\ &= \begin{bmatrix} 1 & 2 & 3 \\ 2 & 3 & 1 \\ 1 & -1 & 0 \end{bmatrix} + \begin{bmatrix} 1+1 & -1+2 & 1+3 \\ 2+2 & -2+3 & 2+1 \\ 3+1 & 1-1 & 3+0 \end{bmatrix} \end{aligned}$$

MATHEMATICS FOR 9TH CLASS (UNIT # 1)

Pilot Superone Mathematics 26 Class 9th

$$= \begin{bmatrix} 1 & 2 & 3 \\ 2 & 3 & 1 \\ 1 & -1 & 0 \end{bmatrix} + \begin{bmatrix} 2 & 1 & 4 \\ 4 & 1 & 3 \\ 4 & 0 & 3 \end{bmatrix}$$

$$= \begin{bmatrix} 1+2 & 2+1 & 3+4 \\ 2+4 & 3+1 & 1+3 \\ 1+4 & -1+0 & 0+3 \end{bmatrix}$$

$$= \begin{bmatrix} 3 & 3 & 7 \\ 6 & 4 & 4 \\ 5 & -1 & 3 \end{bmatrix} \quad (i)$$

R H S = 2A + B

$$= 2 \begin{bmatrix} 1 & 2 & 3 \\ 2 & 3 & 1 \\ 1 & -1 & 0 \end{bmatrix} + \begin{bmatrix} 1 & 1 & 1 \\ 2 & -2 & 2 \\ 3 & 1 & 3 \end{bmatrix}$$

$$= \begin{bmatrix} 2(1) & 2(2) & 2(3) \\ 2(2) & 2(3) & 2(1) \\ 2(1) & 2(-1) & 2(0) \end{bmatrix} + \begin{bmatrix} 1 & 1 & 1 \\ 2 & -2 & 2 \\ 3 & 1 & 3 \end{bmatrix}$$

$$= \begin{bmatrix} 2 & 4 & 6 \\ 4 & 6 & 2 \\ 2 & -2 & 0 \end{bmatrix} + \begin{bmatrix} 1 & 1 & 1 \\ 2 & -2 & 2 \\ 3 & 1 & 3 \end{bmatrix}$$

$$= \begin{bmatrix} 2+1 & 4+1 & 6+1 \\ 4+2 & 6-2 & 2+2 \\ 2+3 & -2+1 & 0+3 \end{bmatrix}$$

$$= \begin{bmatrix} 3 & 5 & 7 \\ 6 & 4 & 4 \\ 5 & -1 & 3 \end{bmatrix} \quad (ii)$$

$$A + (B + A) = 2A + B$$

$$5(v) \quad (C - B) + A = C + (A - B)$$

$$L H S \quad (C - B) + A$$

$$= \left(\begin{bmatrix} 1 & 0 & 0 \\ 0 & 2 & 3 \\ 1 & 1 & 2 \end{bmatrix} - \begin{bmatrix} 1 & -1 & 1 \\ 2 & -2 & 2 \\ 3 & 1 & 3 \end{bmatrix} \right) + \begin{bmatrix} 1 & 2 & 3 \\ 2 & 3 & 1 \\ 1 & 1 & 0 \end{bmatrix}$$

MATHEMATICS FOR 9TH CLASS (UNIT # 1)

Pilot Superzone Mathematics 27 Class 9th

$$= \begin{bmatrix} -1-1 & 0+1 & 0-1 \\ 0-2 & -2+2 & 3-2 \\ 1 & 3 & 1-1 \end{bmatrix} + \begin{bmatrix} 1 & 2 & 3 \\ 2 & 3 & 1 \\ 1 & -1 & 0 \end{bmatrix}$$

$$= \begin{bmatrix} -2 & 1 & -1 \\ -2 & 0 & 1 \\ -2 & 0 & -1 \end{bmatrix} + \begin{bmatrix} 1 & 2 & 3 \\ 2 & 3 & 1 \\ 1 & -1 & 0 \end{bmatrix}$$

$$= \begin{bmatrix} -2+1 & 1+2 & -1+3 \\ -2+2 & 0+3 & 1+1 \\ -2+1 & 0 & -1+0 \end{bmatrix} = \begin{bmatrix} -1 & 3 & 2 \\ 0 & 3 & 2 \\ -1 & -1 & -1 \end{bmatrix} \quad (v)$$

$$\text{R.H.S} = \begin{bmatrix} -1 & 0 & 0 \\ 0 & -2 & 3 \\ 1 & 1 & 2 \end{bmatrix} + \left(\begin{bmatrix} 1 & 2 & 3 \\ 2 & 3 & 1 \\ 1 & -1 & 0 \end{bmatrix} - \begin{bmatrix} 1 & -1 & 1 \\ 2 & 2 & 2 \\ 3 & 1 & 3 \end{bmatrix} \right)$$

$$= \begin{bmatrix} -1 & 0 & 0 \\ 0 & -2 & 3 \\ 1 & 1 & 2 \end{bmatrix} + \begin{bmatrix} 1-1 & 2-1 & 3-1 \\ 2-2 & 3-2 & 1-2 \\ 1-3 & -1-1 & 0-3 \end{bmatrix}$$

$$= \begin{bmatrix} 1 & 0 & 0 \\ 0 & 2 & 3 \\ 1 & 1 & 2 \end{bmatrix} + \begin{bmatrix} 0 & 3 & 2 \\ 0 & 5 & -1 \\ 2 & -2 & -3 \end{bmatrix}$$

$$= \begin{bmatrix} -1+0 & 0+3 & 0+2 \\ 0+0 & -2+5 & 3-1 \\ 1-2 & 1-2 & 2-3 \end{bmatrix}$$

$$= \begin{bmatrix} -1 & 3 & 2 \\ 0 & 3 & 2 \\ -1 & -1 & -1 \end{bmatrix} \quad (ii)$$

$$(C + B) + A = C + (A + B) \quad \text{From (i) and (ii)}$$

$$5 \text{ (vi)} \quad 2A + B = A + (A + B)$$

$$\text{L.H.S} = 2A + B$$

$$= 2 \begin{bmatrix} 1 & 2 & 3 \\ 2 & 3 & 1 \\ 1 & -1 & 0 \end{bmatrix} + \begin{bmatrix} 1 & -1 & 1 \\ 2 & -2 & 2 \\ 3 & 1 & 3 \end{bmatrix}$$

MATHEMATICS FOR 9TH CLASS (UNIT # 1)

Pilot Supercare Mathematics 28 Class 9th

$$\begin{aligned}
 &= \begin{bmatrix} 2(1) & 2(2) & 2(3) \\ 2(2) & 2(3) & 2(1) \\ 2(1) & 2(-1) & 2(0) \end{bmatrix} + \begin{bmatrix} 1 & -1 & 1 \\ 2 & 2 & 2 \\ 3 & 1 & 3 \end{bmatrix} \\
 &= \begin{bmatrix} 2 & 4 & 6 \\ 4 & 6 & 2 \\ 2 & -2 & 0 \end{bmatrix} + \begin{bmatrix} 1 & -1 & 1 \\ 2 & 2 & 2 \\ 3 & 1 & 3 \end{bmatrix} \\
 &= \begin{bmatrix} 2+1 & 4-1 & 6+1 \\ 4+2 & 6+2 & 2+2 \\ 2+3 & -2+1 & 0+3 \end{bmatrix} = \begin{bmatrix} 3 & 3 & 7 \\ 6 & 4 & 4 \\ 5 & -1 & 3 \end{bmatrix} \quad (i)
 \end{aligned}$$

R.H.S. = A + (A + B)

$$\begin{aligned}
 &= \begin{bmatrix} 1 & 2 & 3 \\ 2 & 3 & 1 \\ 1 & -1 & 0 \end{bmatrix} + \left(\begin{bmatrix} 1 & 2 & 3 \\ 2 & 3 & 1 \\ 1 & -1 & 0 \end{bmatrix} + \begin{bmatrix} 1 & -1 & 1 \\ 2 & 2 & 2 \\ 3 & 1 & 3 \end{bmatrix} \right) \\
 &= \begin{bmatrix} 1 & 2 & 3 \\ 2 & 3 & 1 \\ 1 & -1 & 0 \end{bmatrix} + \begin{bmatrix} 1+1 & 2-1 & 3+1 \\ 2+2 & 3+2 & 1+2 \\ 1+3 & -1+1 & 0+3 \end{bmatrix} \\
 &= \begin{bmatrix} 1 & 2 & 3 \\ 2 & 3 & 1 \\ 1 & -1 & 0 \end{bmatrix} + \begin{bmatrix} 2 & 1 & 4 \\ 4 & 1 & 3 \\ 4 & 0 & 3 \end{bmatrix} \\
 &= \begin{bmatrix} 1+2 & 2+1 & 3+4 \\ 2+4 & 3+1 & 1+3 \\ 1+4 & -1+0 & 0+3 \end{bmatrix} = \begin{bmatrix} 3 & 3 & 7 \\ 6 & 4 & 4 \\ 5 & -1 & 3 \end{bmatrix} \quad (ii)
 \end{aligned}$$

$2A + B = A + (A + B)$ From (i) and (ii)

§ (vii) (C - B) - A = (C - A) - B

L.H.S. = (C - B) - A

$$\begin{aligned}
 &= \left(\begin{bmatrix} 1 & 0 & 0 \\ 0 & -2 & 3 \\ 1 & 1 & 2 \end{bmatrix} - \begin{bmatrix} 1 & -1 & 1 \\ 2 & -2 & 2 \\ 3 & 1 & 3 \end{bmatrix} \right) - \begin{bmatrix} 1 & 2 & 3 \\ 2 & 3 & 1 \\ 1 & -1 & 0 \end{bmatrix} \\
 &= \begin{bmatrix} -1-1 & 0+1 & 0-1 \\ 0-2 & -2+2 & 3-2 \\ 1-3 & 1-1 & 2-3 \end{bmatrix} - \begin{bmatrix} 1 & 2 & 3 \\ 2 & 3 & 1 \\ 1 & -1 & 0 \end{bmatrix}
 \end{aligned}$$

MATHEMATICS FOR 9TH CLASS (UNIT # 1)

Pilot Superone Mathematics 29 Class 9th

$$= \begin{bmatrix} 2 & 1 & -1 \\ -2 & 0 & 1 \\ -2 & 0 & 1 \end{bmatrix} - \begin{bmatrix} 1 & 2 & 3 \\ 2 & 3 & 1 \\ 1 & -1 & 0 \end{bmatrix}$$

$$= \begin{bmatrix} 2-1 & 1-2 & -1-3 \\ -2-2 & 0-3 & 1-1 \\ -2-1 & 0+1 & 1-0 \end{bmatrix} = \begin{bmatrix} -3 & -1 & -4 \\ -4 & -3 & 0 \\ -3 & 1 & -1 \end{bmatrix} \quad (i)$$

R.H.S = (C - A) - B

$$= \left(\begin{bmatrix} -1 & 0 & 0 \\ 0 & -2 & 3 \\ 1 & 1 & 2 \end{bmatrix} - \begin{bmatrix} 1 & 2 & 3 \\ 2 & 3 & 1 \\ 1 & -1 & 0 \end{bmatrix} \right) - \begin{bmatrix} 1 & -1 & 1 \\ 2 & -2 & 2 \\ 3 & 1 & 3 \end{bmatrix}$$

$$= \begin{bmatrix} -1-1 & 0-2 & 0-3 \\ 0-2 & -2-3 & 3-1 \\ 1-1 & 1+1 & 2-0 \end{bmatrix} - \begin{bmatrix} 1 & -1 & 1 \\ 2 & -2 & 2 \\ 3 & 1 & 3 \end{bmatrix}$$

$$= \begin{bmatrix} -2 & -2 & -3 \\ 2 & -5 & 2 \\ 0 & 2 & 2 \end{bmatrix} - \begin{bmatrix} 1 & -1 & 1 \\ 2 & -2 & 2 \\ 3 & 1 & 3 \end{bmatrix}$$

$$= \begin{bmatrix} -2-1 & -2+1 & -3-1 \\ -2-2 & -5+2 & 2-2 \\ 0-3 & 2-1 & 2-3 \end{bmatrix} = \begin{bmatrix} -3 & -1 & -4 \\ -4 & -3 & 0 \\ -3 & 1 & -1 \end{bmatrix} \quad (ii)$$

(C - B) - A = (C - A) - B From (i) and (ii)

5 (viii) (A + B) + C = A + (B + C)

L.H.S. = (A + B) + C

$$= \left(\begin{bmatrix} 1 & 2 & 3 \\ 2 & 3 & 1 \\ 1 & 1 & 0 \end{bmatrix} + \begin{bmatrix} 1 & -1 & 1 \\ 2 & -2 & 2 \\ 3 & 1 & 3 \end{bmatrix} \right) + \begin{bmatrix} 1 & 0 & 0 \\ 0 & -2 & 3 \\ 1 & 1 & 2 \end{bmatrix}$$

$$= \begin{bmatrix} 1+1 & 2-1 & 3+1 \\ 2+2 & 3-2 & 1+2 \\ 1+3 & -1+1 & 0+3 \end{bmatrix} + \begin{bmatrix} -1 & 0 & 0 \\ 0 & -2 & 3 \\ 1 & 1 & 2 \end{bmatrix}$$

$$= \begin{bmatrix} 2 & 1 & 4 \\ 4 & 1 & 3 \\ 4 & 0 & 3 \end{bmatrix} + \begin{bmatrix} -1 & 0 & 0 \\ 0 & -2 & 3 \\ 1 & 1 & 2 \end{bmatrix}$$

MATHEMATICS FOR 9TH CLASS (UNIT # 1)

Pilot Superone Mathematics 30 Class 9th

$$= \begin{bmatrix} 2+1 & 1+0 & 4+0 \\ 4+0 & 1 & 2 \\ 4+1 & 0+1 & 3+2 \end{bmatrix} = \begin{bmatrix} 1 & 1 & 4 \\ 4 & -1 & 6 \\ 5 & 1 & 5 \end{bmatrix} \quad (i)$$

R.H.S. = $A + (B + C)$

$$= \left(\begin{bmatrix} 1 & 2 & 3 \\ 2 & 3 & 1 \\ 1 & 1 & 0 \end{bmatrix} + \begin{bmatrix} 1 & -1 & 1 \\ 2 & -2 & 2 \\ 3 & 1 & 3 \end{bmatrix} \right) + \begin{bmatrix} -1 & 0 & 0 \\ 0 & 2 & 3 \\ 1 & 1 & 2 \end{bmatrix}$$

$$= \begin{bmatrix} 1 & 2 & 3 \\ 2 & 3 & 1 \\ 1 & -1 & 0 \end{bmatrix} + \begin{bmatrix} 1+1 & -1+0 & 1+0 \\ 2+0 & 2-2 & 2+3 \\ 3+1 & 1+1 & 3+2 \end{bmatrix}$$

$$= \begin{bmatrix} 1 & 2 & 3 \\ 2 & 3 & 1 \\ 1 & -1 & 0 \end{bmatrix} + \begin{bmatrix} 0 & -1 & 1 \\ 2 & 4 & 5 \\ 4 & 2 & 5 \end{bmatrix}$$

$$= \begin{bmatrix} 1+0 & 2-1 & 3+1 \\ 2+2 & 3-4 & 1+5 \\ 1+4 & -1+2 & 0+5 \end{bmatrix} = \begin{bmatrix} 1 & 1 & 4 \\ 4 & -1 & 6 \\ 5 & 1 & 5 \end{bmatrix} \quad (ii)$$

$(A + B) + C = A + (B + C)$ From (i) and (ii)

5 (ix) $A + (B - C) = (A - C) + B$

L.H.S. = $A + (B - C)$

$$= \begin{bmatrix} 1 & 2 & 3 \\ 2 & 3 & 1 \\ 1 & -1 & 0 \end{bmatrix} + \left(\begin{bmatrix} 1 & -1 & 1 \\ 2 & -2 & 2 \\ 3 & 1 & 3 \end{bmatrix} - \begin{bmatrix} -1 & 0 & 0 \\ 0 & 2 & 3 \\ 1 & 1 & 2 \end{bmatrix} \right)$$

$$= \begin{bmatrix} 1 & 2 & 3 \\ 2 & 3 & 1 \\ 1 & -1 & 0 \end{bmatrix} + \begin{bmatrix} 1+1 & -1-0 & 1-0 \\ 2-0 & -2-2 & 2-3 \\ 3-1 & 1-1 & 3-2 \end{bmatrix}$$

$$= \begin{bmatrix} 1 & 2 & 3 \\ 2 & 3 & 1 \\ 1 & -1 & 0 \end{bmatrix} + \begin{bmatrix} 2 & -1 & 1 \\ 2 & 0 & -1 \\ 2 & 0 & 1 \end{bmatrix}$$

$$= \begin{bmatrix} 1+2 & 2-1 & 3+1 \\ 2+2 & 3+0 & 1+1 \\ 1+2 & -1+0 & 0+1 \end{bmatrix} = \begin{bmatrix} 3 & 1 & 4 \\ 4 & 3 & 2 \\ 3 & -1 & 1 \end{bmatrix} \quad (i)$$

R.H.S. = $(A - C) + B$

MATHEMATICS FOR 9TH CLASS (UNIT # 1)**Pilot Superone Mathematics****31****Class 9th**

$$\begin{aligned}
 &= \left(\begin{bmatrix} 1 & 2 & 3 \\ 2 & 3 & 1 \\ 1 & -1 & 0 \end{bmatrix} - \begin{bmatrix} -1 & 0 & 0 \\ 0 & -2 & 3 \\ 1 & 1 & 2 \end{bmatrix} \right) + \begin{bmatrix} 1 & 1 & 1 \\ 2 & -2 & 2 \\ 3 & 1 & 3 \end{bmatrix} \\
 &= \begin{bmatrix} 1+1 & 2-0 & 3-0 \\ 2-0 & 3+2 & 1-3 \\ 1-1 & -1-1 & 0-2 \end{bmatrix} + \begin{bmatrix} 1 & -1 & 1 \\ 2 & -2 & 2 \\ 3 & 1 & 3 \end{bmatrix} \\
 &= \begin{bmatrix} 2 & 2 & 3 \\ 2 & 5 & -2 \\ 0 & -2 & -2 \end{bmatrix} + \begin{bmatrix} 1 & -1 & 1 \\ 2 & -2 & 2 \\ 3 & 1 & 3 \end{bmatrix} \\
 &= \begin{bmatrix} 2+1 & 2-1 & 3+1 \\ 2+2 & 5-2 & -2+2 \\ 0+3 & -2+1 & -2+3 \end{bmatrix} = \begin{bmatrix} 3 & 1 & 4 \\ 4 & 3 & 0 \\ 3 & 1 & 1 \end{bmatrix} \quad (ii)
 \end{aligned}$$

 $A + (B - C) = (A - C) + B$ From (i) and (ii)**5 (x) $2A + 2B = 2(A + B)$** **L.H.S. = $2A + 2B$**

$$\begin{aligned}
 &= 2 \begin{bmatrix} 1 & 2 & 3 \\ 2 & 3 & 1 \\ 1 & 1 & 0 \end{bmatrix} + 2 \begin{bmatrix} 1 & -1 & 1 \\ 2 & -2 & 2 \\ 3 & 1 & 3 \end{bmatrix} \\
 &= \begin{bmatrix} 2(1) & 2(2) & 2(3) \\ 2(2) & 2(3) & 2(1) \\ 2(1) & 2(-1) & 2(0) \end{bmatrix} + \begin{bmatrix} 2(1) & 2(-1) & 2(1) \\ 2(2) & 2(-2) & 2(2) \\ 2(3) & 2(1) & 2(3) \end{bmatrix} \\
 &= \begin{bmatrix} 2 & 4 & 6 \\ 4 & 6 & 2 \\ 2 & 2 & 0 \end{bmatrix} + \begin{bmatrix} 2 & -2 & 2 \\ 4 & -4 & 4 \\ 6 & 2 & 6 \end{bmatrix} \\
 &= \begin{bmatrix} 2+2 & 4-2 & 6+2 \\ 4+4 & 6-4 & 2+4 \\ 2+6 & -2+2 & 0+6 \end{bmatrix} = \begin{bmatrix} 4 & 2 & 8 \\ 8 & 2 & 6 \\ 8 & 0 & 6 \end{bmatrix} \quad (i)
 \end{aligned}$$

R.H.S. = $2(A + B)$

$$= 2 \left(\begin{bmatrix} 1 & 2 & 3 \\ 2 & 3 & 1 \\ 1 & 1 & 0 \end{bmatrix} + \begin{bmatrix} 1 & -1 & 1 \\ 2 & -2 & 2 \\ 3 & 1 & 3 \end{bmatrix} \right)$$

MATHEMATICS FOR 9TH CLASS (UNIT # 1)

Pilot Superone Mathematics 32 Class 9th

$$= 2 \begin{bmatrix} 1+1 & 2-1 & 3+1 \\ 2+2 & 3-2 & 1+2 \\ 1+3 & -1+1 & 0+3 \end{bmatrix}$$

$$= 2 \begin{bmatrix} 2 & 1 & 4 \\ 4 & 1 & 3 \\ 4 & 0 & 3 \end{bmatrix}$$

$$= \begin{bmatrix} 2(2) & 2(1) & 2(4) \\ 2(4) & 2(1) & 2(3) \\ 2(4) & 2(0) & 2(3) \end{bmatrix} = \begin{bmatrix} 4 & 2 & 8 \\ 8 & 2 & 6 \\ 8 & 0 & 6 \end{bmatrix} \quad (ii)$$

$2A + 2B = 2(A + B)$ From (i) and (ii)

6. If $A = \begin{bmatrix} 1 & -2 \\ 3 & 4 \end{bmatrix}$, $B = \begin{bmatrix} 0 & 7 \\ -3 & 8 \end{bmatrix}$ then find :

(i) $3A - 2B$ (ii) $2A^t - 3B^t$

6 (i) $3A - 2B$

$$= 3 \begin{bmatrix} 1 & 2 \\ 3 & 4 \end{bmatrix} - 2 \begin{bmatrix} 0 & 7 \\ -3 & 8 \end{bmatrix} \quad \text{Putting values of A and B}$$

$$= \begin{bmatrix} 3(1) & 3(-2) \\ 3(3) & 3(4) \end{bmatrix} - \begin{bmatrix} 2(0) & 2(7) \\ 2(-3) & 2(8) \end{bmatrix}$$

$$= \begin{bmatrix} 3 & -6 \\ 9 & 12 \end{bmatrix} - \begin{bmatrix} 0 & 14 \\ -6 & 16 \end{bmatrix}$$

$$= \begin{bmatrix} 3-0 & -6-14 \\ 9+6 & 12-16 \end{bmatrix} = \begin{bmatrix} 3 & 20 \\ 15 & 4 \end{bmatrix}$$

6 (ii) $2A^t - 3B^t$

$$A = \begin{bmatrix} 1 & -2 \\ 3 & 4 \end{bmatrix} \quad \text{We find } A^t \text{ and } B^t$$

MATHEMATICS FOR 9TH CLASS (UNIT # 1)

Pilot Superone Mathematics 33 Class 9th

$$A^t = \begin{bmatrix} 1 & 3 \\ -2 & 4 \end{bmatrix}$$

$$B = \begin{bmatrix} 0 & 7 \\ -3 & 8 \end{bmatrix}$$

$$B^t = \begin{bmatrix} 0 & -3 \\ 7 & 8 \end{bmatrix}$$

$$2A^t - 3B^t$$

$$= 2 \begin{bmatrix} 1 & 3 \\ -2 & 4 \end{bmatrix} - 3 \begin{bmatrix} 0 & -3 \\ 7 & 8 \end{bmatrix} \quad (\text{Putting values } A^t \text{ and } B^t)$$

$$= \begin{bmatrix} 2(1) & 2(3) \\ 2(-2) & 2(4) \end{bmatrix} - \begin{bmatrix} 3(0) & 3(-3) \\ 3(7) & 3(8) \end{bmatrix}$$

$$= \begin{bmatrix} 2 & 6 \\ -4 & 8 \end{bmatrix} - \begin{bmatrix} 0 & -9 \\ 21 & 24 \end{bmatrix}$$

$$= \begin{bmatrix} 2-0 & 6+9 \\ -4-21 & 8-24 \end{bmatrix} = \begin{bmatrix} 2 & 15 \\ -25 & -16 \end{bmatrix}$$

7. If $2 \begin{bmatrix} 2 & 4 \\ -3 & a \end{bmatrix} + 3 \begin{bmatrix} 1 & b \\ 8 & -4 \end{bmatrix} = \begin{bmatrix} 7 & 10 \\ 18 & 1 \end{bmatrix}$, then

find a and b .

$$\begin{bmatrix} 7 & 10 \\ 18 & 1 \end{bmatrix} = 2 \begin{bmatrix} 2 & 4 \\ 3 & a \end{bmatrix} + 3 \begin{bmatrix} 1 & b \\ 8 & -4 \end{bmatrix}$$

$$= \begin{bmatrix} 2(2) & 2(4) \\ 2(-3) & 2a \end{bmatrix} + \begin{bmatrix} 3(1) & 3b \\ 3(8) & 3(-4) \end{bmatrix}$$

MATHEMATICS FOR 9TH CLASS (UNIT # 1)

Pilot Superone Mathematics

34

Class 9th

$$= \begin{bmatrix} 4 & 8 \\ -6 & 2a \end{bmatrix} + \begin{bmatrix} 3 & 3b \\ 24 & -12 \end{bmatrix}$$

$$\begin{bmatrix} 7 & 10 \\ 18 & 1 \end{bmatrix} = \begin{bmatrix} 4+3 & 8+3b \\ -6+24 & 2a-12 \end{bmatrix}$$

$$\begin{bmatrix} 7 & 10 \\ 18 & 1 \end{bmatrix} = \begin{bmatrix} 7 & 8+3b \\ 18 & 2a-12 \end{bmatrix}$$

Therefore $8 + 3b = 10$ (Corresponding elements are same)

$$3b = 2$$

$$b = \frac{2}{3}$$

and

$$1 = 2a - 12$$

$$13 = 2a$$

$$a = \frac{13}{2}$$

8. If $A = \begin{bmatrix} 1 & 2 \\ 0 & 1 \end{bmatrix}$, $B = \begin{bmatrix} 1 & 1 \\ 2 & 0 \end{bmatrix}$ then verify that

(i) $(A + B)^t = A^t + B^t$

(ii) $(A - B)^t = A^t - B^t$

(iii) $A + A^t$ is symmetric

(iv) $A - A^t$ is skew symmetric

(v) $B + B^t$ is symmetric

(vi) $B - B^t$ is skew symmetric

8 (i) $A = \begin{bmatrix} 1 & 2 \\ 0 & 1 \end{bmatrix}$, $B = \begin{bmatrix} 1 & 1 \\ 2 & 0 \end{bmatrix}$

Solution.

$$(A + B)^t = A^t + B^t$$

$$(A + B)^t = ? \quad (\text{Taking L.H.S})$$

$$A + B = ? \quad (\text{Taking L.H.S})$$

MATHEMATICS FOR 9TH CLASS (UNIT # 1)

Pilot Superior Mathematics

35

Class 9th

$$A + B = \begin{bmatrix} 1 & 2 \\ 0 & 1 \end{bmatrix} + \begin{bmatrix} 1 & 1 \\ 2 & 0 \end{bmatrix}$$

$$= \begin{bmatrix} 1+1 & 2+1 \\ 0+2 & 1+0 \end{bmatrix}$$

$$A + B = \begin{bmatrix} 2 & 3 \\ 2 & 1 \end{bmatrix}$$

$$(A + B)^t = \begin{bmatrix} 2 & 2 \\ 3 & 1 \end{bmatrix} \quad (i)$$

$$A^t + B^t = ? \quad (\text{Taking R.H.S})$$

$$A^t = \begin{bmatrix} 1 & 0 \\ 2 & 1 \end{bmatrix} \text{ and}$$

$$B^t = \begin{bmatrix} 1 & 2 \\ 1 & 0 \end{bmatrix}$$

$$A^t + B^t = \begin{bmatrix} 1 & 0 \\ 2 & 1 \end{bmatrix} + \begin{bmatrix} 1 & 2 \\ 1 & 0 \end{bmatrix} = \begin{bmatrix} 1+1 & 0+2 \\ 2+1 & 1+0 \end{bmatrix} = \begin{bmatrix} 2 & 2 \\ 3 & 1 \end{bmatrix} \quad (ii)$$

$$(A + B)^t = A^t + B^t \quad \text{From (i) and (ii)}$$

8 (ii) If $A = \begin{bmatrix} 1 & 2 \\ 0 & 1 \end{bmatrix}$, $B = \begin{bmatrix} 1 & 1 \\ 2 & 0 \end{bmatrix}$

$$\text{Prove that } (A - B)^t = A^t - B^t$$

$$\text{L.H.S} = A - B = \begin{bmatrix} 1 & 2 \\ 0 & 1 \end{bmatrix} - \begin{bmatrix} 1 & 1 \\ 2 & 0 \end{bmatrix}$$

MATHEMATICS FOR 9TH CLASS (UNIT # 1)

Pilot Superone Mathematics

36

Class 9th

$$= \begin{bmatrix} 1-1 & 2-1 \\ 0-2 & 1-0 \end{bmatrix} = \begin{bmatrix} 0 & 1 \\ -2 & 1 \end{bmatrix}$$

$$(A - B)^t = \begin{bmatrix} 0 & -2 \\ 1 & 1 \end{bmatrix}$$

$$\text{R.H.S } A = \begin{bmatrix} 1 & 2 \\ 0 & 1 \end{bmatrix}$$

$$A^t = \begin{bmatrix} 1 & 0 \\ 2 & 1 \end{bmatrix}, B = \begin{bmatrix} 1 & 1 \\ 2 & 0 \end{bmatrix}, B^t = \begin{bmatrix} 1 & 2 \\ 1 & 0 \end{bmatrix}$$

$$A^t - B^t = \begin{bmatrix} 1 & 0 \\ 2 & 1 \end{bmatrix} - \begin{bmatrix} 1 & 2 \\ 1 & 0 \end{bmatrix}$$

$$= \begin{bmatrix} 1-1 & 0-2 \\ 2-1 & 1-0 \end{bmatrix} = \begin{bmatrix} 0 & -2 \\ 1 & 1 \end{bmatrix} \quad (\text{ii})$$

$$(A - B)^t = A^t - B^t \quad \text{From (i) and (ii)}$$

$$\text{8 (iii) If } A = \begin{bmatrix} 1 & 2 \\ 0 & 1 \end{bmatrix}$$

Show $A + A^t$ is a symmetric matrix.

$$A^t = \begin{bmatrix} 1 & 0 \\ 2 & 1 \end{bmatrix}$$

$$A + A^t = \begin{bmatrix} 1 & 2 \\ 0 & 1 \end{bmatrix} + \begin{bmatrix} 1 & 0 \\ 2 & 1 \end{bmatrix}$$

MATHEMATICS FOR 9TH CLASS (UNIT # 1)

Pilot Superone Mathematics

37

Class 9th

$$= \begin{bmatrix} 1+1 & 2+0 \\ 0+2 & 1+1 \end{bmatrix} = \begin{bmatrix} 2 & 2 \\ 2 & 2 \end{bmatrix}$$

$$(A + A^t)^t = \begin{bmatrix} 2 & 2 \\ 2 & 2 \end{bmatrix}$$

So $A + A^t$ is a symmetric matrix.

Note: If $M^t = M$ then M is a symmetric matrix.

8 (iv) If $A = \begin{bmatrix} 1 & 2 \\ 0 & 1 \end{bmatrix}$

Show that $A - A^t$ is symmetric matrix.

$$A^t = \begin{bmatrix} 1 & 0 \\ 2 & 1 \end{bmatrix}$$

$$\begin{aligned} A - A^t &= \begin{bmatrix} 1 & 2 \\ 0 & 1 \end{bmatrix} - \begin{bmatrix} 1 & 0 \\ 2 & 1 \end{bmatrix} \\ &= \begin{bmatrix} 1-1 & 2-0 \\ 0-2 & 1-1 \end{bmatrix} = \begin{bmatrix} 0 & 2 \\ -2 & 0 \end{bmatrix} \end{aligned}$$

$$\text{Now } (A - A^t)^t = \begin{bmatrix} 0 & -2 \\ 2 & 0 \end{bmatrix} \neq A - A^t$$

$A - A^t$ is Not a symmetric matrix

MATHEMATICS FOR 9TH CLASS (UNIT # 1)

=====

Pilot Superone Mathematics 38 Class 9th

8 (v) If $B = \begin{bmatrix} 1 & 1 \\ 2 & 0 \end{bmatrix}$

Show $B + B^t$ is a symmetric matrix.

$$B^t = \begin{bmatrix} 1 & 2 \\ 1 & 0 \end{bmatrix}$$

$$\begin{aligned} B + B^t &= \begin{bmatrix} 1 & 1 \\ 2 & 0 \end{bmatrix} + \begin{bmatrix} 1 & 2 \\ 1 & 0 \end{bmatrix} \\ &= \begin{bmatrix} 1+1 & 1+2 \\ 2+1 & 0+0 \end{bmatrix} = \begin{bmatrix} 2 & 3 \\ 3 & 0 \end{bmatrix} \end{aligned}$$

$$(B + B^t)^t = \begin{bmatrix} 2 & 3 \\ 3 & 0 \end{bmatrix}$$

Now $B + B^t = (B + B^t)^t$

$\therefore B + B^t$ is a symmetric matrix.

8 (vi) $B = \begin{bmatrix} 1 & 1 \\ 2 & 0 \end{bmatrix}$

$$B^t = \begin{bmatrix} 1 & 2 \\ 1 & 0 \end{bmatrix}$$

$$\begin{aligned} B - B^t &= \begin{bmatrix} 1 & 1 \\ 2 & 0 \end{bmatrix} - \begin{bmatrix} 1 & 2 \\ 1 & 0 \end{bmatrix} \\ &= \begin{bmatrix} 1-1 & 1-2 \\ 2-1 & 0-0 \end{bmatrix} \end{aligned}$$

MATHEMATICS FOR 9TH CLASS (UNIT # 1)

Pilot Superone Mathematics

39

Class 9th

$$= \begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix}$$

$$(B - B^t)^t = \begin{bmatrix} 0 & 1 \\ -1 & 0 \end{bmatrix} \neq B - B^t$$

$B - B^t$ is Not a symmetric matrix

Multiplication of Matrices

- (i) Two matrices are conformable for multiplication if the number of columns of the first matrix is equal to the number of rows of the second matrix
- (ii) **Association Law of multiplication**
 If A, B and C are three matrices conformable for multiplication then associative law of multiplication is
 $(AB)C = A(BC)$
- (iii) **Distribution Laws of Multiplication over Addition and Subtraction.**
 - i. $A(B + C) = AB + AC$
 - ii. $(A + B)C = AC + BC$
 - iii. $A(B - C) = AB - AC$
 - iv. $(A - B)C = AC - BC$
- (iv) **Multiplication Identity of a matrix.** If A is a matrix and B another matrix. B is called the identity matrix of A under multiplication if:
 $AB = A = BA$
- (v) **Singular and Non - singular Matrix**

A square matrix A called singular

if $|A| = 0$ and if $|A| \neq 0$ then

A is called non - singular matrix

MATHEMATICS FOR 9TH CLASS (UNIT # 1)

=====

Pilot Superong Mathematics 40 Class 9th

(vi) Adjoint of a Matrix

Adjoint of a square matrix is obtained by interchanging the diagonal entries and changing the sign of other entries. Adjoint of matrix A is denoted as Adj A

(vii) Multiplication inverse of a non - singular square matrix if A and B will multiplication of each other if:

$$AB = BA = I$$

Multiplication inverse of A is written as A^{-1}

(viii) Inverse of matrix using Adjoint.

Let M be a square matrix

Then $M^{-1} = \frac{\text{Adj } M}{|M|}$

www.downloadclassnotes.com

Exercise 1.4

1. Which of the following product of matrices is conformable for multiplication?

(i) $\begin{bmatrix} 1 & -1 \\ 0 & 2 \end{bmatrix} \begin{bmatrix} -2 \\ 3 \end{bmatrix}$ (ii) $\begin{bmatrix} 1 & -1 \\ 1 & 0 \end{bmatrix} \begin{bmatrix} 2 & 1 \\ 1 & 3 \end{bmatrix}$

(iii) $\begin{bmatrix} 1 \\ -1 \end{bmatrix} \begin{bmatrix} 0 & 1 \\ -1 & 2 \end{bmatrix}$ (iv) $\begin{bmatrix} 1 & 2 \\ 0 & -1 \\ -1 & -2 \end{bmatrix} \begin{bmatrix} 1 & 0 & -1 \\ 0 & 1 & 2 \end{bmatrix}$

(v) $\begin{bmatrix} 3 & 2 & 1 \\ 0 & 1 & -1 \end{bmatrix} \begin{bmatrix} 1 & -1 \\ 0 & 2 \\ 2 & 3 \end{bmatrix}$

1 (i) Let $A = \begin{bmatrix} 1 & -1 \\ 0 & 2 \end{bmatrix}$, $B = \begin{bmatrix} 2 \\ 3 \end{bmatrix}$

Number of columns of A = 2

Number of rows of B = 2

Multiplication is possible.

1 (ii) Let $A = \begin{bmatrix} 1 & -1 \\ 1 & 0 \end{bmatrix}$, $B = \begin{bmatrix} 2 & -1 \\ 1 & 3 \end{bmatrix}$

Number of columns in A = 2

Number of rows in B = 2

Multiplication is possible.

1 (iii) Let $A = \begin{bmatrix} 1 \\ -1 \end{bmatrix}$, $B = \begin{bmatrix} 0 & 1 \\ -1 & 2 \end{bmatrix}$

Number of columns in A = 1

Number of rows in B = 2

Multiplication is NOT possible.

MATHEMATICS FOR 9TH CLASS (UNIT # 1)

Pilot Superone Mathematics

42

Class 9th

1 (iv) Let $A = \begin{bmatrix} 1 & 2 \\ 0 & 1 \\ -1 & 2 \end{bmatrix}$, $B = \begin{bmatrix} 1 & 0 & -1 \\ 0 & 1 & 2 \end{bmatrix}$

Number of columns in A = 2

Number of rows in B = 2

Multiplication is possible.

1 (v) Let $A = \begin{bmatrix} 3 & 2 & 1 \\ 0 & 1 & -1 \end{bmatrix}$, $B = \begin{bmatrix} 1 & -1 \\ 0 & 2 \\ -2 & 3 \end{bmatrix}$

Number of columns in A = 3

Number of rows in B = 3

Multiplication AB is possible.

2. If $A = \begin{bmatrix} 3 & 0 \\ 1 & 2 \end{bmatrix}$, $B = \begin{bmatrix} 6 \\ 5 \end{bmatrix}$ find (i) AB (ii) BA (if possible)

2(i) $AB = \begin{bmatrix} 3 & 0 \\ -1 & 2 \end{bmatrix} \begin{bmatrix} 6 \\ 5 \end{bmatrix}$
 $= \begin{bmatrix} 1(6) & +0(5) \\ -1(6) & +2(5) \end{bmatrix} = \begin{bmatrix} 18+0 \\ -6+10 \end{bmatrix} = \begin{bmatrix} 18 \\ 4 \end{bmatrix}$

2(ii) BA is NOT possible

3. Find the following products.

(i) $\begin{bmatrix} 1 & 2 \end{bmatrix} \begin{bmatrix} 4 \\ 0 \end{bmatrix}$ (ii) $\begin{bmatrix} 1 & 2 \end{bmatrix} \begin{bmatrix} 5 \\ 4 \end{bmatrix}$

(iii) $\begin{bmatrix} 3 & 0 \end{bmatrix} \begin{bmatrix} 4 \\ 0 \end{bmatrix}$ (iv) $\begin{bmatrix} 6 & -0 \end{bmatrix} \begin{bmatrix} 4 \\ 0 \end{bmatrix}$

(v) $\begin{bmatrix} 1 & 2 \\ -3 & 0 \\ 6 & 1 \end{bmatrix} \begin{bmatrix} 4 & 5 \\ 0 & -4 \end{bmatrix}$

MATHEMATICS FOR 9TH CLASS (UNIT # 1)**Pt. of Superone Mathematics 43****Class 9th****Solution:**

$$\begin{aligned} 3 \text{ (i)} \quad & [1 \quad 2] \begin{bmatrix} 4 \\ 0 \end{bmatrix} \\ &= [1(4) + 2(0)] = [4 + 0] = [4] \end{aligned}$$

$$\begin{aligned} 3 \text{ (ii)} \quad & [1 \quad 2] \begin{bmatrix} 5 \\ -4 \end{bmatrix} \\ &= [1(5) + 2(-4)] = [5 - 8] = [-3] \end{aligned}$$

$$\begin{aligned} 3 \text{ (iii)} \quad & [-3 \quad 0] \begin{bmatrix} 4 \\ 0 \end{bmatrix} \\ &= [-3(4) + 0(0)] = [-12 + 0] = [-12] \end{aligned}$$

$$\begin{aligned} 3 \text{ (iv)} \quad & [6 \quad 0] \begin{bmatrix} 4 \\ 0 \end{bmatrix} \\ &AB = [6(4) + 0(0)] = [24 + 0] = [24] \end{aligned}$$

$$\begin{aligned} 3 \text{ (v)} \quad & \begin{bmatrix} 1 & 2 \\ -3 & 0 \\ 6 & -1 \end{bmatrix} \begin{bmatrix} 4 & 5 \\ 0 & -4 \end{bmatrix} \\ &= \begin{bmatrix} 1(4)+2(0) & 1(5)+2(-4) \\ -3(4)+0(0) & -3(5)+0(-4) \\ 6(4)+(-1)(0) & 6(5)+(-1)(-4) \end{bmatrix} \\ &= \begin{bmatrix} 4+0 & 5-8 \\ 12+0 & -15+0 \\ 24+0 & 30+4 \end{bmatrix} = \begin{bmatrix} 4 & -3 \\ 12 & -15 \\ 24 & 34 \end{bmatrix} \end{aligned}$$

4. Multiply the following matrices.

$$(a) \begin{bmatrix} 2 & 3 \\ 1 & 1 \\ 0 & -2 \end{bmatrix} \begin{bmatrix} 2 & 1 \\ 3 & 0 \end{bmatrix} \quad (b) \begin{bmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \end{bmatrix} \begin{bmatrix} 1 & 2 \\ 3 & 4 \\ -1 & 1 \end{bmatrix}$$

MATHEMATICS FOR 9TH CLASS (UNIT # 1)

Pilot Superone Mathematics 44 Class 9th

$$(c) \begin{bmatrix} 1 & 2 \\ 3 & 4 \\ -1 & 1 \end{bmatrix} \begin{bmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \end{bmatrix} \quad (d) \begin{bmatrix} 8 & 5 \\ 6 & 4 \end{bmatrix} \begin{bmatrix} 2 & -\frac{5}{2} \\ -4 & 4 \end{bmatrix}$$

$$(e) \begin{bmatrix} -1 & 2 \\ 1 & 3 \end{bmatrix} \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix}$$

4 (a) Solution:

$$\begin{aligned} & \begin{bmatrix} 2 & 3 \\ 1 & 1 \\ 0 & 2 \end{bmatrix} \begin{bmatrix} 2 & -1 \\ 3 & 0 \end{bmatrix} \\ &= \begin{bmatrix} 2(2)+3(3) & 2(-1)+3(0) \\ 1(2)+1(3) & 1(-1)+1(0) \\ 0(2)+(-2)(3) & 0(-1)+(-2)(0) \end{bmatrix} \\ &= \begin{bmatrix} 4+9 & -2+0 \\ 2+3 & -1+0 \\ 0 & 6 \end{bmatrix} = \begin{bmatrix} 13 & -2 \\ 5 & -1 \\ -6 & 0 \end{bmatrix} \end{aligned}$$

$$\begin{aligned} 4(b) \quad & \begin{bmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \end{bmatrix} \begin{bmatrix} 1 & 2 \\ 3 & 4 \\ -1 & 1 \end{bmatrix} \\ &= \begin{bmatrix} 1(1)+2(3)+3(-1) & 1(2)+2(4)+3(1) \\ 4(1)+5(3)+6(-1) & 4(2)+5(4)+6(1) \end{bmatrix} \\ &= \begin{bmatrix} 1+6-3 & 2+8+3 \\ 4+15-6 & 8+20+6 \end{bmatrix} \\ &= \begin{bmatrix} 4 & 13 \\ 13 & 34 \end{bmatrix} \end{aligned}$$

$$4(c) \quad \begin{bmatrix} 1 & 2 \\ 3 & 4 \\ -1 & 1 \end{bmatrix} \begin{bmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \end{bmatrix}$$

MATHEMATICS FOR 9TH CLASS (UNIT # 1)

Pilot Superone Mathematics 45 Class 9th

$$= \begin{bmatrix} 1(1) + 2(4) & 1(2) + 2(5) & 1(3) + 2(6) \\ 3(1) + 4(4) & 3(2) + 4(5) & 3(3) + 4(6) \\ 1(1) + 1(4) & -1(2) + 1(5) & -1(3) + 1(6) \end{bmatrix}$$

$$= \begin{bmatrix} 1+8 & 2+10 & 3+12 \\ 3+16 & 6+20 & 9+24 \\ -1+4 & -2+5 & -3+6 \end{bmatrix}$$

$$= \begin{bmatrix} 9 & 12 & 15 \\ 19 & 26 & 33 \\ 3 & 3 & 3 \end{bmatrix}$$

$$4(d) \quad \begin{bmatrix} 8 & 5 \\ 6 & 4 \end{bmatrix} \begin{bmatrix} 2 & -\frac{5}{2} \\ -4 & 4 \end{bmatrix}$$

$$= \begin{bmatrix} 8(2) + 5(-4) & 8\left(-\frac{5}{2}\right) + 5(4) \\ 6(2) + 4(-4) & 6\left(-\frac{5}{2}\right) + 4(4) \end{bmatrix}$$

$$= \begin{bmatrix} 16 & 20 & -20+20 \\ 12-16 & -15+16 \end{bmatrix}$$

$$= \begin{bmatrix} -4 & 0 \\ -4 & 1 \end{bmatrix}$$

$$4(e) \quad \begin{bmatrix} -1 & 2 \\ 1 & 3 \end{bmatrix} \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix}$$

$$= \begin{bmatrix} -1(0) + 2(0) & -1(0) + 2(0) \\ 1(0) + 3(0) & 1(0) + 3(0) \end{bmatrix}$$

$$= \begin{bmatrix} 0+0 & 0+0 \\ 0+0 & 0+0 \end{bmatrix} = \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix}$$

MATHEMATICS FOR 9TH CLASS (UNIT # 1)

Pilot Superone Mathematics 46 Class 9th

5. Let $A = \begin{bmatrix} 1 & 3 \\ 2 & 0 \end{bmatrix}$, $B = \begin{bmatrix} 1 & 2 \\ -3 & -5 \end{bmatrix}$ and $C = \begin{bmatrix} 2 & 1 \\ 1 & 3 \end{bmatrix}$

verify whether

- (i) $AB = BA$ (ii) $A(BC) = (AB)C$
 (iii) $A(B+C) = AB+AC$ (iv) $A(B-C) = AB-AC$

5(i) $A = \begin{bmatrix} 1 & 3 \\ 2 & 0 \end{bmatrix}$, $B = \begin{bmatrix} 1 & 2 \\ -3 & -5 \end{bmatrix}$

$$AB = \begin{bmatrix} -1(1) + 3(-3) & -1(2) + 3(-5) \\ 2(1) + 0(-3) & 2(2) + 0(-5) \end{bmatrix}$$

$$= \begin{bmatrix} -1-9 & -2-15 \\ 2+0 & 4+0 \end{bmatrix} = \begin{bmatrix} -10 & -17 \\ 2 & 4 \end{bmatrix} \quad (i)$$

and $BA = \begin{bmatrix} 1 & 2 \\ -3 & -5 \end{bmatrix} \begin{bmatrix} -1 & 3 \\ 2 & 0 \end{bmatrix}$

$$= \begin{bmatrix} 1(-1) + 2(2) & 1(3) + 2(0) \\ -3(-1) + (-5)(2) & -3(3) + (-5)(0) \end{bmatrix}$$

$$= \begin{bmatrix} -1+4 & 3+0 \\ 3-10 & -9+0 \end{bmatrix} = \begin{bmatrix} 3 & 3 \\ 7 & 9 \end{bmatrix} \quad (ii)$$

$AB \neq BA$ From (i) and (ii)

- 5(ii) Prove that $A(BC) = (AB)C$

$A = \begin{bmatrix} -1 & 3 \\ 2 & 0 \end{bmatrix}$, $B = \begin{bmatrix} 1 & 2 \\ -3 & -5 \end{bmatrix}$, $C = \begin{bmatrix} 2 & 1 \\ 1 & 3 \end{bmatrix}$

$A(BC) = ?$

BC (First we find BC)

$$= \begin{bmatrix} 1 & 2 \\ -3 & -5 \end{bmatrix} \begin{bmatrix} 2 & 1 \\ 1 & 3 \end{bmatrix}$$

$$= \begin{bmatrix} 1(2) + 2(1) & 1(1) + 2(3) \\ -3(2) + (-5)(1) & -3(1) + (-5)(3) \end{bmatrix}$$

MATHEMATICS FOR 9TH CLASS (UNIT # 1)

Pilot Superone Mathematics 47

Class 9th

$$BC = \begin{bmatrix} 2+2 & 1+6 \\ -6 & 5 & -3 & 15 \end{bmatrix}$$

$$= \begin{bmatrix} 4 & 7 \\ -11 & -18 \end{bmatrix}$$

$$\text{Now } A(BC) = \begin{bmatrix} 1 & 3 \\ 2 & 0 \end{bmatrix} \begin{bmatrix} 4 & 7 \\ -11 & -18 \end{bmatrix}$$

$$= \begin{bmatrix} 1(4) + 3(-11) & -1(7) + 3(-18) \\ 2(4) + 0(-11) & 2(7) + 0(-18) \end{bmatrix}$$

$$= \begin{bmatrix} 4 - 33 & -7 - 54 \\ 8 + 0 & 14 + 0 \end{bmatrix}$$

$$= \begin{bmatrix} -29 & -61 \\ 8 & 14 \end{bmatrix} \quad (I)$$

$(AB)C = ?$

$$AB = \begin{bmatrix} -1 & 3 \\ 2 & 0 \end{bmatrix} \begin{bmatrix} 1 & 2 \\ 3 & -5 \end{bmatrix} \quad \text{First we find } AB$$

$$= \begin{bmatrix} -1(1) + 3(-3) & -1(2) + 3(-5) \\ 2(1) + 0(-3) & 2(2) + 0(-5) \end{bmatrix}$$

$$= \begin{bmatrix} -1 - 9 & -2 - 15 \\ 2 + 0 & 4 + 0 \end{bmatrix}$$

$$= \begin{bmatrix} -10 & -17 \\ 2 & 4 \end{bmatrix}$$

Now, we find $(AB)C$

$$(AB)C = \begin{bmatrix} -10 & -17 \\ 2 & 4 \end{bmatrix} \begin{bmatrix} 2 & 1 \\ 1 & 3 \end{bmatrix}$$

$$= \begin{bmatrix} -10(2) + (-17)(1) & -10(1) + (-17)(3) \\ 2(2) + 4(1) & 2(1) + 4(3) \end{bmatrix}$$

MATHEMATICS FOR 9TH CLASS (UNIT # 1)

Pilot Superone Mathematics 48 Class 9th

$$= \begin{bmatrix} -20-17 & -10-51 \\ 4+4 & 2+12 \end{bmatrix}$$

$$= \begin{bmatrix} 37 & -61 \\ 8 & 14 \end{bmatrix} \quad (ii)$$

$$A(BC) = (AB)C \quad \text{From (i) and (ii)}$$

5(iii) Prove that $A(B+C) = AB + AC$

$$A(B+C) = ?$$

$(B+C)$. First we find $(B+C)$

$$= \begin{bmatrix} 1 & 2 \\ 3 & -5 \end{bmatrix} + \begin{bmatrix} 2 & 1 \\ 1 & 3 \end{bmatrix}$$

$$= \begin{bmatrix} 1+2 & 2+1 \\ -3+1 & 5+3 \end{bmatrix}$$

$$= \begin{bmatrix} 3 & 3 \\ -2 & -2 \end{bmatrix}$$

$$A(B+C) = \begin{bmatrix} -1 & 3 \\ 2 & 0 \end{bmatrix} \begin{bmatrix} 3 & 3 \\ -2 & -2 \end{bmatrix}$$

$$= \begin{bmatrix} -1(3)+3(-2) & -1(3)+3(-2) \\ 2(3)+0(-2) & 2(3)+0(-2) \end{bmatrix}$$

$$= \begin{bmatrix} -3 & 6 & -3 & -6 \\ 6+0 & 6+0 \end{bmatrix}$$

$$= \begin{bmatrix} -9 & -9 \\ 6 & 6 \end{bmatrix} \quad (i)$$

$$AB + AC = ? \quad \text{Taking L.H.S.}$$

$$AB = \begin{bmatrix} -1 & 3 \\ 2 & 0 \end{bmatrix} \begin{bmatrix} 1 & 2 \\ -3 & -5 \end{bmatrix}$$

$$= \begin{bmatrix} -1(1)+3(-3) & -1(2)+3(-5) \\ 2(1)+0(-3) & 2(2)+0(-5) \end{bmatrix}$$

MATHEMATICS FOR 9TH CLASS (UNIT # 1)

Pilot Superone Mathematics

65

Class 9th

$$A^{-1} = \frac{\text{Adj } A}{|A|}$$

$$= \frac{\begin{bmatrix} 2 & 0 \\ 1 & 4 \end{bmatrix}}{8} = \begin{bmatrix} \frac{2}{8} & 0 \\ \frac{1}{8} & \frac{4}{8} \end{bmatrix}$$

$$= \begin{bmatrix} \frac{1}{4} & 0 \\ \frac{1}{8} & \frac{1}{2} \end{bmatrix}$$

$$B^{-1}A^{-1} = \begin{bmatrix} -1 & 1 \\ -6 & 3 \end{bmatrix} \begin{bmatrix} \frac{1}{4} & 0 \\ \frac{1}{8} & \frac{1}{2} \end{bmatrix}$$

$$= \begin{bmatrix} -\frac{1}{6}(\frac{1}{4}) + \frac{1}{3}(\frac{1}{8}) & \frac{1}{6}(0) + (\frac{1}{3})(\frac{1}{2}) \\ -\frac{1}{6}(\frac{1}{4}) + (-\frac{2}{3})(\frac{1}{8}) & \frac{1}{6}(0) + (-\frac{2}{3})(\frac{1}{2}) \end{bmatrix}$$

$$= \begin{bmatrix} -\frac{1}{24} + \frac{1}{24} & 0 + \frac{1}{6} \\ -\frac{1}{24} - \frac{1}{12} & 0 - \frac{1}{3} \end{bmatrix}$$

$$= \begin{bmatrix} 0 & \frac{1}{6} \\ -\frac{1}{8} & -\frac{1}{3} \end{bmatrix}$$

(ii)

$$(AB)^{-1} = B^{-1}A^{-1}$$

From (i) and (ii)

MATHEMATICS FOR 9TH CLASS (UNIT # 1)

Vol Superone Mathematics 66 Class 9th

6(ii) $A = \begin{bmatrix} 4 & 0 \\ 1 & 2 \end{bmatrix}$, $D = \begin{bmatrix} 3 & 1 \\ 2 & 2 \end{bmatrix}$, verify that:

$(DA)^{-1} = A^{-1}D^{-1}$

$$(DA)^{-1} = \begin{bmatrix} 3 & 1 \\ -2 & 2 \end{bmatrix}^{-1} \begin{bmatrix} 4 & 0 \\ 1 & 2 \end{bmatrix}^{-1}$$

$$= \begin{bmatrix} 3(4) + 1(-1) & 3(0) + 1(2) \\ -2(4) + 2(-1) & -2(0) + 2(2) \end{bmatrix}^{-1}$$

$$= \begin{bmatrix} 12 - 1 & 0 + 2 \\ -8 - 2 & 0 + 4 \end{bmatrix}^{-1}$$

$$= \begin{bmatrix} 11 & 2 \\ -10 & 4 \end{bmatrix}^{-1}$$

$$(DA)^{-1} = \begin{bmatrix} 11 & 2 \\ -10 & 4 \end{bmatrix}^{-1}$$

$$= \frac{(11)(4) - (2)(-10)}{44 - 20} = \frac{64}{24}$$

$$\text{Adj}(DA) = \begin{bmatrix} 4 & 2 \\ 10 & 11 \end{bmatrix}$$

$$(DA)^{-1} = \frac{\text{Adj}(DA)}{DA}$$

$$= \frac{\begin{bmatrix} 4 & 2 \\ 10 & 11 \end{bmatrix}}{64} = \begin{bmatrix} \frac{4}{64} & \frac{2}{64} \\ \frac{10}{64} & \frac{11}{64} \end{bmatrix}$$

MATHEMATICS FOR 9TH CLASS (UNIT # 1)

Pilot Superone Mathematics 67

Class 9th

$$\begin{bmatrix} 1 & -1 \\ 16 & -32 \\ 5 & 11 \\ 32 & 64 \end{bmatrix} \quad (1)$$

$$A = \begin{bmatrix} 1 & 0 \\ -10 & 2 \end{bmatrix}$$

$$|A| = (1)(2) - (-1)(0) = 2 \quad 0 \neq 2$$

$$\text{Adj } A = \begin{bmatrix} 2 & 0 \\ 1 & 1 \end{bmatrix}$$

$$A^{-1} = \frac{\text{Adj } A}{|A|} = \frac{\begin{bmatrix} 2 & 0 \\ 1 & 1 \end{bmatrix}}{2} = \begin{bmatrix} 1 & 0 \\ 0.5 & 0.5 \end{bmatrix}$$

$$\begin{bmatrix} 1 & 0 \\ 1 & 1 \\ 8 & 2 \end{bmatrix}$$

$$D = \begin{bmatrix} 3 & 1 \\ -2 & 2 \end{bmatrix}$$

$$|D| = \begin{vmatrix} 3 & 1 \\ 2 & 2 \end{vmatrix}$$

$$\text{Adj } D = \begin{bmatrix} 2 & -1 \\ 2 & 3 \end{bmatrix}$$

$$D^{-1} = \frac{\text{Adj } D}{|D|} = \frac{\begin{bmatrix} 2 & -1 \\ 2 & 3 \end{bmatrix}}{8}$$

MATHEMATICS FOR 9TH CLASS (UNIT # 1)

Pilot Superone Mathematics

68

Class 9th

$$\begin{aligned}
 &= \begin{bmatrix} 2 & -1 \\ 8 & 8 \\ 2 & 3 \\ 8 & 8 \end{bmatrix} \times \begin{bmatrix} 1 & -1 \\ 4 & 8 \\ 1 & 3 \\ 4 & 8 \end{bmatrix} \\
 A^{-1}D^{-1} &= \begin{bmatrix} 1 & 0 \\ 4 & 0 \\ 1 & 1 \\ 8 & 2 \end{bmatrix} \begin{bmatrix} 1 & -1 \\ 4 & 8 \\ 1 & 3 \\ 4 & 8 \end{bmatrix} \\
 &= \begin{bmatrix} \frac{1}{4}\left(\frac{1}{4}\right) + 0\left(\frac{1}{4}\right) & \frac{1}{4}\left(-\frac{1}{8}\right) + 0\left(\frac{3}{8}\right) \\ \frac{1}{8}\left(\frac{1}{4}\right) + \frac{1}{2}\left(\frac{1}{4}\right) & \frac{1}{8}\left(-\frac{1}{8}\right) + \frac{1}{2}\left(\frac{3}{8}\right) \end{bmatrix} \\
 &= \begin{bmatrix} \frac{1}{16} + 0 & -\frac{1}{32} + 0 \\ \frac{1}{32} + \frac{1}{8} & -\frac{1}{64} + \frac{3}{16} \end{bmatrix} \\
 &= \begin{bmatrix} \frac{1}{16} & -\frac{1}{32} \\ \frac{5}{32} & \frac{11}{64} \end{bmatrix} \quad (ii)
 \end{aligned}$$

$$(DA)^{-1} = A^{-1}D^{-1}$$

From (i), (ii)

SOLUTION OF SIMULTANEOUS LINEAR EQUATIONS

System of two linear equations in two variable in general form are,

$ax + by = m$, $cx + dy = n$ where a, b, c, d, m, n are real numbers.

This system is also called simultaneous linear equations.

Cramer's Rule

Let $ax + by = m$

$cx + dy = n$

writing the equations in matrix form

$$\begin{bmatrix} ax + by \\ cx + dy \end{bmatrix} = \begin{bmatrix} m \\ n \end{bmatrix}$$

MATHEMATICS FOR 9TH CLASS (UNIT # 1)

Pilot Supreme Mathematics

69

Class 9th

$$\text{or} \quad \begin{bmatrix} a & b \\ c & d \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} m \\ n \end{bmatrix}$$

$$AX = B$$

$$\text{When } A = \begin{bmatrix} a & b \\ c & d \end{bmatrix}, X = \begin{bmatrix} x \\ y \end{bmatrix}, B = \begin{bmatrix} m \\ n \end{bmatrix}$$

$$AX = B$$

$$X = A^{-1}B$$

$$X = \frac{\text{Adj } A}{|A|} \times B \quad [A \neq 0]$$

$$\begin{bmatrix} x \\ y \end{bmatrix} = \frac{\begin{bmatrix} d & -b \\ -c & a \end{bmatrix}}{|A|} \begin{bmatrix} m \\ n \end{bmatrix}$$

$$\begin{bmatrix} x \\ y \end{bmatrix} = \frac{\begin{bmatrix} dm - bn \\ -cm + an \end{bmatrix}}{|A|}$$

$$\text{Thus } x = \frac{dm - bn}{|A|} = \frac{|A_x|}{|A|}$$

$$\text{And } y = \frac{an - cm}{|A|} = \frac{|A_y|}{|A|}$$

$$\text{When } |A_x| = \begin{vmatrix} m & b \\ n & d \end{vmatrix}$$

$$\text{and } |A_y| = \begin{vmatrix} a & m \\ c & n \end{vmatrix}$$

MATHEMATICS FOR 9TH CLASS (UNIT # 1)

Past Superior Mathematics 70 Class 9

Exercise 1.6

Q.1. Use matrices if possible to solve the following system of linear equations by

- (i) The matrix inverse method
 (ii) The Cramer's Rule

(i) $2x - 2y = 4$
 $3x + 2y = 6$

(iii) $4x + 2y = 8$
 $3x - y = 1$

(v) $3x - 2y = 4$
 $-6x + 4y = 7$

(vii) $2x - 2y = 4$
 $-5x - 2y = 10$

(ii) $2x + y = 3$
 $6x + 5y = 1$

(iv) $3x - 2y = 6$
 $5x - 2y = -10$

(vi) $4x + y = 9$
 $3x - y = -5$

(viii) $3x - 4y = 4$
 $x + 2y = 8$

Solution:-

1(i) $2x - 2y = 4$
 $3x + 2y = 6$

Writing the equations in matrix form.

$$\begin{bmatrix} 2 & -2 \\ 3 & 2 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 4 \\ 6 \end{bmatrix}$$

or $\begin{bmatrix} 2 & -2 \\ 3 & 2 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 4 \\ 6 \end{bmatrix}$

Let $AX = B$

Where $A = \begin{bmatrix} 2 & -2 \\ 3 & 2 \end{bmatrix}$, $X = \begin{bmatrix} x \\ y \end{bmatrix}$, $B = \begin{bmatrix} 4 \\ 6 \end{bmatrix}$

$$|A| = \begin{vmatrix} 2 & -2 \\ 3 & 2 \end{vmatrix}$$

$$= (2)(2) - (-2)(3)$$

$$4 + 6 = 10 \neq 0$$

A is a non-singular matrix. Its multiplicative inverse is A^{-1}

Now $A^{-1}AX = A^{-1}B$

MATHEMATICS FOR 9TH CLASS (UNIT # 1)

Pick Superon: Mathematics 71 Class 9th

or

$$X = A^{-1}B$$

$$\text{Adj } A = \begin{bmatrix} 2 & 2 \\ 3 & 3 \end{bmatrix} \quad A = 10$$

$$A^{-1} = \frac{\text{Adj } A}{|A|}$$

$$\begin{bmatrix} 2 & 2 \\ 3 & 3 \end{bmatrix}$$

10

$$\begin{bmatrix} 2 & 2 \\ 10 & 10 \end{bmatrix} \quad \begin{bmatrix} 1 & 1 \\ 5 & 5 \end{bmatrix}$$

Now

$$X = A^{-1}B$$

$$\begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 1 & 1 \\ 5 & 5 \end{bmatrix} \begin{bmatrix} 4 \\ 6 \end{bmatrix}$$

$$\begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 1(4) & 1(6) \\ 5(4) & 5(6) \end{bmatrix}$$

$$\begin{bmatrix} 4 & 6 \\ 20 & 30 \end{bmatrix}$$

$$\text{So, } x = 2, y = 0$$

Using Cramer's Rule:

$$2x - 2y = 4$$

$$3x + 2y = 6$$

MATHEMATICS FOR 9TH CLASS (UNIT # 1)

Prior Supreme Mathematics

72

Class 9th

$$A = \begin{bmatrix} 2 & -2 \\ 3 & 2 \end{bmatrix}$$

$$A_x = \begin{bmatrix} 4 & -2 \\ 6 & 2 \end{bmatrix}$$

$$A_y = \begin{bmatrix} 2 & 4 \\ 3 & 6 \end{bmatrix}$$

$$|A| = \begin{vmatrix} 2 & -2 \\ 3 & 2 \end{vmatrix}$$

$$x = \frac{|A_x|}{|A|} = \frac{\begin{vmatrix} 4 & -2 \\ 6 & 2 \end{vmatrix}}{10}$$

$$= \frac{(4)(2) - (-2)(6)}{10}$$

$$= \frac{8 + 12}{10} = 2$$

$$y = \frac{|A_y|}{|A|} = \frac{\begin{vmatrix} 2 & 4 \\ 3 & 6 \end{vmatrix}}{10}$$

$$= \frac{(2)(6) - (3)(4)}{10}$$

$$= \frac{12 - 12}{10} = 0$$

Sol. 1(iii)
$$\begin{aligned} x - 2y &= 0 \\ 2x + y &= 3 \\ 6x + 5y &= 1 \end{aligned}$$

Writing the equations in the matrix form

$$\begin{bmatrix} 2x & +y \\ 6x & +5y \end{bmatrix} = \begin{bmatrix} 3 \\ 1 \end{bmatrix}$$

or
$$\begin{bmatrix} 2 & 1 \\ 6 & 5 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 3 \\ 1 \end{bmatrix}$$

MATHEMATICS FOR 9TH CLASS (UNIT # 1)**Pilot Superone Mathematics 73****Class 9th**Let $AX = B$ Where $A = \begin{bmatrix} 2 & 1 \\ 6 & 5 \end{bmatrix}$, $X = \begin{bmatrix} x \\ y \end{bmatrix}$, $B = \begin{bmatrix} 3 \\ 1 \end{bmatrix}$

$$|A| = \begin{vmatrix} 2 & 1 \\ 6 & 5 \end{vmatrix}$$

$$(2 \times 5) - (1 \times 6) = 10 - 6 = 4$$

A is a non-singular matrix

$$AX = B$$

$$X = A^{-1}B$$

$$\text{Adj } A = \begin{bmatrix} 5 & -1 \\ -6 & 2 \end{bmatrix}$$

$$A^{-1} = \frac{\text{Adj } A}{|A|} = \frac{\begin{bmatrix} 5 & -1 \\ -6 & 2 \end{bmatrix}}{4}$$

$$\begin{bmatrix} 5 & -1 \\ -6 & 2 \end{bmatrix} \div 4 = \begin{bmatrix} \frac{5}{4} & -\frac{1}{4} \\ -\frac{3}{2} & \frac{1}{2} \end{bmatrix}$$

Now

$$X = A^{-1}B$$

$$\begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} \frac{5}{4} & -\frac{1}{4} \\ -\frac{3}{2} & \frac{1}{2} \end{bmatrix} \begin{bmatrix} 3 \\ 1 \end{bmatrix}$$

$$\begin{bmatrix} \frac{5}{4}(3) + \left(-\frac{1}{4}\right)(1) \\ -\frac{3}{2}(3) + \left(\frac{1}{2}\right)(1) \end{bmatrix}$$

MATHEMATICS FOR 9TH CLASS (UNIT # 1)

Pilot Superior Mathematics

74

Class 9th

$$\begin{bmatrix} 15 & 1 \\ 4 & 4 \\ 9 & 1 \\ 2 & 2 \\ 11 & 1 \end{bmatrix}$$

$$\begin{bmatrix} 1 & 1 & 2 \\ 2 & 1 & 1 \\ 3 & 1 & 1 \end{bmatrix}$$

$$x = \frac{7}{2}, y = 4$$

Using Cramer's Rule

$$\begin{aligned} 2x + y &= 3 \\ 6x + 5y &= 1 \end{aligned}$$

$$A = \begin{bmatrix} 2 & 1 \\ 6 & 5 \end{bmatrix}$$

$$A_1 = \begin{bmatrix} 3 & 1 \\ 1 & 5 \end{bmatrix}$$

$$A_2 = \begin{bmatrix} 2 & 3 \\ 6 & 1 \end{bmatrix}$$

$$A = \begin{bmatrix} 2 & 1 \\ 6 & 5 \end{bmatrix}$$

$$(2 \times 5) - (1 \times 6) = 10 - 6 = 4$$

$$x = \frac{|A_1|}{|A|} = \frac{10 - 5}{4}$$

$$\frac{(3 \times 5) - (1 \times 1)}{4} = \frac{15 - 1}{4}$$

$$\frac{14}{4} = \frac{7}{2}$$

www.downloadclassnotes.com

MATHEMATICS FOR 9TH CLASS (UNIT # 1)

Pkx Superone Mathematics

75

Class 9th

$$y = \frac{A_{31}}{|A|} = \frac{\begin{vmatrix} 2 & 3 \\ 6 & 1 \end{vmatrix}}{4} = \frac{(2)(1) - (3)(6)}{4} = \frac{2 - 18}{4} = \frac{-16}{4} = -4$$

So, $x = \frac{7}{2}, y = -4$

1(iii) $4x + 2y = 8$
 $3x - y = -1$

Writing the equations in matrix form

$$\begin{bmatrix} 4 & 2 \\ 3 & -1 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 8 \\ -1 \end{bmatrix}$$

or $\begin{bmatrix} 4 & 2 \\ 3 & -1 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 8 \\ -1 \end{bmatrix}$

Let $A X = B$

where $A = \begin{bmatrix} 4 & 2 \\ 3 & -1 \end{bmatrix}, X = \begin{bmatrix} x \\ y \end{bmatrix}, B = \begin{bmatrix} 8 \\ -1 \end{bmatrix}$

$$|A| = \begin{vmatrix} 4 & 2 \\ 3 & -1 \end{vmatrix} = (4)(-1) - (2)(3) = -4 - 6 = -10$$

$$A X = B$$

$$X = A^{-1} B$$

$$\text{Adj } A = \begin{bmatrix} 1 & 2 \\ 3 & 4 \end{bmatrix}$$

$$A^{-1} = \frac{\text{Adj } A}{|A|} = \frac{\begin{bmatrix} 1 & 2 \\ 3 & 4 \end{bmatrix}}{-10}$$

MATHEMATICS FOR 9TH CLASS (UNIT # 1)

Pilot Superone Mathematics

76

Class 9th

$$= \begin{bmatrix} \frac{-1}{-10} & \frac{-2}{-10} \\ \frac{-3}{-10} & \frac{4}{-10} \end{bmatrix} = \begin{bmatrix} \frac{1}{10} & \frac{1}{5} \\ \frac{3}{10} & \frac{2}{5} \end{bmatrix}$$

$$X = A^{-1}B$$

$$= \begin{bmatrix} \frac{1}{10} & \frac{1}{5} \\ \frac{3}{10} & \frac{2}{5} \end{bmatrix} \begin{bmatrix} 8 \\ -1 \end{bmatrix}$$

$$= \begin{bmatrix} \frac{1}{10}(8) + \frac{1}{5}(-1) \\ \frac{3}{10}(8) + \frac{2}{5}(-1) \end{bmatrix}$$

$$= \begin{bmatrix} \frac{4}{5} - \frac{1}{5} \\ \frac{12}{5} - \frac{2}{5} \end{bmatrix}$$

$$\begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} \frac{3}{5} \\ \frac{10}{5} \end{bmatrix}$$

So,

$$x = \frac{3}{5}$$

$$y = \frac{10}{5}$$

Using Cramer's Rule:

$$4x + 2y = 8$$

$$3x - y = 1$$

$$A = \begin{bmatrix} 4 & 2 \\ 3 & -1 \end{bmatrix}$$

MATHEMATICS FOR 9TH CLASS (UNIT # 1)

Pilot Superone Mathematics

77

Class 9th

$$A_x = \begin{bmatrix} 8 & 2 \\ -1 & -1 \end{bmatrix}$$

$$A_y = \begin{bmatrix} 4 & 8 \\ 3 & -1 \end{bmatrix}$$

$$|A| = \begin{vmatrix} 4 & 2 \\ 3 & -1 \end{vmatrix}$$

$$= (4)(-1) - (2)(3)$$

$$= -4 - 6 = -10$$

$$x = \frac{|A_x|}{|A|} = \frac{\begin{vmatrix} 8 & 2 \\ -1 & -1 \end{vmatrix}}{-10}$$

$$= \frac{(8)(-1) - (2)(-1)}{-10}$$

$$= \frac{-8 + 2}{-10} = \frac{-6}{-10} = \frac{3}{5}$$

$$y = \frac{|A_y|}{|A|} = \frac{\begin{vmatrix} 4 & 8 \\ 3 & -1 \end{vmatrix}}{-10}$$

$$= \frac{(4)(-1) - (8)(3)}{-10} = \frac{-4 - 24}{-10}$$

$$= \frac{28}{-10} = \frac{14}{5}$$

So,

$$x = \frac{3}{5}, y = \frac{14}{5}$$

$$3x - 2y = -6$$

$$5x - 2y = -10$$

Writing the equations in matrix form

$$\begin{bmatrix} 3x & -2y \end{bmatrix} = \begin{bmatrix} -6 \\ -10 \end{bmatrix}$$

MATHEMATICS FOR 9TH CLASS (UNIT # 1)

Pilot Superone Mathematics 78 Class 9th

or $\begin{bmatrix} 3 & -2 \\ 5 & -2 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} -6 \\ -10 \end{bmatrix}$

Let $AX = B$

Where $A = \begin{bmatrix} 3 & -2 \\ 5 & -2 \end{bmatrix}$ $X = \begin{bmatrix} x \\ y \end{bmatrix}$ $B = \begin{bmatrix} -6 \\ -10 \end{bmatrix}$

$$|A| = \begin{vmatrix} 3 & -2 \\ 5 & -2 \end{vmatrix}$$

$$= (3)(-2) - (-2)(5)$$

$$= -6 + 10 = 4$$

$$AX = B$$

$$X = A^{-1}B$$

$$\text{Adj } A = \begin{bmatrix} -2 & 2 \\ -5 & 3 \end{bmatrix}$$

$$A^{-1} = \frac{\text{Adj } A}{|A|} = \frac{\begin{bmatrix} -2 & 2 \\ -5 & 3 \end{bmatrix}}{4}$$

$$= \begin{bmatrix} \frac{-2}{4} & \frac{2}{4} \\ \frac{-5}{4} & \frac{3}{4} \end{bmatrix}$$

$$X = A^{-1}B$$

$$= \begin{bmatrix} \frac{-2}{4} & \frac{2}{4} \\ \frac{-5}{4} & \frac{3}{4} \end{bmatrix} \begin{bmatrix} -6 \\ -10 \end{bmatrix}$$

$$= \begin{bmatrix} -\frac{2}{4}(-6) + \frac{2}{4}(-10) \\ \frac{-5}{4}(-6) + \frac{3}{4}(-10) \end{bmatrix}$$

MATHEMATICS FOR 9TH CLASS (UNIT # 1)

Pilot Superone Mathematics

79

Class 9th

$$\begin{bmatrix} 17 & -20 \\ 4 & 4 \\ 30 & 30 \\ 4 & 4 \end{bmatrix}$$

$$\begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 8 \\ 0 \end{bmatrix} = \begin{bmatrix} 2 \\ 0 \end{bmatrix}$$

So, $x = 2, y = 0$

Using Cramer's Rule:

$$3x - 2y = 6$$

$$5x - 2y = -10$$

$$A = \begin{bmatrix} 3 & -2 \\ 5 & -2 \end{bmatrix}$$

$$A_1 = \begin{bmatrix} -6 & 2 \\ 10 & -2 \end{bmatrix}$$

$$A_2 = \begin{bmatrix} 3 & -6 \\ 5 & -10 \end{bmatrix}$$

$$|A| = \begin{vmatrix} 3 & -2 \\ 5 & -2 \end{vmatrix}$$

$$= (3)(-2) - (5)(-2)$$

$$= -6 + 10 = 4$$

$$x = \frac{|A_1|}{|A|} = \frac{\begin{vmatrix} -6 & 2 \\ 10 & -2 \end{vmatrix}}{4}$$

$$= \frac{(-6)(-2) - (2)(-10)}{4}$$

$$= \frac{+12 - 20}{4} = \frac{-8}{4} = -2$$

MATHEMATICS FOR 9TH CLASS (UNIT # 1)

Pilot Superone Mathematics

80

Class 9th

$$\begin{aligned} y &= \frac{|A_1|}{|A|} = \frac{\begin{vmatrix} 3 & -6 \\ 5 & -10 \end{vmatrix}}{4} \\ &= \frac{(3)(-10) - (-6)(5)}{4} \\ &= \frac{-30 + 30}{4} = 0 \end{aligned}$$

So, $x = -2, y = 0$

1(v) $\begin{aligned} 3x - 2y &= 4 \\ -6x + 4y &= 7 \end{aligned}$

Writing the equations in matrix form

$$\begin{bmatrix} 3 & -2 \\ -6 & 4 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 4 \\ 7 \end{bmatrix}$$

$$\begin{bmatrix} 3 & -2 \\ -6 & 4 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 4 \\ 7 \end{bmatrix}$$

Let $AX = B$

where $A = \begin{bmatrix} 3 & -2 \\ -6 & 4 \end{bmatrix}, X = \begin{bmatrix} x \\ y \end{bmatrix}, B = \begin{bmatrix} 4 \\ 7 \end{bmatrix}$

$$\begin{aligned} |A| &= \begin{vmatrix} 3 & -2 \\ -6 & 4 \end{vmatrix} \\ &= (3)(4) - (-2)(-6) \\ &= 12 - 12 = 0 \end{aligned}$$

A is a singular matrix, therefore, solution is not possible

1(vi) $\begin{aligned} 4x + y &= 9 \\ -3x - y &= 5 \end{aligned}$

Writing the equations in matrix form.

$$\begin{bmatrix} 4 & 1 \\ -3 & -1 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 9 \\ -5 \end{bmatrix}$$

or $\begin{bmatrix} 4 & 1 \\ -3 & -1 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 9 \\ -5 \end{bmatrix}$

MATHEMATICS FOR 9TH CLASS (UNIT # 1)

Pilot Superone Mathematics 81 Class 9th

Let $AX = B$

Where $A = \begin{bmatrix} 4 & 1 \\ 3 & 1 \end{bmatrix}$, $X = \begin{bmatrix} x \\ y \end{bmatrix}$, $B = \begin{bmatrix} 9 \\ -5 \end{bmatrix}$

$$\begin{aligned} |A| &= \begin{vmatrix} 4 & 1 \\ 3 & 1 \end{vmatrix} \\ &= (4)(1) - (1)(3) \\ &= -4 + 3 = -1 \end{aligned}$$

$$AX = B$$

$$X = A^{-1}B$$

$$\text{Adj } A = \begin{bmatrix} -1 & -1 \\ 3 & 4 \end{bmatrix}$$

$$\begin{aligned} A^{-1} &= \frac{\text{Adj } A}{|A|} = \begin{bmatrix} -1 & -1 \\ 3 & 4 \end{bmatrix} \cdot \frac{1}{-1} \\ &= \begin{bmatrix} 1 & 1 \\ -3 & -4 \end{bmatrix} = \begin{bmatrix} 1 & 1 \\ -3 & -4 \end{bmatrix} \end{aligned}$$

$$X = A^{-1}B$$

$$\begin{aligned} &= \begin{bmatrix} 1 & 1 \\ -3 & -4 \end{bmatrix} \begin{bmatrix} 9 \\ -5 \end{bmatrix} \\ &= \begin{bmatrix} 1(9) + 1(-5) \\ -3(9) + (-4)(-5) \end{bmatrix} \end{aligned}$$

$$\begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 9 - 5 \\ -27 + 20 \end{bmatrix} = \begin{bmatrix} 4 \\ -7 \end{bmatrix}$$

So, $x = 4, y = -7$

Using Cramer's Rule:

$$\begin{aligned} 4x + y &= 9 \\ -3x - y &= -5 \end{aligned}$$

MATHEMATICS FOR 9TH CLASS (UNIT # 1)

Pilot Superone Mathematics 82 Class 9th

$$A = \begin{bmatrix} 4 & 1 \\ -3 & -1 \end{bmatrix}$$

$$A_x = \begin{bmatrix} 9 & 1 \\ 5 & -1 \end{bmatrix}$$

$$A_y = \begin{bmatrix} 4 & 9 \\ -3 & -5 \end{bmatrix}$$

$$|A| = \begin{vmatrix} 4 & 1 \\ -3 & -1 \end{vmatrix}$$

$$(4)(-1) - (1)(-3) \\ = -4 + 3 = -1$$

$$x = \frac{|A_x|}{|A|} = \frac{\begin{vmatrix} 9 & 1 \\ 5 & -1 \end{vmatrix}}{-1} \\ = \frac{(9)(-1) - (1)(-5)}{-1} \\ = \frac{-9 + 5}{-1} = \frac{-4}{-1} = 4$$

$$y = \frac{|A_y|}{|A|} = \frac{\begin{vmatrix} 4 & 9 \\ -3 & -5 \end{vmatrix}}{-1} \\ = \frac{(4)(-5) - (9)(-3)}{-1} \\ = \frac{-20 + 27}{-1} = \frac{7}{-1} = -7$$

So, $x = 4, y = -7$
 1(vii) $2x - 2y = 4$
 $5x - 2y = -10$

Writing the equations in matrix form.

MATHEMATICS FOR 9TH CLASS (UNIT # 1)

Prior Superone Mathematics

83

Class 9th

$$\begin{bmatrix} 2x & 2y \\ -5x & 2y \end{bmatrix} = \begin{bmatrix} 4 \\ 10 \end{bmatrix}$$

or $\begin{bmatrix} 2 & 2 \\ 5 & 2 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 4 \\ 10 \end{bmatrix}$

Let $A = \begin{bmatrix} 2 & 2 \\ 5 & 2 \end{bmatrix}$, $X = \begin{bmatrix} x \\ y \end{bmatrix}$, $B = \begin{bmatrix} 4 \\ 10 \end{bmatrix}$

Where $A = \begin{bmatrix} 2 & 2 \\ 5 & 2 \end{bmatrix}$, $X = \begin{bmatrix} x \\ y \end{bmatrix}$, $B = \begin{bmatrix} 4 \\ 10 \end{bmatrix}$

$$|A| = \begin{vmatrix} 2 & 2 \\ 5 & 2 \end{vmatrix}$$

$$= (2)(2) - (2)(5)$$

$$= 4 - 10 = -6$$

$$AX = B$$

$$X = A^{-1}B$$

$$\text{Adj } A = \begin{bmatrix} -2 & 2 \\ 5 & 2 \end{bmatrix}$$

$$A^{-1} = \frac{\text{Adj } A}{|A|} = \frac{\begin{bmatrix} -2 & 2 \\ 5 & 2 \end{bmatrix}}{-6}$$

$$\begin{bmatrix} -2 & 2 \\ 5 & 2 \end{bmatrix} \cdot \begin{bmatrix} 1 & 1 \\ 7 & -1 \end{bmatrix}$$

$$X = A^{-1}B$$

$$= \begin{bmatrix} 1 & 1 \\ -5 & 1 \end{bmatrix} \begin{bmatrix} 4 \\ -10 \end{bmatrix}$$

www.downloadclassnotes.com

MATHEMATICS FOR 9TH CLASS (UNIT # 1)

Pilot Superone Mathematics

84

Class 9th

$$= \begin{bmatrix} \frac{1}{7}(4) + \left(-\frac{1}{7}\right)(-10) \\ -\frac{5}{14}(4) + \left(-\frac{1}{7}\right)(-10) \end{bmatrix}$$

$$= \begin{bmatrix} \frac{4}{7} + \frac{10}{7} \\ -\frac{10}{7} + \frac{10}{7} \end{bmatrix} = \begin{bmatrix} \frac{14}{7} \\ 0 \end{bmatrix}$$

$$\begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 2 \\ 0 \end{bmatrix}$$

So, $x = 2, y = 0$

Using Cramer's Rule:

$$2x - 2y = 4$$

$$5x - 2y = -10$$

$$A = \begin{bmatrix} 2 & -2 \\ -10 & 2 \end{bmatrix}$$

$$A_1 = \begin{bmatrix} 4 & -2 \\ 10 & -2 \end{bmatrix}$$

$$A_2 = \begin{bmatrix} 2 & 4 \\ 5 & -10 \end{bmatrix}$$

$$|A| = \begin{vmatrix} 2 & -2 \\ -5 & -2 \end{vmatrix}$$

$$= (2)(-2) - (-2)(-5)$$

$$= -4 - 10 = -14$$

$$x = \frac{|A_1|}{|A|} = \frac{\begin{vmatrix} 4 & -2 \\ -10 & -2 \end{vmatrix}}{-14}$$

$$= \frac{4(-2) - (-2)(-10)}{-14}$$

MATHEMATICS FOR 9TH CLASS (UNIT # 1)

Pilot Superone Mathematics 85 Class 9th

$$= \frac{-8 \pm 20}{14} = \frac{28}{14} = 2$$

$$y = \frac{|A_y|}{|A|} = \frac{\begin{vmatrix} 3 & 10 \\ 1 & 4 \end{vmatrix}}{\begin{vmatrix} 3 & 4 \\ 1 & 2 \end{vmatrix}} = \frac{(2)(-10) - (4)(-5)}{14} = \frac{-20 + 20}{-4} = 0$$

So, $x = 2, y = 0$
 1(viii) $3x - 4y = 4$
 $x + 2y = 8$

writing the equations in matrix form

$$\begin{bmatrix} 3 & -4 \\ 1 & 2 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 4 \\ 8 \end{bmatrix}$$

or $\begin{bmatrix} 3 & -4 \\ 1 & 2 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 4 \\ 8 \end{bmatrix}$

Let $AX = B$

Where $A = \begin{bmatrix} 3 & -4 \\ 1 & 2 \end{bmatrix}$ $X = \begin{bmatrix} x \\ y \end{bmatrix}$ $B = \begin{bmatrix} 4 \\ 8 \end{bmatrix}$

$$|A| = \begin{vmatrix} 3 & -4 \\ 1 & 2 \end{vmatrix} = 3(2) - (4)(1) = 6 - 4 = 2$$

$$AX = B$$

$$X = A^{-1}B$$

$$\text{Adj } A = \begin{bmatrix} 2 & 4 \\ -1 & 3 \end{bmatrix}$$

www.dn.com

MATHEMATICS FOR 9TH CLASS (UNIT # 1)

Pilot Superone Mathematics

86

Class 9th

$$X = \frac{\text{Adj } A}{|A|} \cdot \begin{bmatrix} 2 & 4 \\ -1 & 3 \end{bmatrix}$$

$$\begin{bmatrix} 2 & 4 \\ 10 & 10 \\ -1 & 3 \\ 10 & 10 \end{bmatrix}$$

$$X = A^{-1}B$$

$$= \begin{bmatrix} 2 & 4 \\ 10 & 10 \\ -1 & 3 \\ 10 & 10 \end{bmatrix} \begin{bmatrix} 4 \\ 8 \end{bmatrix}$$

$$= \begin{bmatrix} \frac{2}{10}(4) + \frac{4}{10}(8) \\ -\frac{1}{10}(4) + \frac{3}{10}(8) \end{bmatrix}$$

$$= \begin{bmatrix} \frac{8}{10} + \frac{32}{10} \\ -\frac{4}{10} + \frac{24}{10} \end{bmatrix}$$

$$\begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} \frac{40}{10} \\ \frac{20}{10} \end{bmatrix} = \begin{bmatrix} 4 \\ 2 \end{bmatrix}$$

So, $x = 4, y = 2$

Using Cramer's Rule:

$$3x - 4y = 4$$

$$x + 2y = 8$$

MATHEMATICS FOR 9TH CLASS (UNIT # 1)

Pilot Supreme Mathematics 87 Class 9th

$$A = \begin{bmatrix} 3 & -4 \\ 1 & 2 \end{bmatrix}$$

$$A_1 = \begin{bmatrix} 4 & -4 \\ 8 & 2 \end{bmatrix}$$

$$A_2 = \begin{bmatrix} 3 & 4 \\ 1 & 8 \end{bmatrix}$$

$$|A| = \begin{vmatrix} 3 & -4 \\ 1 & 2 \end{vmatrix}$$

$$x = \frac{|A_1|}{|A|} = \frac{\begin{vmatrix} 4 & -4 \\ 8 & 2 \end{vmatrix}}{10}$$

$$= \frac{(4)(2) - (4)(8)}{10}$$

$$= \frac{8 - 32}{10} = \frac{-24}{10} = -2.4$$

$$y = \frac{|A_2|}{|A|} = \frac{\begin{vmatrix} 3 & 4 \\ 1 & 8 \end{vmatrix}}{10}$$

$$= \frac{(3)(8) - (4)(1)}{10}$$

$$= \frac{24 - 4}{10} = \frac{20}{10} = 2$$

So, $x = -2.4, y = 2$

Solve the following word problems by using

- matrix inversion method
- Cramer's rule

- The length of a rectangle is 4 times its width. The perimeter of the rectangle is 150cm. Find the dimensions of the rectangle.

MATHEMATICS FOR 9TH CLASS (UNIT # 1)

Pilot Superone Mathematics

88

Class 9th

3. Two sides of a rectangle differ by 35cm. Find the dimensions of the rectangle its perimeter is 67cm.
4. The third angle of an isosceles triangle is 16° less than the sum of the two equal angles. Find three angles of the triangle
5. One acute angle of a right triangle is 12° more than twice the other acute angle. Find the acute angles of the right triangle.
6. Two cars that are 600 km apart are moving towards each other. Their speeds differ by 6 km per hour and the cars are 123 km apart after $4\frac{1}{2}$ hours. Find the speed of each car.

Q.2. The length of a rectangle is 4 times its width. The perimeter of the rectangle is 150 cm. find the dimensions of the rectangle.

Suppose length of rectangle

$$= x \text{ cm}$$

$$= y \text{ cm}$$

According to the given conditions

$$4x = y$$

$$\text{and } 2x + 2y = 150$$

$$\text{Thus } 4x - y = 0 \quad (i)$$

$$2x + 2y = 150 \quad (ii)$$

Writing the equations in matrix form.

$$\begin{bmatrix} 4x & y \\ 2x & +2y \end{bmatrix} = \begin{bmatrix} 0 \\ 150 \end{bmatrix}$$

$$\text{or } \begin{bmatrix} 4 & -1 \\ 2 & 2 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 0 \\ 150 \end{bmatrix}$$

$$\text{Let } A X = B$$

$$\text{Where } A = \begin{bmatrix} 4 & -1 \\ 2 & 2 \end{bmatrix}, X = \begin{bmatrix} x \\ y \end{bmatrix}, B = \begin{bmatrix} 0 \\ 150 \end{bmatrix}$$

MATHEMATICS FOR 9TH CLASS (UNIT # 1)

Pilot Superone Mathematics

89

Class 9th

$$|A| = \begin{vmatrix} 4 & 1 \\ 2 & 2 \end{vmatrix}$$

$$= (4)(2) - (-1)(2)$$

$$= 8 + 2 = 10$$

$$AX = B$$

$$X = A^{-1}B$$

$$\text{Adj } A = \begin{bmatrix} 2 & 1 \\ -2 & 4 \end{bmatrix}$$

$$A^{-1} = \frac{\text{Adj } A}{|A|} = \frac{\begin{bmatrix} 2 & 1 \\ -2 & 4 \end{bmatrix}}{10}$$

$$= \begin{bmatrix} \frac{2}{10} & \frac{1}{10} \\ \frac{-2}{10} & \frac{4}{10} \end{bmatrix}$$

$$X = A^{-1}B$$

$$= \begin{bmatrix} \frac{2}{10} & \frac{1}{10} \\ \frac{-2}{10} & \frac{4}{10} \end{bmatrix} \begin{bmatrix} 0 \\ 150 \end{bmatrix}$$

$$= \begin{bmatrix} \frac{2}{10}(0) + \frac{1}{10}(150) \\ \frac{-2}{10}(0) + \frac{4}{10}(150) \end{bmatrix}$$

$$= \begin{bmatrix} 0 & +15 \\ 0 & +60 \end{bmatrix}$$

$$\begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 15 \\ 60 \end{bmatrix}$$

So,

$$x = 15, y = 60$$

Thus

$$\text{width} = 15 \text{ cm}$$

MATHEMATICS FOR 9TH CLASS (UNIT # 1)

Pilot Superone Mathematics 90 Class 9th

Length = 6 cm

Using Cramer's Rule:

$$\begin{aligned} 4x - y &= 0 \\ 2x + 2y &= 150 \end{aligned}$$

$$A = \begin{bmatrix} 4 & -1 \\ 2 & 2 \end{bmatrix}$$

$$A_x = \begin{bmatrix} 0 & -1 \\ 150 & 2 \end{bmatrix}$$

$$A_y = \begin{bmatrix} 4 & 0 \\ 2 & 150 \end{bmatrix}$$

$$\begin{aligned} |A| &= \begin{vmatrix} 4 & -1 \\ 2 & 2 \end{vmatrix} \\ &= (4)(2) - (-1)(2) \\ &= 8 + 2 = 10 \end{aligned}$$

$$\begin{aligned} x &= \frac{|A_x|}{|A|} = \frac{\begin{vmatrix} 0 & -1 \\ 150 & 2 \end{vmatrix}}{10} \\ &= \frac{(0)(2) - (-1)(150)}{10} \\ &= \frac{0 + 150}{10} = \frac{150}{10} = 15 \end{aligned}$$

$$\begin{aligned} y &= \frac{|A_y|}{|A|} = \frac{\begin{vmatrix} 4 & 0 \\ 2 & 150 \end{vmatrix}}{10} \\ &= \frac{(4)(150) - (0)(2)}{10} = \frac{600}{10} = 60 \end{aligned}$$

So, $x = 15, y = 60$

Q.3. Two sides of a rectangle differ by 3.5 cm. Find the dimensions of the rectangle if its perimeter is 67 cm.

Suppose:

MATHEMATICS FOR 9TH CLASS (UNIT # 1)

Pilot Superone Mathematics 91 Class 9th

Length of a rectangle = x cm

Width of a rectangle = y cm

According to the conditions

$$x + y = 35$$

and $2x + 2y = 67$

Writing the equations in matrix form

$$\begin{bmatrix} x & y \\ 2x & 2y \end{bmatrix} = \begin{bmatrix} 35 \\ 67 \end{bmatrix}$$

or $\begin{bmatrix} 1 & -1 \\ 2 & 2 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 35 \\ 67 \end{bmatrix}$

Let $AX = B$

Where $A = \begin{bmatrix} 1 & -1 \\ 2 & 2 \end{bmatrix}$, $X = \begin{bmatrix} x \\ y \end{bmatrix}$, $B = \begin{bmatrix} 35 \\ 67 \end{bmatrix}$

$$A = \begin{bmatrix} 1 & -1 \\ 2 & 2 \end{bmatrix}$$

$$\therefore (1 \times 2) - (1 \times 2)$$

$$= 2 + 2 = 4$$

$$AX = B$$

$$X = A^{-1}B$$

$$\text{Adj } A = \begin{bmatrix} 2 & 1 \\ -2 & 1 \end{bmatrix}$$

$$A^{-1} = \frac{\text{Adj } A}{|A|} = \frac{\begin{bmatrix} 2 & 1 \\ -2 & 1 \end{bmatrix}}{4} = \begin{bmatrix} \frac{2}{4} & \frac{1}{4} \\ \frac{-2}{4} & \frac{1}{4} \end{bmatrix}$$

$$X = A^{-1}B$$

$$= \begin{bmatrix} \frac{2}{4} & \frac{1}{4} \\ \frac{-2}{4} & \frac{1}{4} \end{bmatrix} \begin{bmatrix} 35 \\ 67 \end{bmatrix}$$

MATHEMATICS FOR 9TH CLASS (UNIT # 1)

Pilot Superone Mathematics 92 Class 9th

$$= \begin{bmatrix} \frac{2}{4}(35) + \frac{1}{4}(67) \\ \frac{2}{4}(35) + \frac{1}{4}(67) \end{bmatrix}$$

$$= \begin{bmatrix} \frac{70}{4} + \frac{67}{4} \\ \frac{70}{4} + \frac{67}{4} \end{bmatrix} \quad \text{②}$$

$$\begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} \frac{70}{4} \\ \frac{60}{4} \end{bmatrix} = \begin{bmatrix} 18.5 \\ 15 \end{bmatrix}$$

So $x = 18.5$

$y = 15$

Length = 18.5 cm

Width = 15 cm

Using Cramer's Rule:

$$x - y = 3.5$$

$$2x + 2y = 67$$

$$A = \begin{bmatrix} 1 & -1 \\ 2 & 2 \end{bmatrix}$$

$$A_x = \begin{bmatrix} 3.5 & -1 \\ 67 & 2 \end{bmatrix}$$

$$A_y = \begin{bmatrix} 1 & 3.5 \\ 2 & 67 \end{bmatrix}$$

$$A_1 = \begin{vmatrix} 1 & -1 \\ 2 & 2 \end{vmatrix}$$

$$(1)(2) - (-1)(2)$$

$$= 2 + 2 = 4$$

www.dn

MATHEMATICS FOR 9TH CLASS (UNIT # 1)

Pilot Superone Mathematics

93

Class 9th

$$x = \frac{A_{x1}}{|A|} = \frac{\begin{vmatrix} 3.5 & -1 \\ 67 & 2 \end{vmatrix}}{4}$$

$$= \frac{(3.5)(2) - (-1)(67)}{4} = \frac{7 + 67}{4}$$

$$= \frac{74}{4} = 18.5$$

$$y = \frac{A_{y1}}{|A|} = \frac{\begin{vmatrix} 1 & 3.5 \\ 2 & 67 \end{vmatrix}}{4}$$

$$= \frac{(1)(67) - (2)(3.5)}{4} = \frac{67 - 7}{4}$$

$$= \frac{60}{4} = 15$$

So, $x = 18.5, y = 15$
 Length = 18.5 cm
 Width = 15 cm

Q.4. The third angle of an isosceles is 16° less than the sum of the two equal angles. Find three angles of the triangle.

Solution:-

Suppose measure of each equal angle

x°

Let third angle = y°

According to the conditions

$$2x - y = 16$$

$$2x + y = 180^\circ \quad (\text{sum of the angles})$$

Writing the equations in matrix form.

$$\begin{bmatrix} 2x - y \\ 2x + y \end{bmatrix} = \begin{bmatrix} 16 \\ 180 \end{bmatrix}$$

MATHEMATICS FOR 9TH CLASS (UNIT # 1)

Pilot Superone Mathematics 94 Class 9th

$$\begin{bmatrix} 2 & 1 \\ 2 & 1 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 16 \\ 180 \end{bmatrix}$$

Let $A X = B$

Where $A = \begin{bmatrix} 2 & -1 \\ 2 & 1 \end{bmatrix}$ $X = \begin{bmatrix} x \\ y \end{bmatrix}$ $B = \begin{bmatrix} 16 \\ 180 \end{bmatrix}$

$$|A| = \begin{vmatrix} 2 & -1 \\ 2 & 1 \end{vmatrix}$$

$$= (2)(1) - (1)(2)$$

$$= 2 + 2 = 4$$

$$A X = B$$

$$X = A^{-1} B$$

$$\text{Adj } A = \begin{bmatrix} 1 & 1 \\ -2 & 2 \end{bmatrix}$$

$$A^{-1} = \frac{\text{Adj } A}{|A|} = \frac{\begin{bmatrix} 1 & 1 \\ -2 & 2 \end{bmatrix}}{4}$$

$$= \begin{bmatrix} \frac{1}{4} & \frac{1}{4} \\ -\frac{2}{4} & \frac{2}{4} \end{bmatrix}$$

$$X = A^{-1} B$$

$$= \begin{bmatrix} \frac{1}{4} & \frac{1}{4} \\ -\frac{2}{4} & \frac{2}{4} \end{bmatrix} \begin{bmatrix} 16 \\ 180 \end{bmatrix}$$

$$= \begin{bmatrix} \frac{1}{4}(16) + \frac{1}{4}(180) \\ -\frac{2}{4}(16) + \frac{2}{4}(180) \end{bmatrix}$$

MATHEMATICS FOR 9TH CLASS (UNIT # 1)

=====

Pilot Superone Mathematics

95

Class 9th

$$\begin{bmatrix} 4 + 45 \\ x + 90 \end{bmatrix}$$

$$\begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 49 \\ 82 \end{bmatrix}$$

So,

$$x = 49^\circ$$

$$y = 82^\circ$$

Measure of each equal angle 49°

Measure of third angle 82°

Using Cramer's Rule:

$$2x - y = 16$$

$$2x + y = 180$$

$$A = \begin{bmatrix} 2 & -1 \\ 2 & 1 \end{bmatrix}$$

$$A_x = \begin{bmatrix} 16 & -1 \\ 180 & 1 \end{bmatrix}$$

$$A_y = \begin{bmatrix} 2 & 16 \\ 2 & 180 \end{bmatrix}$$

$$|A| = \begin{vmatrix} 2 & -1 \\ 2 & 1 \end{vmatrix}$$

$$= (2)(1) - (-1)(2)$$

$$= 2 + 2 = 4$$

$$x = \frac{|A_x|}{|A|} = \frac{\begin{vmatrix} 16 & -1 \\ 180 & 1 \end{vmatrix}}{4}$$

$$= \frac{(16)(1) - (-1)(180)}{4}$$

$$= \frac{16 + 180}{4}$$

$$= \frac{196}{4} = 49$$

MATHEMATICS FOR 9TH CLASS (UNIT # 1)

Pilot Supercare Mathematics

96

Class 9th

$$y = \frac{A_2}{A_1} = \frac{\begin{vmatrix} 2 & 16 \\ 2 & 180 \end{vmatrix}}{J} \\ = \frac{(2)(180) - (16)(2)}{4} \\ = \frac{360 - 32}{4} = \frac{328}{4} = 82$$

So, $x = 49, y = 82^\circ$

Measure of each equal angle 49°

Measure of third angle $= 82^\circ$

Q.5. One acute angle of a right triangle is 12° more than twice the other acute angle. Find the acute angles of the right triangle.

Solution.-

Let one acute angle $= x^\circ$

and the other acute

angle $= y^\circ$

according to the conditions

$$x + y = 90^\circ$$

$$2x = y - 12$$

or $2x - y = -12$

and $x + y = 90^\circ$

Writing the equations in matrix form

$$\begin{bmatrix} 2x & y \end{bmatrix} = \begin{bmatrix} 12 \\ 90 \end{bmatrix}$$

or $\begin{bmatrix} 2 & -1 \\ 1 & 1 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} -12 \\ 90 \end{bmatrix}$

Let $A X = B$

Where $A = \begin{bmatrix} 2 & -1 \\ 1 & 1 \end{bmatrix}$ $X = \begin{bmatrix} x \\ y \end{bmatrix}$ $B = \begin{bmatrix} -12 \\ 90 \end{bmatrix}$

MATHEMATICS FOR 9TH CLASS (UNIT # 1)

Pilot Syllabus Mathematics

97

Class 9th

$$A = \begin{bmatrix} 2 & -1 \\ 1 & 1 \end{bmatrix}$$

$$= (2)(1) - (-1)(1) \\ = 2 + 1 = 3$$

$$A X = B$$

$$X = A^{-1} B$$

$$\text{Adj } A = \begin{bmatrix} 1 & 1 \\ -1 & 2 \end{bmatrix}$$

$$A^{-1} = \frac{\text{Adj } A}{A} = \begin{bmatrix} 1 & 1 \\ -1 & 2 \end{bmatrix}$$

$$= \begin{bmatrix} \frac{1}{3} & \frac{1}{3} \\ -1 & 2 \end{bmatrix}$$

$$X = A^{-1} B$$

$$= \begin{bmatrix} 1 & 1 \\ 3 & 3 \\ -1 & 2 \\ 3 & 3 \end{bmatrix} \begin{bmatrix} 12 \\ 90 \end{bmatrix}$$

$$\begin{bmatrix} \frac{1}{3}(-12) + \frac{1}{3}(90) \\ -1(-12) + 2(90) \end{bmatrix}$$

$$\begin{bmatrix} -4 + 30 \\ 12 + 180 \end{bmatrix}$$

$$\begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 26 \\ 192 \end{bmatrix}$$

So $x = 26$ & $y = 192$

One of the acute angle 26°

MATHEMATICS FOR 9TH CLASS (UNIT # 1)

Pilot Superior Mathematics 98 _____ Class '9' _____

Other acute angle 64

Using Cramer's Rule:

$$2x + y = 12$$

$$x + y = 90$$

$$A = \begin{bmatrix} 2 & 1 \\ 1 & 1 \end{bmatrix}$$

$$A_1 = \begin{bmatrix} 12 & 1 \\ 90 & 1 \end{bmatrix}$$

$$A_2 = \begin{bmatrix} 2 & 12 \\ 1 & 90 \end{bmatrix}$$

2

$$|A| = \begin{vmatrix} 2 & 1 \\ 1 & 1 \end{vmatrix} = (2)(1) - (1)(1)$$

$$= 2 - 1 = 1$$

$$x = \frac{|A_1|}{|A|} = \frac{\begin{vmatrix} -12 & -1 \\ 90 & 1 \end{vmatrix}}{1}$$

$$= \frac{(-12)(1) - (-1)(90)}{1}$$

$$= \frac{12 + 90}{1} = \frac{78}{1} = 78$$

$$y = \frac{|A_2|}{|A|} = \frac{\begin{vmatrix} 2 & 12 \\ 1 & 90 \end{vmatrix}}{1}$$

$$= \frac{(2)(90) - (-12)(1)}{1}$$

www.

MATHEMATICS FOR 9TH CLASS (UNIT # 1)

$$\frac{\text{Pilot Supermark Mathematics}}{180 \times 1^{\circ}} = \frac{99}{5} \quad \text{Class 9th}$$

$$\text{So } x = 26, y = 64$$

$$\text{One acute angle} = 26^{\circ}$$

$$\text{Other acute angle} = 64^{\circ}$$

Q.6 Two cars that are 600km apart are moving towards each other. Their speeds differ by 6km per hour and cars are 123km apart after $4\frac{1}{2}$ hours. Find the speed of each car.

Solution:

Let speed of one car = x km/hr

Speed of the other car = y km/hr

Distance covered by first car in $4\frac{1}{2}$ hours = $\frac{9}{2}x$ km

Distance covered by the other car in

$4\frac{1}{2}$ hours = $\frac{9}{2}y$ km

$$x - y = 6 \quad \text{(First condition)}$$

$$\frac{9}{2}x + \frac{9}{2}y = 600 - 123 \quad \text{(Second condition)}$$

$$\frac{9}{2}(x + y) = 477$$

$$x + y = 477 \times \frac{2}{9}$$

$$x + y = 106$$

and

$$x - y = 6$$

Writing the equations in matrix form

MATHEMATICS FOR 9TH CLASS (UNIT # 1)

Pilot Supermatic Mathematics 100 Class 9th

$$x + y = 106$$

$$x - y = 6$$

Or
$$\begin{bmatrix} 1 & 1 \\ 1 & -1 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 106 \\ 6 \end{bmatrix}$$

Let $AX = B$

Where $A = \begin{bmatrix} 1 & 1 \\ 1 & -1 \end{bmatrix}$, $X = \begin{bmatrix} x \\ y \end{bmatrix}$, $B = \begin{bmatrix} 106 \\ 6 \end{bmatrix}$

$$A^{-1} = \frac{1}{\begin{vmatrix} 1 & 1 \\ 1 & -1 \end{vmatrix}} \begin{bmatrix} -1 & 1 \\ 1 & 1 \end{bmatrix} \quad (\det A = (1)(-1) - (1)(1))$$

$$= \frac{1}{-2}$$

$$AX = B$$

$$X = A^{-1}B$$

$$\text{Id} = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$$

$$A^{-1} = \frac{Adj A}{\det A}$$

$$= \frac{1}{-2} \begin{bmatrix} -1 & 1 \\ 1 & 1 \end{bmatrix}$$

$$= \begin{bmatrix} \frac{-1}{-2} & \frac{1}{-2} \\ \frac{1}{-2} & \frac{1}{-2} \end{bmatrix} = \begin{bmatrix} \frac{1}{2} & -\frac{1}{2} \\ -\frac{1}{2} & -\frac{1}{2} \end{bmatrix}$$

MATHEMATICS FOR 9TH CLASS (UNIT # 1)

Pilot Superone Mathematics 101

..... Class 9th

$$A^{-1} B$$

$$\begin{bmatrix} 1 & 1 \\ 2 & 2 \\ 1 & 1 \\ 2 & 2 \end{bmatrix} \begin{bmatrix} 106 \\ 6 \end{bmatrix}$$

$$\begin{bmatrix} 1 & 1 \\ 2 & 2 \\ 1 & 1 \\ 2 & 2 \end{bmatrix} \begin{bmatrix} 106 \\ 6 \end{bmatrix} = \begin{bmatrix} 106 \\ 6 \end{bmatrix}$$

$$\begin{bmatrix} 1 & 1 \\ 2 & 2 \\ 1 & 1 \\ 2 & 2 \end{bmatrix} \begin{bmatrix} 106 \\ 6 \end{bmatrix} = \begin{bmatrix} 106 \\ 6 \end{bmatrix}$$

Thus $x = 56$

$y = 50$

speed of the first car = 56km/hr

speed of the other car = 50km/hr

using Cramer's rule

$$x + y = 106$$

$$x - y = 6$$

$$I = \begin{bmatrix} 1 & 1 \\ 1 & -1 \end{bmatrix}$$

$$I_1 = \begin{bmatrix} 106 & 1 \\ 6 & -1 \end{bmatrix}$$

$$I_2 = \begin{bmatrix} 1 & 106 \\ 1 & 6 \end{bmatrix}$$

$$I_3 = \begin{bmatrix} 1 & 1 \\ 1 & -1 \end{bmatrix} = (1)(-1) - (1)(1)$$

MATHEMATICS FOR 9TH CLASS (UNIT # 2)

Pilot Subject: Mathematics — 11th

**Unit
2**

REAL AND COMPLEX NUMBERS

Real Numbers

Set of Natural Numbers

$N = \{1, 2, 3, \dots\}$ Natural Numbers

Set of whole Numbers

$W = \{0, 1, 2, 3, \dots\}$

Set of Integers

$Z = \{\dots, -3, -2, -1, 0, 1, 2, 3, \dots\}$

Rational Numbers

$Q = \left\{ \frac{p}{q} : p, q \in Z, q \neq 0 \right\}$

Irrational Numbers

$Q' = \{x : x \neq \frac{p}{q}, p, q \in Z, q \neq 0\}$

For example $\pi, \sqrt{5}, \sqrt{3}, \sqrt{2}$

Real Numbers

The union of Q and Q' is a set of real numbers

It is denoted by R

$R = Q \cup Q'$

Remember:

(i) $N \subset W \subset Z \subset Q$

(ii) For each prime number p , \sqrt{p} is an irrational number

(iii) Square root of all positive non-square integers are irrational

(a) Types of Rational Numbers

The decimal representation of rational numbers are of two types: terminating and recurring

MATHEMATICS FOR 9TH CLASS (UNIT # 2)

Pilot Superone Mathematics 112 Class 9th

(i) **Terminating Decimal Fractions**

The decimal fraction in which there are finite number of digits in its decimal part is called a terminating decimal fraction. For example $\frac{4}{5} = 0.8$ and $\frac{3}{4} = 0.75$

(ii) **Recurring and Non-terminating Decimal Fraction**

The decimal fraction (non-terminating) in which some digits in the same repeated again and again in the same order in its decimal part is called a recurring decimal fraction. For example $\frac{1}{9} = 0.1111$

(b) **Irrational Numbers:**

Decimal representation for irrational numbers are neither terminating nor repeating in blocks. For example $\sqrt{2} = 1.414213562$
 π and e

MATHEMATICS FOR 9TH CLASS (UNIT # 2)

Pilot Superone Mathematics 113 Class 9th

Exercise 2.1

Q.1. Identify which of the following are rational and irrational numbers

- (i) $\sqrt{3}$ (ii) $\frac{1}{6}$ (iii) π
 (iv) $\frac{15}{2}$ (v) 7.25 (vi) $\sqrt{29}$

Solutions:

$\frac{1}{6}$ $\frac{15}{2}$ 7.25 (Rational)

$\sqrt{3}$ π , $\sqrt{29}$ (Irrational)

Q.2. Convert the following fractions into decimal fractions.

- (i) $\frac{17}{25}$ (ii) $\frac{19}{4}$ (iii) $\frac{57}{8}$
 (iv) $\frac{205}{18}$ (v) $\frac{5}{8}$ (vi) $\frac{25}{38}$

Solution:-

2(i) $\frac{17}{25}$
 0.68

25 $\frac{68}{100}$
 1.700
 150
 200
 200

2(ii) $\frac{19}{4}$
 $= 4.75$

$\frac{19}{4}$
 4.75
 4.75
 19.00
 16
 30
 28
 20
 20
 x

MATHEMATICS FOR 9TH CLASS (UNIT # 2)

Pilot Superone Mathematics	114	Class 9 th
2(iii) $\frac{57}{8}$		$\frac{125}{8}$
	7 125	57 000
		56
		10
		8
		20
		16
		40
		40
		x
2(iv) $\frac{205}{18}$		11 3889
	11 3889	18 205 0000
		18
		25
		18
		70
		54
		160
		144
		160
		144
		16
2(v) $\frac{5}{8}$		625
	0.625	8 5 0000
		48
		20
		16
		40
		40
		x
2(vi) $\frac{25}{38}$		65789
	65789	38 25 00000
		228
		220
		190
		300
		266
		340
		304
		360
		342
		18

MATHEMATICS FOR 9TH CLASS (UNIT # 2)

P lot Superone Mathematics 115 Class 9th

Q.3 Which of the statements are true and which are false?

- (i) $\frac{2}{3}$ is an irrational number
- (ii) π is an irrational number
- (iii) $\frac{1}{9}$ is a terminating fraction.
- (iv) $\frac{3}{4}$ is a terminating fraction

Solution:-

- (i) $\frac{2}{3}$ is an irrational number **False**
- (ii) π is an irrational number **True**
- (iii) $\frac{1}{9}$ is a terminating fraction **False**

Explanation:

$$\begin{array}{r} 111 \\ 9 \overline{) 1000} \\ \underline{9} \\ 10 \\ \underline{9} \\ 1 \end{array}$$

$\frac{1}{9}$ 0.1 non-terminating.

- (iv) $\frac{3}{4}$ is a terminating fraction **True**

Explanation:

- (v) $\frac{4}{5}$ is a recurring fraction. **False**


MATHEMATICS FOR 9TH CLASS (UNIT # 2)

Pilot Superone Mathematics 116 Class 9th

Q 4. Represent the following numbers on the number line.


- (i) $\frac{2}{3}$ (ii) $-\frac{4}{5}$ (iii) $1\frac{3}{4}$
 (iv) $-2\frac{5}{8}$ (v) $2\frac{3}{4}$ (vi) $\sqrt{5}$

- 4(i)** To represent the rational number $\frac{2}{3}$ divide unit length between 0 and 1 into 3 equal parts
(ii) Take 2 parts on right of 0
(iii) Point M represents $\frac{2}{3}$ on the number line in the following figure.



4(ii) Represent $-\frac{4}{5}$ on the number line.

- (i)** To represent the rational number $-\frac{4}{5}$ divide unit length between 0 and 1 into 5 equal parts
(ii) Take 4 parts on left of '0'
(iii) Point M represents $-\frac{4}{5}$ on the number line in the following figure




MATHEMATICS FOR 9TH CLASS (UNIT # 2)

Pilo, Saneetone Mathematics 117 Class 9th

4(iii) Represent $1\frac{3}{4}$ on the number line

Solution:-

$1\frac{3}{4} = 1 + \frac{3}{4}$ therefore $1\frac{3}{4}$ lies between 1 and 2 on the number line.




The distance between 1 and 2 is divided into four equal parts. From 1, we take 3 parts.

Point M represents $1\frac{3}{4}$ on the number line.

4(iv) Represent $-2\frac{5}{8}$ on the number line.


Rational number $-2\frac{5}{8}$ lies between -2 and -3.

We divide the distance between -2 and -3 into eight equal parts and take 5 parts from -2.



4(v) Represent $2\frac{3}{4}$ on the number line.

Number $2\frac{3}{4}$ lies between 2 and 3. Divided the distance between 2 and 3 into 4 equal parts. Take 3 parts from 2.




MATHEMATICS FOR 9TH CLASS (UNIT # 2)

Pilot Superone Mathematics 118 Class 9th

Point W shows $2\frac{3}{4}$ on the number line in the above figure

4(vi) Show $\sqrt{5}$ on the number line.




- (i) Construct a $\triangle OAB$ which $m\overline{OA} = 2$ units and $1\overline{BA} = 1$
 (ii) Take O as centre and draw an arc of radius OB. It cuts the number line at M. $m\overline{OM} = \sqrt{5}$.
 $(m\overline{OB})^2 = (m\overline{OA})^2 + (m\overline{AB})^2$ (Pythagorean Theorem)

$$(m\overline{OB})^2 = 4 + 1 = 5$$

$$m\overline{OB} = \sqrt{5}$$

$$\text{Thus } m\overline{OB} = m\overline{OM} = \sqrt{5}$$

Q.5. Give a rational number between $\frac{3}{4}$ and $\frac{5}{9}$



- (i) on line l a number $\frac{5}{9}$ has been shown between 0 and 1 by M.

MATHEMATICS FOR 9TH CLASS (UNIT # 2)

Pilot Superone Mathematics 115 Class 9th

- (ii) on line l a number $\frac{3}{4}$ has been shown between 0 and 1 by M (the same interval)

Result

There are a number of rational numbers between $\frac{3}{4}$ and $\frac{5}{9}$. For example $\frac{57}{100}, \frac{58}{100}, \frac{59}{100}, \frac{60}{100}, \frac{74}{100}$ etc.

- Q.6.** Express the following recurring decimals as the rational number $\frac{p}{q}$ where p, q are integers and $q \neq 0$.

- (i) $0.\overline{5}$ (ii) $0.\overline{13}$ (iii) $0.\overline{67}$

Solution:-

- 6(i)** $0.\overline{5}$

Let $x = 0.\overline{5}$

That $x = 0.5555$

Only one digit 5 is being repeated, we multiply by 10 on both sides.

$$x = 0.5555 \quad (i)$$

and $10x = (0.5555) \times 10$

That $x = 5.5555 \quad (ii)$

Subtracting (i) from (ii)

$$9x = 5$$

$$x = \frac{5}{9}$$

- 6(ii)** $0.\overline{13}$

Let $x = 0.\overline{13}$

That $x = 0.13131313$

Here, a block of 13 is being repeated, we multiply both sides by 100.

MATHEMATICS FOR 9TH CLASS (UNIT # 2)

Pilot Superone Mathematics 120 Class 9th

$$x = 0.\overline{13} \quad \overline{13} \quad \overline{13} \quad \overline{13} \quad \overline{13} \quad \dots (i)$$

$$100x = 13.\overline{13} \quad \overline{13} \quad \overline{13} \quad \overline{13} \quad \overline{13} \quad \overline{13} \quad \dots (ii)$$

Subtracting (i) from (ii)

$$100x - x = 13$$

$$99x = 13$$

$$\text{Thus } x = \frac{13}{99}$$

$$0.\overline{13} = \frac{13}{99}$$

$$6(iii) \quad 0.\overline{67}$$

$$\text{Let } x = 0.\overline{67}$$

$$\text{or } x = 0.67676767$$

Here, a block of 67 is being repeated, we multiply both sides by 100.

$$x = 0.67676767 \quad \dots (i)$$

$$100x = 67.676767 \quad \dots (ii)$$

Subtracting (i) from (ii)

$$100x - x = 67$$

$$99x = 67$$

$$\text{Thus } x = \frac{67}{99}$$

$$0.\overline{67} = \frac{67}{99}$$

Properties of Real Numbers

(a) **Properties of real numbers w.r.t. addition.**

(i) **Closure Property**

$$a + b \in R, \forall a, b \in R$$

(ii) **Commutative Property**

$$a + b = b + a, \forall a, b \in R$$

(iii) **Associative Property**

$$(a + b) + c = a + (b + c), \forall a, b, c \in R$$

MATHEMATICS FOR 9TH CLASS (UNIT # 2)

Pilot Supreme Mathematics 121 Class 9th

(iv) **Additive Identity**

$$a + 0 = 0 + a = a \quad \forall a \in R$$

(0 is called additive identity)

(v) **Additive Inverse**

$$a + (-a) = 0 \quad \forall a \in R$$

Properties of real numbers under multiplication

(i) **Closure Property**

$$ab \in R \quad \forall a, b \in R$$

(ii) **Commutative Property**

$$ab = ba \quad \forall a, b \in R$$

(iii) **Associative Property**

$$(ab)c = a(bc) \quad \forall a, b, c \in R$$

(iv) **Multiplicative Identity**

$$a \times 1 = 1 \times a = a \quad \forall a \in R$$

(v) **Multiplicative Inverse**

$$a \times \frac{1}{a} = 1 = \frac{1}{a} \times a \quad \forall a \in R \text{ and } a \neq 0$$

(vi) **Multiplication is distributive over addition and**

$$\forall a, b, c \in R$$

$$a(b + c) = ab + ac \quad \text{(Left distributive law)}$$

$$(a + b)c = ac + bc \quad \text{(Right distributive law)}$$

Properties of Equality of Real Numbers

(i) **Reflexive property**

$$a = a \quad \forall a \in R$$

(ii) **Symmetric Property**

$$\text{If } a = b \text{ then } b = a \quad \forall a, b \in R$$

(iii) **Transitive Property**

$$\text{If } a = b \text{ and } b = c, \text{ then } a = c, \quad \forall a, b, c \in R$$

(iv) **Cancellation Property for Addition**

$$\text{If } a + c = b + c, \text{ then } a = b, \quad \forall a, b, c \in R$$

(v) **Multiplicative Property**

$$\text{If } a = b, \text{ then } ac = bc, \quad \forall a, b, c \in R$$

(vi) **Cancellation Property for Addition**

$$\text{If } a + b = b + c, \text{ then } a = c, \quad \forall a, b, c \in R$$

(vii) **Cancellation Property for Multiplication**

MATHEMATICS FOR 9TH CLASS (UNIT # 2)

Pilot Superope Mathematics 122 Class 9th

- If $ac < bc$ $c \neq 0$ then $a < b$, $\forall a, b, c \in \mathbb{R}$
- (c) **Properties of Inequalities of Real Numbers**
 Properties of inequalities of real numbers are as follows.
- (i) **Trichotomy Property**
 $\forall a, b \in \mathbb{R}$
 $a < b$ or $a = b$ or $a > b$
- (ii) **Transitive Property**
 $\forall a, b, c \in \mathbb{R}$
 (a) $a < b$ and $b < c \Rightarrow a < c$
 (b) $a > b$ and $b > c \Rightarrow a > c$
- (iii) **Additive Property**
 $\forall a, b, c \in \mathbb{R}$
 (a) $a < b \Rightarrow a + c < b + c$ and
 $a < b \Rightarrow c + a < c + b$
 (b) $a > b \Rightarrow a + c > b + c$
 $a > b \Rightarrow c + a > c + b$
- (iv) **Multiplicative Property**
 (a) $\forall a, b, c \in \mathbb{R}$ and $c > 0$
 (i) $a > b \Rightarrow ac > bc$ and
 $a > b \Rightarrow ca > cb$
 (ii) $a < b \Rightarrow ac < bc$ and
 $a < b \Rightarrow ca < cb$
 (b) $\forall a, b, c \in \mathbb{R}$ and $c < 0$
 (i) $a > b \Rightarrow ac < bc$ and
 $a > b \Rightarrow ca < cb$
 (ii) $a < b \Rightarrow ac > bc$ and
 $a < b \Rightarrow ca > cb$
- (v) **Multiplicative Inverse Property**
 $\forall a, b \in \mathbb{R}$ and $a \neq 0, b \neq 0$
 (a) $a < b \Leftrightarrow \frac{1}{a} > \frac{1}{b}$
 (b) $a > b \Leftrightarrow \frac{1}{a} < \frac{1}{b}$

MATHEMATICS FOR 9TH CLASS (UNIT # 2)

Pilot Superone Mathematics

123

Class 9th

Exercise 2.2

Q.1. Identify the property used in the following.

(i) $a + b = b + a$..

(ii) $(ab)c = a(bc)$..

(iii) $7 \times 1 = 7$..

(iv) $x > y$ or $x = y$ or $x < y$..

(v) $ab = ba$..

(vi) $a + c + b + c \Rightarrow a = b$..

(vii) $5 + (-5) = 0$..

(viii) $7 \times \frac{1}{7} = 1$..

(ix) $a > b \Rightarrow ac > bc$ ($c > 0$)

Solution.-

(i) $a + b = b + a$.. (Commutative w.r.t addition)

(ii) $(ab)c = a(bc)$.. (Associative w.r.t multiplication)

(iii) $7 \times 1 = 7$.. (Multiplicative w.r.t Identity)

(iv) $x > y$ or $x = y$ or $x < y$.. (Trichotomy)

(v) $ab = ba$.. (Commutative w.r.t Multiplication)

(vi) $a + c = b + c \Rightarrow a = b$
(Cancellation property of addition)

(vii) $5 + (-5) = 0$.. (Additive Inverse)

(viii) $7 \times \frac{1}{7} = 1$.. (Multiplicative inverse)

(ix) $a > b \Rightarrow ac > bc$ ($c > 0$) ..
(Multiplicative property)

Q.2. Fill in the following blanks by stating the properties of real numbers used.

$3x + 3(y - x)$

$= 3x + 3y - 3x$.. (i)

$= 3x - 3x + 3y$.. (ii)

$= 0 + 3y$.. (iii)

$= 3y$.. (iv)

MATHEMATICS FOR 9TH CLASS (UNIT # 2)

Pilot Superone Mathematics 124 Class 9th

- 2(i) $3x + 3(y - x)$
 $= 3x + 3y - 3x$
 Distributive property of multiplication over subtraction.
- (ii) $3x + 3y = 3x + 3y$ (Commutative)
 $= 3x + 3y$
- (iii) $3x - 3x + 3y$
 $= 0 + 3y$ (Addition Inverse)
- (iv) $0 + 3y$
 $= 3y$ (Addition Identity)

Q.3. Give the name of property used in the following.

- (i) $\sqrt{24} + 0 = \sqrt{24}$
- (ii) $\frac{2}{3} \left(5 + \frac{7}{2} \right) = \left(-\frac{2}{3} \right) (5) + \left(-\frac{2}{3} \right) \left(\frac{7}{2} \right)$
- (iii) $\pi + (-\pi) = 0$
- (iv) $\sqrt{3} \cdot \sqrt{3}$
- (v) $\left(\frac{5}{8} \right) \left(-\frac{8}{5} \right) = 1$

Solution:-

- (i) $\sqrt{24} + 0 = \sqrt{24}$ (Additive Identity)
- (ii) $-\frac{2}{3} \left(5 + \frac{7}{2} \right) = \left(-\frac{2}{3} \right) (5) + \left(-\frac{2}{3} \right) \left(\frac{7}{2} \right)$
 Distributive property of multiplication over addition
- (iii) $\pi + (-\pi) = 0$ (Additive Inverse)
- (iv) $\sqrt{3} \cdot \sqrt{3}$ is a real number
 (Closure property w.r.t multiplication)
- (v) $\left(\frac{5}{8} \right) \left(-\frac{8}{5} \right) = 1$
 (Multiplication Inverse)

Concept of Radicals and Radicands

If n is positive integer greater than 1 and a is a real number, then any real number x such that $x^n = a$ is called the n th root of a , and in symbols is written as

MATHEMATICS FOR 9TH CLASS (UNIT # 2)

Pilot Superone Mathematics 125 Class 9th

$$x = \sqrt[n]{a} \quad \text{or} \quad x = (a)^{1/n}$$

In the radical $\sqrt[n]{a}$, the symbol $\sqrt{\quad}$ is called the radical sign, n is called the index of the radical and the real number a under the radical sign is called the radicand or base.

Note

$$\sqrt[n]{a} \text{ is usually written as } \sqrt{a}$$

Properties of Radicals

Let $a, b \in \mathbb{R}$ and m, n be positive integers. Then,

- | | |
|--|--|
| (i) $\sqrt[n]{ab} = \sqrt[n]{a} \sqrt[n]{b}$ | (ii) $\sqrt[n]{\frac{a}{b}} = \frac{\sqrt[n]{a}}{\sqrt[n]{b}}$ |
| (iii) $\sqrt[n]{\sqrt[m]{a}} = \sqrt[nm]{a}$ | (iv) $\sqrt[n]{a^m} = (\sqrt[n]{a})^m$ |
| (v) $\sqrt[n]{a^n} = a$ | |

MATHEMATICS FOR 9TH CLASS (UNIT # 2)

Pilot Superone Mathematics 126

Class 9th

Exercise 2.3

1. Write each radical expression in exponential notation and each exponential expression in radical notation. Do not simplify.

(i) $\sqrt[3]{-64}$

(ii) $2^{1/5}$

(iii) $-7^{1/3}$

(iv) $y^{2/3}$

(i) $\sqrt[3]{-64} = (-64)^{1/3}$ (Exponential form)

(ii) $2^{1/5} = \sqrt[5]{2}$ (Radical form)

(iii) $-7^{1/3} = -\sqrt[3]{7}$ (Radical form)

(iv) $y^{2/3} = \sqrt[3]{y^2}$ (Radical form)

- Q.2. Tell whether the following statements are true or false?

(i) $5^{1/5} = \sqrt{5}$

(ii) $2^{2/3} = \sqrt[3]{4}$

(iii) $\sqrt{49} = \sqrt{7}$

(iv) $\sqrt[3]{x^{27}} = x^3$

Solutions:-

2(i) $5^{1/5} \neq \sqrt{5}$

(False)

$5^{1/5} = \sqrt[5]{5}$

Explanation

2(ii) $2^{2/3} = \sqrt[3]{4}$

(True)

$2^{2/3} = \sqrt[3]{2^2} = \sqrt[3]{4}$

Explanation

2(iii) $\sqrt{49} = \sqrt{7}$

(False)

$\sqrt{49} = 7$

Explanation

2(iv) $\sqrt[3]{x^{27}} = x^3$

(False)

$\sqrt[3]{x^{27}} = (x^{27})^{1/3}$

Explanation

MATHEMATICS FOR 9TH CLASS (UNIT # 2)

Pilot Superone Mathematics

127

Class 9th

$$= x^{2 \times 3} \times x^3 = x^9$$

Q.3 Simplify the following radical expressions.

(i) $\sqrt[3]{125}$

(ii) $\sqrt[4]{32}$

(iii) $\sqrt[5]{\frac{3}{32}}$

(iv) $\sqrt[3]{-\frac{8}{27}}$

Solutions:

3(i) $\sqrt[3]{125}$

$$\begin{aligned} &= \sqrt[3]{(-5)^3} = [(-5)^3]^{1/3} \\ &= (-5)^{3 \times 1/3} \\ &= (-5) \end{aligned}$$

3(ii) $\sqrt[4]{32}$

$$\begin{aligned} &= \sqrt[4]{2^5} = \sqrt[4]{2^4 \times 2} \\ &= (2^4 \times 2)^{1/4} \\ &= 2^{4 \times 1/4} \times 2^{1/4} \\ &= 2\sqrt[4]{2} \end{aligned}$$

3(iii) $\sqrt[3]{\frac{3}{32}}$

$$\begin{aligned} \sqrt[3]{\frac{3}{32}} &= \left(\frac{3}{2^5}\right)^{1/3} \\ &= \frac{\sqrt[3]{3}}{2} \end{aligned}$$

3(iv) $\sqrt[3]{-\frac{8}{27}}$

$$= \sqrt[3]{\left(-\frac{2}{3}\right)^3} = \left[\left(-\frac{2}{3}\right)^3\right]^{1/3}$$

MATHEMATICS FOR 9TH CLASS (UNIT # 2)

Pilot Superior Mathematics 128 Class 9th

$$\left(-\frac{2}{3} \right)^{3 \times (-1)} = -\frac{2}{3}$$

Base and Exponent

In the exponential notation a^n (read as a to the n th power) we call ' a ' as the base and ' n ' exponent or the power to which the base is raised.

Laws of Exponents

If $a, b \in R$ and m, n are positive integers then

$$(a^m)^n = a^{mn}, (ab)^n = a^n b^n$$

$$\left(\frac{a}{b} \right)^n = \frac{a^n}{b^n} \quad b \neq 0$$

$$a^m a^n = a^{m+n}$$

$$\frac{a^m}{a^n} = a^{m-n} \quad m \geq n, a \neq 0$$

$$a^0 = 1, a^{-n} = \frac{1}{a^n}, a \neq 0$$

MATHEMATICS FOR 9TH CLASS (UNIT # 2)

Pilot Superone Mathematics

129

Class 9th

Exercise 2.4

Q.1. Use laws of exponents to simplify.

(i) $\frac{(243)^{2/3}(32)^{-1/5}}{\sqrt{(196)^{-1}}}$

(ii) $(2x^3y^{-4})(-8x^{-3}y^2)$

(iii) $\left(\frac{x^{-2}y^{-1}z^{-3}}{x^4y^{-3}z^6}\right)^{-3}$

(iv) $\frac{(81)^n 3^5 (3)^{4n-1} (243)}{(9^{2n})(3^8)}$

Solution:-

$$\begin{aligned} 1(i) \quad & \frac{(243)^{2/3}(32)^{-1/5}}{\sqrt{(196)^{-1}}} \\ & \frac{(3^5)^{2/3}(2^5)^{-1/5}}{[(14^2)^{-1}]^{1/2}} \\ & \frac{3^{10/3} 2^{-1/5}}{14^{-1} (14)^{1/2}} \\ & = \frac{3^{10/3} 2^{-1/5}}{14^{-1/2}} = \frac{3^{10/3} 2^{-1/5}}{2^1 \times 7^{-1/2}} \\ & = \frac{3^{10/3}}{2^1} \times \frac{7^{1/2}}{2^{-1/5}} \\ & = \frac{3^{10/3} 7^{1/2}}{2^{1-1/5}} = 3^{10/3} \cdot 7^{1/2} \cdot 2^{4/5} \\ & = \frac{3^3 \sqrt[3]{3} \cdot 27 \sqrt[3]{3}}{2^1 \cdot 2^{4/5}} \end{aligned}$$

$$\begin{aligned} 1(ii) \quad & (2x^3y^{-4})(-8x^{-3}y^2) \\ & = (2)(-8) x^3 y^{-4} x^{-3} y^2 \\ & = 16 x^3 y^{-4+2} \\ & = 16 x^3 y^{-2} \\ & = \frac{16x^3}{y^2} \end{aligned}$$

$$1(iii) \quad \left(\frac{x^{-2}y^{-1}z^{-3}}{x^4y^{-3}z^6}\right)^{-3}$$

MATHEMATICS FOR 9TH CLASS (UNIT # 2)

Pilot Superone Mathematics 130 Class 9th

$$= \left(\frac{y^{1+3}}{x^{1+2} z^{1+0}} \right)^{-3}$$

$$= \left(\frac{y^4}{x^3 z^1} \right)^{-3}$$

$$= \frac{y^{4 \times (-3)}}{x^{3 \times (-3)} z^{1 \times (-3)}} = \frac{y^{-12}}{x^{-9} z^{-3}} = \frac{x^9 y^{-12}}{z^{-3}}$$

$$(iv) = \frac{(81)^n \cdot 3^3 - (3)^{4n+1} (243)}{(9^{2n}) (3^3)}$$

$$= \frac{(3^4)^n \cdot 3^3 - 3^{4n+1} (3)^3}{(3^2)^{2n} (3^3)}$$

$$= \frac{3^{4n+3} - 3^{4n+1+3}}{3^{4n} \times 3^3}$$

$$= \frac{3^{4n+3} - 3^{4n+4}}{3^{4n+3}} = \frac{3^{4n+3} (3 - 3)}{3^{4n+3}}$$

$$= \frac{3^{4n+3} (3 - 1)}{3^{4n+3}} = 3^{4n+3-4n-3} \times (2)$$

$$= 3 \times 2 = 6$$

Q.2. Show that:

$$\left(\frac{x^a}{x^b} \right)^{a+b} \times \left(\frac{x^b}{x^c} \right)^{b+c} \times \left(\frac{x^c}{x^a} \right)^{c+a} = 1$$

$$L.H.S = \left(\frac{x^a}{x^b} \right)^{a+b} \times \left(\frac{x^b}{x^c} \right)^{b+c} \times \left(\frac{x^c}{x^a} \right)^{c+a}$$

$$= (x^{a-b})^{a+b} \times (x^{b-c})^{b+c} \times (x^{c-a})^{c+a}$$

$$= x^{(a-b)(a+b)} \times x^{(b-c)(b+c)} \times x^{(c-a)(c+a)}$$

$$= x^{a^2-b^2} \times x^{b^2-c^2} \times x^{c^2-a^2}$$

$$= x^{a^2-b^2+b^2-c^2+c^2-a^2}$$

$$= x^0 = 1$$

Q.3. Simplify.

$$(i) \frac{2^{1/3} (27)^{1/3} (60)^{1/2}}{(180)^{1/2} (4)^{-1/3} (9)^{1/4}} \quad (ii) \sqrt{\frac{(216)^{2/3} (25)^{1/2}}{(04)^{1/2}}}$$

$$(iii) 5^{1/3} \div (5^2)^{1/3} \quad (iv) (x^3)^2 \div x^5, x \neq 0$$

MATHEMATICS FOR 9TH CLASS (UNIT # 2)

Pilot Superone Mathematics 131 Class 9th

$$\begin{aligned}
 3(i) \quad & \frac{2^{1/3} \cdot (27)^{1/3} \cdot (60)^{1/2}}{(180)^{1/2} \cdot (4)^{1/3} \cdot (9)^{1/4}} \\
 & \frac{2^{1/3} \cdot (3^3)^{1/3} \cdot (2^2 \cdot 3 \cdot 5)^{1/2}}{(2^2 \cdot 3^2 \cdot 5)^{1/2} \cdot (2^2)^{1/3} \cdot (3^2)^{1/4}} \\
 & = \frac{2^{1/3} \cdot 3^{3 \times 1/3} \cdot 2^{2 \times 1/2} \cdot 3^{1/2} \cdot 5^{1/2}}{2^{2 \times 1/2} \cdot 3^{2 \times 1/2} \cdot 5^{1/2} \cdot 2^{2(1/3)} \cdot 3^{2 \times 1/4}} \\
 & = \frac{2^{1/3} \cdot 3 \cdot 2 \cdot 3^{1/2} \cdot 5^{1/2}}{2 \cdot 3 \cdot 5^{1/2} \cdot 2^{2/3} \cdot 3^{1/2}} = 2^{1/3 + 2/3} \\
 & = 2^{1+2/3} = 2^3 = 2^3 = 2^3 = 2
 \end{aligned}$$

$$\begin{aligned}
 3(ii) \quad & \sqrt{\frac{(216)^{2/3} \cdot (25)^{1/2}}{(64)^{1/2}}} \\
 & = \sqrt{\frac{(6^3)^{2/3} \cdot (5^2)^{1/2}}{(4)^{1/2}}} \\
 & = \sqrt{\frac{6^{3 \times 2/3} \cdot 5^{2 \times 1/2}}{(1/25)^{1/2}}} \\
 & = \sqrt{\frac{6^2 \times 5}{(1/5)^{1/2}}} \\
 & = \sqrt{\frac{6^2 \times 5}{(5^{-3})^{1/2}}} = \sqrt{\frac{6^2 \times 5}{5^{(-3/2)}}} \\
 & = \sqrt{\frac{6^2 \times 5}{5}} = (6^2)^{1/2} = 6
 \end{aligned}$$

$$\begin{aligned}
 3(iii) \quad & 5^{2^3} \div (5^2)^3 \\
 & = 5^8 \div 5^{2(3)} \\
 & = 5^8 \div 5^6 \\
 & = \frac{5^8}{5^6} = 5^{8-6} = 5^2 = 25
 \end{aligned}$$

$$\begin{aligned}
 3(iv) \quad & (x^3)^2 \div x^{3^2} \quad x \neq 0 \\
 & = x^{3(2)} \div x^9
 \end{aligned}$$

MATHEMATICS FOR 9TH CLASS (UNIT # 2)

Pilot Superone Mathematics 132 Class 9th

$$x^0 \div x^0 = \frac{x^0}{x^0} \quad \frac{1}{x^0} = \frac{1}{x^0}$$

Complex Numbers

A number $\sqrt{-1}$ has been found that square of it is “-1” It is called the imaginary unit. It is denoted by i (iota) that $i^2 = -1$ It is not a real number. The Swiss mathematician Leonard Euler (1707 – 1783) was the first to use the symbol i for the number $\sqrt{-1}$.

Numbers like $\sqrt{-1}$, $\sqrt{-3}$, $\sqrt{-4}$ etc are called imaginary numbers.

Definition of a Complex Numbers

A number of the form $z = a + bi$ where a and b are real numbers and $i = \sqrt{-1}$ is called a complex number.

Set of Complex Numbers

Set of complex numbers is denoted by C

$$C = \{ z = a + bi, \text{ where } a, b \in R \text{ and } i = \sqrt{-1} \}$$

The number ‘ a ’ is real part and b is imaginary part of the complex number $a + bi$.

These are denoted as $a = \text{Re}(z)$ and $b = \text{Im}(z)$

Conjugate of a Complex Number

Conjugate of $a + bi$ is $a - bi$, then $a + bi$ and

$$\bar{z} = a - bi$$

Now $\bar{\bar{z}} = a + bi$ i.e., conjugate of a original complex number

Equality of Complex Numbers

$$\forall a, b, c, d \in R$$

If $a + bi = c + di$ then $a = c$ and $b = d$

Properties of Complex Numbers

- (i) $z_1 = z_1$ (Reflexive Law)
- (ii) $z_1 = z_2 \rightarrow z_2 = z_1$ (Symmetric Law)
- (iii) $z_1 = z_2, z_2 = z_3 \rightarrow z_1 = z_3$ (Transitive Law)

MATHEMATICS FOR 9TH CLASS (UNIT # 2)

Pilot Superone Mathematics

133

Class 9th

Exercise 2.5

Q.1. Evaluate:

(i) i^7 (ii) i^{50} (iii) i^{12}

(iv) $(-i)^8$ (v) $(-i)^5$ (vi) i^{27}

1(i) i^7
 $= i^6 \cdot i$
 $(i^2)^3 \cdot i$
 $(-1)^3 \cdot i = -i$

1(ii) i^{50}
 $= (i^4)^{12} \cdot i^2$
 $(-1)^{12} \cdot (-1) = -1$

1(iii) i^{12}
 $= (i^2)^6$
 $= (-1)^6 = 1$

1(iv) $(-i)^8$
 $(-i)^8$
 $= [(-i)^2]^4 = (-i)^4 = 1$

1(v) $(-i)^5$
 $= (-i)^4 \times (-i)$
 $(-i)^4 \times (-i) = (i^2)^2 \times (-i)$
 $= (-1)^2 \times (-i) = -i$

1(vi) i^{27}
 $= (i^2)^{13} \times i$
 $= (-1)^{13} \times i = -i$

MATHEMATICS FOR 9TH CLASS (UNIT # 2)

Plot Superone Mathematics 134 Class 9th

Q.2. Write the conjugate of the following numbers.

(i) $2 + 3i$ (ii) $3 - 5i$ (iii) i

(iv) $3 + 4i$ (v) $-4 - i$ (vi) $1 - 3i$

Solution:-

2(i) $z = 2 + 3i$

$\bar{z} = 2 - 3i$

2(ii) $z = 3 - 5i$

$\bar{z} = 3 + 5i$

2(iii) $z = i$

$\bar{z} = -i$

2(vi) $z = -3 + 4i$

$\bar{z} = -3 - 4i$

2(v) $z = -4 - i$

$\bar{z} = -4 + i$

2(vi) $z = 1 - 3i$

$\bar{z} = -1 + 3i$

Q.3. Write the real and imaginary part of the following numbers.

(i) $1 + i$ (ii) $-1 + 2i$ (iii) $3i + 2$

(iv) $2 - 2i$ (v) $-3i$ (vi) $2 + 0i$

Solutions:-

3(i) $z = 1 + i$

MATHEMATICS FOR 9TH CLASS (UNIT # 2)

Pilot Superone Mathematics 135 Class 9th

$$Re(z) = 1 \quad Im(z) = 1$$

$$3(ii) \quad z = -1 + 2i$$

$$Re(z) = -1 \quad Im(z) = 2$$

$$3(iii) \quad z = 3i + 2$$

$$Re(z) = 2 \quad Im(z) = 3$$

$$3(iv) \quad z = -2 - 2i$$

$$Re(z) = -2, \quad Im(z) = -2$$

$$3(v) \quad z = -3i$$

$$Re(z) = 0 \quad Im(z) = -3$$

$$3(vi) \quad z = 2 + 0i$$

$$Re(z) = 2, \quad Im(z) = 0$$

Q.4. Find the value of x and y if

$$x + iy + 1 = 4 - 3i$$

$$\text{Here } iy = -3i$$

$$y = -3$$

$$\text{and } x + 1 = 4$$

$$x = 4 - 1$$

$$x = 3$$

MATHEMATICS FOR 9TH CLASS (UNIT # 2)

Pilot Superone Mathematics

146

Class 9th

Exercise 2.6

Q.1. Identify the following statements as true or false.

(i) $\sqrt{-3} \sqrt{-3} = 3$

(ii) $i^{23} = i$

(iii) $i^{10} = -1$

(iv) Complex conjugate of $(-6i + i^2)$ is $(-1 + 6i)$

(v) Difference of a complex number $z = a + bi$ and its conjugate is a real number

(vi) If $(a - 1) - (b + 3)i = 5 + 8i$, then $a = 6$ and $b = -11$

(vii) Product of a complex number and its conjugate is always a non-negative real number

Solutions:-

I(i) $\sqrt{-3} \sqrt{-3} = 3$

False

$\sqrt{3i} \times \sqrt{3i}$

Explanation

$\sqrt{-3} \sqrt{-3} = i^2$

$= 3 (-1) = -3$

I(ii) $i^7 = -i$

False

i^{23}

Explanation

$= (i)^{22} \times i$

$= (i^2)^{11} \times i$

$= (-1)^{11} \times i = -i$

I(iii) $i^{10} = -1$

True

i^{10}

Explanation

$(i^2)^5$

$= (-1)^5 = -1$

I(iv) Complex Conjugate of $(-6i + i^2)$ is $(-1 + 6i)$

True

$6i + i^2$

Explanation

$= 6i - 1 = -1 + 6i$

I(v) Difference of a complex number $z = a + bi$ and its conjugate is a real number

False

MATHEMATICS FOR 9TH CLASS (UNIT # 2)

Pilot Superone Mathematics

137

Class 9th

$$z = a + bi$$

and

$$\bar{z} = a - bi$$

$$z - \bar{z} = (a + bi) - (a - bi)$$

$$= a + bi - a + bi$$

$$= 2bi \text{ (an imaginary number)}$$

1(vi) If $(a - i) (b + 3i) = 5 + 8i$

Solution:-

$$(a - i) (b + 3i) = 5 + 8i$$

$$a - i \quad b + 3i = 5 + 8i$$

$$a - b = 5 + 8i + i - 3i$$

$$a - b = 6 + 4i$$

$$a = 6$$

and

$$b = -14 \text{ (True)}$$

1(vii) Product of a complex number and its conjugate is always a real non-negative real number

Example:

$$(2 - 3i) (2 + 3i)$$

$$= 2(2 + 3i) - 3i(2 + 3i)$$

$$= 4 + 6i - 6i - 9i^2$$

$$= 4 - 9i^2$$

$$= 4 - 9(-1) = 4 + 9 = 13$$

It is a non-negative real number (True)

Q.2. Express each complex number in the standard form $a + bi$, where a and b are real numbers.

(i) $(2 + 3i) + (7 - 2i)$

(ii) $2(5 + 4i) - 3(7 + 4i)$

(iii) $(-3 + 5i) - (4 + 9i)$

(iv) $2i^2 + 6i^3 + 3i^{16} - 6i^{19} + 4i^{21}$

Solution:-

2(i) $(2 + 3i) + (7 - 2i)$

$$= 2 + 3i + 7 - 2i$$

MATHEMATICS FOR 9TH CLASS (UNIT # 2)

Pilot Superone Mathematics 138 Class 9th

$$= 2 + 7 + 3i \quad 2i = 9 + i$$

$$2(\text{iii}) \quad -(3 + 5i) - (4 + 9i)$$

$$3 \quad 5i \quad + \quad 9i$$

$$3 \quad 4 - 5i \quad 9i = -1 \quad 14i$$

$$2(\text{iv}) \quad 2i + 6i^3 + 3i^{16} - 6i^{19} + 4i^{25}$$

$$2i^2 + 6i^3 + 3i^{16} - 6i^{19} + 4i^{25}$$

$$= 2(-1) + 6i^3 + 3(i^2)^8 - 6(i^2)^9 + 4(i^2)^{12}$$

$$= -2 + 6(-1)i + 3(-1)^8 - 6(-1)^9 + 4(-1)^{12}$$

$$= -2 - 6i + 3(1) - 6(-1)i + 4(1)i$$

$$-2 - 6i + 3 + 6i + 4i$$

$$-2 + 3 - 6i + 6i + 4i$$

$$1 + 4i$$

Q.3. Simplify and write your answer in the form $a + bi$.

$$(i) \quad (-7 + 3i)(-3 + 2i)$$

$$(ii) \quad (2 - \sqrt{-4})(3 - \sqrt{-4})$$

$$(iii) \quad (\sqrt{5} - 3i)^2$$

$$(iv) \quad (2 - 3i)(\overline{3 - 2i})$$

Solution:

$$3(i) \quad (-7 + 3i)(-3 + 2i)$$

$$(-7)(-3 + 2i) + 3i(-3 + 2i)$$

$$21 - 14i - 9i + 6i^2$$

$$21 - 14i - 9i + 6(-1)$$

$$21 - 23i - 6 - 15 - 23i$$

$$3(ii) \quad (2 - \sqrt{-4})(3 - \sqrt{-4})$$

$$(2 - 2i)(3 - 2i)$$

$$2(3 - 2i) - 2i(3 - 2i)$$

$$6 - 4i - 6i + 4i^2$$

$$6 - 10i + 4(-1)$$

$$6 - 10i - 4 - 2 - 10i$$

$$3(iii) \quad (\sqrt{5} - 3i)^2$$

MATHEMATICS FOR 9TH CLASS (UNIT # 2)

Pilot Superone Mathematics 139

Class 9th

$$\begin{aligned} & (\sqrt{5}i)^2 + (3i)^2 - 2\sqrt{5} \times 3i \\ & = 5 + 9i^2 - 6\sqrt{5}i \\ & = 5 + 9(-1) - 6\sqrt{5}i \\ & = 5 - 9 - 6\sqrt{5}i \\ & = -4 - 6\sqrt{5}i \end{aligned}$$

$$\begin{aligned} 3(iv) \quad & (2 - 3i)(\overline{3 - 2i}) \\ & = (2 - 3i)(3 + 2i) \quad (\text{Taking conjugate}) \\ & = 2(3 + 2i) - 3i(3 + 2i) \\ & = 6 + 4i - 9i - 6i^2 \\ & = 6 + 4i - 9i - 6(-1) \\ & = 6 + 4i - 9i + 6 \\ & = 12 - 5i \end{aligned}$$

Q.4. Simplify and write your answer in the form $a + bi$.

$$(i) \quad \frac{2}{1+i}$$

$$(ii) \quad \frac{2+3i}{4-i}$$

$$(iii) \quad \frac{9-7i}{3+i}$$

$$(iv) \quad \frac{2-6i}{3+i} \cdot \frac{4+i}{3-i}$$

$$(v) \quad \left(\frac{1+i}{1-i} \right)^2$$

$$(vi) \quad \frac{1}{(2+3i)(1-i)}$$

Solution:-

$$\begin{aligned} 4(i) \quad & \frac{2}{1+i} \\ & = \frac{2}{1+i} \times \frac{1-i}{1-i} \\ & = \frac{-2(1-i)}{1-i^2} = \frac{-2+2i}{1-(-1)} \\ & = \frac{-2+2i}{1+1} = \frac{-2+2i}{2} \\ & = -1+i = -1+i \end{aligned}$$

$$4(ii) \quad \frac{2+3i}{4-i}$$

MATHEMATICS FOR 9TH CLASS (UNIT # 2)

Pilot Superone Mathematics 140 Class 9th

$$= \frac{2+i}{4-i} \times \frac{4+i}{4+i}$$

$$= \frac{(2+i)(4+i)}{4^2 - i^2}$$

$$\frac{2(4+i) + 3i(4+i)}{16 - (-1)}$$

$$= \frac{8 + 2i + 12i + 3i^2}{16 + 1}$$

$$\frac{8 + 14i + 3(-1)}{17}$$

$$= \frac{8 + 14i - 3}{17}$$

$$= \frac{5 + 14i}{17} = \frac{5}{17} + \frac{14}{17}i$$

4(iii) $\frac{9-7i}{3+i}$

$$\frac{9-7i}{3+i} \times \frac{3-i}{3-i}$$

$$\frac{(9-7i) \times (3-i)}{3^2 - i^2}$$

$$\frac{3(-i) - 7i(3-i)}{9 - (-1)}$$

$$= \frac{27 - 9i - 21i + 7i^2}{9 + 1}$$

$$= \frac{27 - 30i + 7(-1)}{10} = \frac{20 - 30i}{10} = 2 - 3i$$

4(iv) $\frac{2-6i}{3+i} - \frac{4+i}{3+i}$

$$= \left(\frac{2-6i}{3+i} \times \frac{3-i}{3-i} \right) - \left(\frac{4+i}{3+i} \times \frac{3-i}{3-i} \right)$$

$$= \left(\frac{2(3-i) - 6i(3-i)}{3^2 - i^2} \right) - \left(\frac{4(3-i) + i(3-i)}{3^2 - i^2} \right)$$

MATHEMATICS FOR 9TH CLASS (UNIT # 2)

Pilot Superone Mathematics 141 Class 9th

$$= \frac{6 - 2i - 18i + 6i^2}{9 - (-1)} = \frac{12 - 4i + 3i - i^2}{9 - (-1)}$$

$$= \frac{6 - 20i + 6(-1)}{10} = \frac{12 - i - (-1)}{10}$$

$$= \frac{6 - 20i - 6}{10} = \frac{12 - i + 1}{10}$$

$$= \frac{-20i}{10} = \frac{13 - i}{10} = \frac{-20i - 13 + 1}{10}$$

$$= \frac{13 - 19i}{10} = \frac{-13}{10} - \frac{19}{10}i$$

$$\begin{aligned} 4(v) \quad & \left(\frac{1-i}{1+i} \right)^2 \\ &= \left[\frac{1-i}{1+i} \times \frac{1+i}{1+i} \right]^2 \\ &= \left[\frac{(1-i)^2}{1-i^2} \right]^2 = \left[\frac{1+2i+i^2}{1-(-1)} \right]^2 \\ &= \left[\frac{1+2i-1}{2} \right]^2 \\ &= \left[\frac{2i}{2} \right]^2 = i^2 = -1 \end{aligned}$$

$$\begin{aligned} 4(vi) \quad & \frac{1}{(2+3i)(1-i)} \\ &= \frac{1}{2+3i} \times \frac{1}{1-i} \\ &= \frac{1(2-3i)}{(2+3i)(2-3i)} \times \frac{(1+i)}{(1-i)(1+i)} \\ &= \frac{(2-3i)(1+i)}{(4-9i^2)(1-i^2)} = \frac{2(1+i)-3i(1+i)}{[4-9(-1)][1-(-1)]} \\ &= \frac{2+2i-3i-3i^2}{(4+9)(1+1)} = \frac{2-i-3(-1)}{(13)(2)} \\ &= \frac{2+3-i}{26} = \frac{5-i}{26} = \frac{5}{26} - \frac{i}{26} \end{aligned}$$

MATHEMATICS FOR 9TH CLASS (UNIT # 2)

Pilot Superone Mathematics 142 Class 9th

Q.5. Calculate (a) \bar{z} (b) $z + \bar{z}$ (c) $z - \bar{z}$ (d) $z \bar{z}$, for each of the following.

(i) $z = -i$

(ii) $z = 2 + i$

(iii) $z = \frac{1+i}{1-i}$

(iv) $z = \frac{4+3i}{2+4i}$

Solution:-

S(i) $z = -i$

(a) $z = 0 - i$

$\bar{z} = 0 + i$

(c) $z - \bar{z}$
 $= -i - (i)$
 $= -i - i$
 $= -2i$

(b) $z + \bar{z}$

$= -i + i$
 $= 0$

(d) $z \bar{z}$
 $= (-i)(i)$
 $= -i^2$
 $= -(-1)$
 $= 1$

S(ii) $z = 2 + i$

(a) $\bar{z} = 2 - i$

(b) $z + \bar{z}$
 $= (2 + i) + (2 - i)$
 $= 2 + i + 2 - i$
 $= 4$

(c) $z - \bar{z}$
 $= (2 + i) - (2 - i)$
 $= 2 + i - 2 + i$
 $= 2i$

(d) $z \bar{z}$
 $= (2 + i)(2 - i)$
 $= 4 - (i^2)$
 $= 4 - (-1)$
 $= 4 + 1 = 5$

S(iii) $z = \frac{1+i}{1-i}$

(a) $= \frac{1+i}{1-i} \times \frac{1+i}{1+i}$

MATHEMATICS FOR 9TH CLASS (UNIT # 2)

Pilot Superone Mathematics 143 Class 9th

$$= \frac{(1+i)^2}{1-i^2} = \frac{1-i^2+2i}{1+1}$$

$$= \frac{1-1+2i}{2} = i = 0+i$$

$$\bar{z} = 0 - i$$

(b) $z + \bar{z}$

$$= 1 - i - i = 0$$

(c) $z - \bar{z}$

$$= 1 - (-i)$$

$$= 1 + i = 2i$$

(d) $z \bar{z}$

$$= (i) (-i)$$

$$= (-i)^2 = -(-1) = 1$$

5(iv) $z = \frac{4-3i}{2+4i}$

$$\frac{4-3i}{2+4i} \times \frac{(2-4i)}{(2-4i)}$$

$$= \frac{(4-3i)(2-4i)}{(2)^2 - (4i)^2}$$

$$= \frac{4(2-4i) - 3i(2-4i)}{4 - 16i^2}$$

$$= \frac{8 - 16i - 6i + 12i^2}{4 - 16(-1)}$$

$$= \frac{8 - 22i + 12(-1)}{4 + 16} = \frac{-4 - 22i}{20}$$

$$= \frac{-4}{20} - \frac{22}{20}i$$

$$z = -\frac{1}{5} - \frac{11}{10}i$$

(a) $\bar{z} = \frac{1}{5} - \frac{11}{10}i$

MATHEMATICS FOR 9TH CLASS (UNIT # 2)

Pilot Superior Mathematics 144 Class 9th

$$(b) \quad z + \bar{z} = \left(-\frac{1}{5} - \frac{11}{10}i\right) + \left(-\frac{1}{5} + \frac{11}{10}i\right)$$

$$= -\frac{1}{5} - \frac{11}{10}i - \frac{1}{5} + \frac{11}{10}i$$

$$= -\frac{1}{5} - \frac{1}{5} = -\frac{2}{5}$$

$$(c) \quad z \cdot \bar{z} = \left(-\frac{1}{5} - \frac{11}{10}i\right) \left(-\frac{1}{5} + \frac{11}{10}i\right)$$

$$= \frac{1}{5} - \frac{11}{10}i + \frac{1}{5} - \frac{11}{10}i$$

$$= \frac{11}{10}i - \frac{11}{10}i$$

$$= -\frac{22}{10}i = -\frac{11}{5}i$$

$$(d) \quad z \cdot \bar{z} = \left(-\frac{1}{5} - \frac{11}{10}i\right) \left(-\frac{1}{5} + \frac{11}{10}i\right)$$

$$= \left(-\frac{1}{5}\right)^2 - \left(\frac{11}{10}i\right)^2$$

$$= \frac{1}{25} - \frac{121}{100}i^2$$

$$= \frac{1}{25} - \frac{121}{100}(-1)$$

$$= \frac{1}{25} + \frac{121}{100} = \frac{4 + 121}{100}$$

$$= \frac{125}{100} = \frac{5}{4}$$

Q.6. If $z = 2 + 3i$ and $w = 5 - 4i$, show that

$$(i) \quad z + w = \bar{z} + \bar{w} \quad (ii) \quad \overline{z - w} = \bar{z} - \bar{w}$$

$$(iii) \quad zw = \overline{z} \cdot \overline{w} \quad (iv) \quad \left(\frac{z}{w}\right) = \frac{\bar{z}}{\bar{w}} \quad w \neq 0$$

$$(v) \quad \frac{1}{2}(z + \bar{z}) \quad \text{is a real part of } z$$

MATHEMATICS FOR 9TH CLASS (UNIT # 2)

Prior Superone Mathematics 145 Class 9th

(vi) $\frac{1}{2i}(z - \bar{z})$ is the imaginary part of z

Solutions:

6(i) $z = 2 + 3i$

and

$w = 5 - 4i$

$z + w = \bar{z} + \bar{w}$ (To show)

$\bar{z} = 2 - 3i$

$\bar{w} = 5 + 4i$

Now,

$z + w = 2 + 3i + 5 - 4i$
 $= 7 - i$

$\overline{z + w} = 7 + i$ (i)

$z + w = (2 + 3i) + (5 - 4i)$
 $= 2 + 3i + 5 - 4i$
 $= 7 - i$ (ii)

$z + w = \overline{z + w} = \bar{z} + \bar{w}$ (from i and ii)

6(ii) $\overline{z - w} = \bar{z} - \bar{w}$

We find $z - w$

$z - w = (2 + 3i) - (5 - 4i)$
 $= 2 + 3i - 5 + 4i$
 $= -3 + 7i$

$\overline{z - w} = -3 - 7i$ (i)

$\bar{z} = 2 - 3i$

$\bar{w} = 5 + 4i$

$\bar{z} - \bar{w} = (2 - 3i) - (5 + 4i)$
 $= 2 - 3i - 5 - 4i$

MATHEMATICS FOR 9TH CLASS (UNIT # 2)

Pilot Superone Mathematics 146 Class 9th

$$= 3 - 7i \quad (ii)$$

$$\overline{z - w} = \overline{z} - \overline{w} \quad (\text{From i and ii})$$

$$6(iii) \quad \overline{zw} = \overline{z} \cdot \overline{w}$$

We find zw

$$\begin{aligned} zw &= (2 + 3i)(5 - 4i) \\ &= 2(5 - 4i) + 3i(5 - 4i) \\ &= 10 - 8i + 15i - 12i^2 \\ &= 10 + 7i - 12(-1) \\ &= 10 + 7i + 12 \\ &= 22 + 7i \end{aligned}$$

$$\overline{zw} = 22 - 7i \quad (i)$$

$$\overline{z} = 2 - 3i$$

$$\overline{w} = 5 + 4i$$

$$\begin{aligned} \overline{z} \cdot \overline{w} &= (2 - 3i)(5 + 4i) \\ &= 2(5 + 4i) - 3i(5 + 4i) \\ &= 10 + 8i - 15i - 12i^2 \\ &= 10 - 7i - 12(-1) \\ &= 10 - 7i + 12 \\ &= 22 - 7i \quad (ii) \end{aligned}$$

$$\overline{zw} = \overline{z} \cdot \overline{w} \quad (\text{From i and ii})$$

$$6(iv) \quad \left(\frac{z}{w} \right) = \frac{\overline{z}}{\overline{w}} \quad (w \neq 0)$$

We find $\frac{z}{w}$

$$\begin{aligned} \frac{z}{w} &= \frac{2 + 3i}{5 - 4i} \\ &= \frac{2 + 3i}{5 - 4i} \times \frac{5 + 4i}{5 + 4i} \end{aligned}$$

MATHEMATICS FOR 9TH CLASS (UNIT # 2)

Pilot Superone Mathematics 147 Class 9th

$$\begin{aligned} &= \frac{(2+3i)(5+4i)}{(5)^2 + (4i)^2} \\ &= \frac{2(5+4i) + 3i(5+4i)}{25 - 16i^2} \\ &= \frac{10 + 8i + 15i + 12i^2}{25 - 16(-1)} \\ &= \frac{10 + 23i + 12(-1)}{41} = \frac{10 - 23i - 12}{41} \\ &= \frac{-2 - 23i}{41} \end{aligned}$$

$$\frac{z}{w} = \frac{-2}{41} + \frac{23}{41}i$$

$$\left(\frac{z}{w}\right)^- = \frac{2}{41} - \frac{23}{41}i \quad (i)$$

Now,

$$\begin{aligned} \frac{\bar{z}}{\bar{w}} &= \frac{2-3i}{5+4i} \\ &= \frac{2-3i}{5+4i} \times \frac{5-4i}{5-4i} \\ \frac{\bar{z}}{\bar{w}} &= \frac{(2-3i)(5-4i)}{(5)^2 + (4i)^2} \\ &= \frac{2(5-4i) - 3i(5-4i)}{25 - 16i^2} \\ &= \frac{10 - 8i - 15i + 12i^2}{25 - 16(-1)} \\ &= \frac{10 - 23i - 12}{41} \\ &= \frac{-2 - 23i}{41} \\ &= \frac{-2}{41} - \frac{23}{41}i \quad (ii) \end{aligned}$$

MATHEMATICS FOR 9TH CLASS (UNIT # 2)

Plot Superone Mathematics 148 Class 9th

$$\left(\frac{z}{w}\right) = \frac{\bar{z}}{\bar{w}} \quad (\text{From i and ii})$$

$$\begin{aligned} 6(v) \quad & \frac{1}{2}(z + \bar{z}) \\ & \frac{1}{2}\{2 + 3i + 2 - 3i\} \\ & \frac{1}{2}\{4\} \\ & 2 \quad (2 \text{ is a real part of } z) \end{aligned}$$

$$\begin{aligned} 6(vi) \quad & \frac{1}{2}(z - \bar{z}) \\ & \frac{1}{2}\{(2 + 3i) - (2 - 3i)\} \\ & \frac{1}{2}\{2 + 3i - 2 + 3i\} \\ & \frac{1}{2}\{6i\} \\ & 3i \quad (3 \text{ is imaginary part of } z) \end{aligned}$$

Q. 7. Solve the following equations for real x and y.

- (i) $(2 - 3i)(x + yi) = 4 + i$
 (ii) $(3 - 2i)(x + yi) = 2(x - 2yi) + 2i - 1$
 (iii) $(3 + 4i)^2 = 2(x - yi) = x + yi$

Solution:-

$$\begin{aligned} 7(i) \quad & (2 - 3i)(x + yi) = 4 + i \\ & 2(x + yi) - 3i(x + yi) = 4 + i \\ & 2x + 2yi - 3xi - 3yi^2 = 4 + i \\ & 2x + 2yi - 3xi - 3y(-1) = 4 + i \\ & 2x + 2yi - 3xi + 3y = 4 + i \\ & 2x + 3y + 2yi - 3xi = 4 + i \\ & (2x + 3y) + (2y - 3x)i = 4 + i \\ & \quad \quad \quad 2x - 3y = 4 \quad (i) \\ \text{and} \quad & 2y - 3x = 1 \quad (ii) \end{aligned}$$

MATHEMATICS FOR 9TH CLASS (UNIT # 2)

Pilot Superone Mathematics 149 Class 9th

Multiply eq (i) by 3 and eq (ii) by 2

$$6x + 9y = 12 \quad (iii)$$

$$4y - 6x = 2 \quad (iv)$$

From (iv) + (iii)

$$13y = 14$$

$$y = \frac{14}{13}$$

Putting $y = \frac{14}{13}$ in (i)

$$2x + 3\left(\frac{14}{13}\right) = 4$$

$$2x + \frac{42}{13} = 4$$

$$2x = 4 - \frac{42}{13}$$

$$= \frac{52 - 42}{13}$$

$$2x = \frac{10}{13}$$

$$x = \frac{10}{2 \times 13}$$

$$x = \frac{5}{13}$$

Thus $x = \frac{5}{13}, y = \frac{14}{13}$

7(ii)

$$(3 - 2i)(x + yi) = 2(x - 2yi) + 2i - 1$$

$$3x + 3yi - 2xi - 2yi^2 = 2x - 4yi + 2i - 1$$

$$3x + 3yi - 2xi - 2y(-1) = 2x - 4yi + 2i - 1$$

$$3x + 3yi - 2xi + 2y = 2x - 4yi + 2i - 1$$

$$3x - 2x + 3yi + 4yi - 2xi + 2y = -1 + 2i$$

$$\rightarrow x + 7yi - 2xi + 2y = -1 + 2i$$

$$(x + 2y) + (7y - 2x)i = -1 + 2i$$

Thus $x + 2y = -1 \quad (i)$

MATHEMATICS FOR 9TH CLASS (UNIT # 2)

Pilot Superiore Mathematics 150 Class 9th

and $7y - 2x = 2$ (ii)

(iii) $2x + 4y = -2$

Multiplying eq (i) by 2

$11y = 0$ Adding (iii), (ii)

$y = 0$

$x + 2(0) = 1$

Putting $y = 0$ in eq (i)

$x + 0 = -1$

$x = -1$

Thus $x = -1, y = 0$

7(iii) $(3 + 4i)^2 - 2(x - yi) = x + yi$

$9 + 24i + 16i^2 - 2x + 2yi = x + yi$

$9 + 24i + 16(-1) - 2x + 2yi = x + yi = 0$

$9 + 24i - 16 - 2x + 2yi = 0$

$-1x + yi = 16 - 9 - 24i$

$-1x + yi = 7 - 24i$

Thus $-1x = 7$ (i)

and $y = 24$ (ii)

$x = -\frac{7}{1}$ From (i)

Thus $x = -\frac{7}{3}, y = 24$

MATHEMATICS FOR 9TH CLASS (UNIT # 3)

Pilot Superior Mathematics 157

Class 9th

**Unit
3**

LOGARITHMS

Scientific Notation:

Scientists have developed a concise precise and convenient method to write very small or very large numbers, that is called scientific notation of expressing an ordinary number

A number in the form $a \times 10^n$ where $1 \leq a \leq 16$ and n is an integer is called the scientific notation.

Exercise 3.1

Q.1. Express each of the following numbers in scientific notation.

(i) 5700

(ii) 49,800,000

(iii) 96 000, 000

(iv) 416 9

(v) 83 000

(vi) 0.00643

(vii) 0.0074

(viii) 60 000, 000

(ix) 0.00000000395

(x) $\frac{275,000}{0.0025}$

Solution:-

1(i) 5700

$$= 5\ 700 \times 1000$$

$$= 5\ 7 \times 10^3$$

1(ii) 49, 800, 000

$$= 4\ 9800000 \times 10000000$$

$$4\ 98 \times 10^7$$

MATHEMATICS FOR 9TH CLASS (UNIT # 3)

Pilot Superone Mathematics 158 Class 9th

1(iii) 96,000,000

$$96000000 \times 10000000$$

$$= 96 \times 10^7$$

1(iv) 416.9

$$4169 \times 100$$

$$= 4169 \times 10^2$$

1(v) 83000

$$= 83000 \times 10000$$

$$= 83 \times 10^4$$

1(vi) 0.00643

$$= \frac{643}{1000} = \frac{6.43}{10^2} = 6.43 \times 10^{-3}$$

1(vii) 0.074

$$= \frac{74}{1000} = \frac{7.4}{10^2} = 7.4 \times 10^{-2}$$

1(viii) 60,000,000

$$= 60000000 \times 10000000$$

$$= 6 \times 10^7$$

1(ix) 0.00000000395

$$\frac{395}{1000000000} = \frac{3.95}{10^9}$$

$$3.95 \times 10^{-9}$$

1(x) $\frac{275000}{0.025}$

$$= \frac{27500 \times 100000}{1000}$$

$$= \frac{275}{1000}$$

$$= \frac{275 \times 10^5}{25 \times 10^{-3}}$$

$$= \frac{275 \times 10^5}{25 \times 10^{-3}}$$

Q.2. Express the following numbers in ordinary notation.

(i) 6×10^{-4}

(ii) 5.06×10^{10}

MATHEMATICS FOR 9TH CLASS (UNIT # 3)

Pilot Superone Mathematics 159 Class 9th

(iii) 9.018×10^{-6} (iv) -7.865×10^8

Solution:-

2(i) 6×10^{-4}

$$6 \times \frac{1}{10^4} = 0.0006$$

2(ii) 5.06×10^{10}

$$= 50600000000$$

2(iii) 9.018×10^{-6}

$$= 9.018 \times \frac{1}{10^6} = 0.000009018$$

2(iv) 7.865×10^8

$$= 786500000$$

Logarithm of a Real Number

If $a^x = y$, then x is called the logarithm of 'y' to the base 'a' and is written as $\log_a y = x$, where $a > 0$, $a \neq 1$ and $y > 0$.

If $a^x = y$

then $\log_a y = x$ and

if $\log_a y = x$

then $a^x = y$

Thus $a^x = y$ and $\log_a y = x$ are equivalent equations

$a^x = y$ is exponential form and

$\log_a y = x$ is logarithmic form

Remember:

$$a^0 = 1 \Rightarrow \log_a 1 = 0 \quad \text{and}$$

$$a^1 = a \Rightarrow \log_a a = 1$$

MATHEMATICS FOR 9TH CLASS (UNIT # 3)

Pilot Superior Mathematics 160

Class 9th

Exercise 3.2

Q.1. Find the common logarithms of the following numbers.

- (i) 232 92 (ii) 29.326
 (iii) 0.00032 (iv) 0.3206

Solution:-

1(i) $\log 232\ 92$

(i) Integral part has three (3) digits
 $ch\ 3\ l = 2$

(ii) To find mantissa, we round off 232 92 to 232 9

Log Table										Mean Differences										
	0	1	2	3	4	5	6	7	8	9	1	2	3	4	5	6	7	8	9	
23			↓																↓	17
			3650																	

(iii) We see the log table and find the row corresponding to 23.

(iv) We proceed horizontally and reach column corresponding to 2. The number at the intersection is 3655.

(v) Proceeding ahead we find a number in the mean difference column under 9. It is 17. Mantissa is 3655 + 17 = 3672.

$$\log 232\ 92 = 2.3672$$

1(ii) $\log 29.326$

(i) Integral part has (2) digits.
 $ch\ 2\ l = 1$

(ii) To find mantissa, we round off 29.326 to 29.32

Log Table										Mean Differences										
	0	1	2	3	4	5	6	7	8	9		1	2	3	4	5	6	7	8	9
29				↓										↓						
				669										4						

MATHEMATICS FOR 9TH CLASS (UNIT # 3)

Pilot Supreme Mathematics 161 Class 9th

- (iii) We read the log table and find the row corresponding to 29
- (iv) We proceed horizontally and reach column corresponding to 3. The number at the intersection is 4669
- (v) Proceed ahead and we find a number in the mean differences column under 3. It is 4. Mantissa is $4669 + 4 = 4673$
- $\therefore \log 29.326 = 1.4673$
- 1(iii) $\log 0.00032$
- (i) There are three zeros on the right of the decimal point:-

$$ch = \bar{4}$$

Log Table										Mean Differences	
	0	1	2	3	4	5	6	7	8	9	123456789
32	↓	5051									

- (ii) Corresponding to 32 in the log table, we proceed horizontally and under 0 find a number 5051
- Mantissa = 5051

Thus $\log 0.00032 = \bar{4}.5051$

1(iv) $\log 0.3206$

- (i) The integral part of 0.3206 is '0' and there is no zero on the right of decimal point

$$ch = 1$$

Log Table											Mean Differences
	0	1	2	3	4	5	6	7	8	9	123456789
32	↓	5051									↓
											8

MATHEMATICS FOR 9TH CLASS (UNIT # 3)

Pilot Superone Mathematics 162 Class 9th

- (ii) Corresponding to 32 in the log table, we proceed horizontally and under 0 find a number 5051
 (iii) We proceed ahead and find in the mean difference column under 6 number 8.
 $\text{mantissa} = 5051 + 8 = 5059$

$$\log 0.3206 = \bar{1} 5059$$

Q.2. If $\log 31.09 = 1.4926$, find the values of following.

- (i) $\log 3109$ (ii) $\log 310.9$
 (iii) $\log 0.003109$ (iv) $\log 0.3109$

Solution:-

- (2) $\log 31.09 = 1.4926$ (Given)
 2(i) $\log 3109 = 0.4926$ (Mantissa does not change)
 2(ii) $\log 3109 = 2.4926$

$$2(\text{iii}) \quad \log 0.003109 = \bar{3} 4926$$

$$2(\text{iv}) \quad \log 0.3109 = \bar{1} 4926$$

Q.3. Find the numbers whose common logarithms are:

- (i) 3.5621 (ii) 1.7427

- 3(i) Let the number = x
 $\log x = 3.5622$
 here ch 3

Anti-log Table											Mean Differences
	0	1	2	3	4	5	6	7	8	9	123456789
56	3648										2

- (i) Corresponding to 56 in the antilog table, we proceed horizontally and under 2, find a number 3648
 (ii) We proceed ahead and find in the mean difference column under 2 number 2

MATHEMATICS FOR 9TH CLASS (UNIT # 3)

Pilot Superone Mathematics 163 Class 9th

$$\text{Now } 3648 \div 2 = 3650$$

ch 3 therefore, there are four digits in the number

$$\text{Required Number } x = 3650$$

3(ii) Let the number x

$$\log x = \bar{2}.7427$$

$$\text{here } ch = \bar{2}$$

Therefore, there will be ONE zeros on the right side of decimal point

Anti-log Table

											Mean Differences
	0	1	2	3	4	5	6	7	8	9	1 2 3 4 5 6 7 8 9
74				5521							9

(i) Corresponding to 74 in the ant.log table, we proceed horizontally and under 2, find a number 5521

(ii) We proceed ahead and find in the mean difference column under 7 number 9

$$\text{Now } 5521 + 9 = 5530$$

ch = 2, therefore, there are four digits in the number

$$\text{Required Number } x = 05530$$

Q.4. What replacement for the unknown in each of the following will make the statement true?

$$(i) \log_3 81 = L \quad (ii) \log_a 6 = 0.5$$

$$(iii) \log_5 n = 2 \quad (iv) 10^x = 40$$

$$4(i) \log_3 81 = L$$

$$3^L = 81 \quad (\text{Writing in the exponential form})$$

$$3^L = 3^4$$

$$L = 4$$

$$4(ii) \log_a 6 = 0.5$$

$$6 = a^{0.5} \quad (\text{Writing in the exponential form})$$

$$6 = a^{5/10}$$

MATHEMATICS FOR 9TH CLASS (UNIT # 3)

Pilot Superior Mathematics 164 Class 9th

$$6 = a^{1/2}$$

$$6^2 = (a^{1/2})^2 \quad (\text{Squaring})$$

$$36 = a^{1/2 \times 2}$$

$$36 = a$$

Thus $a = 36$

4(iii) $\log_5 n = 2$

$$n = 5^2 \quad (\text{Writing in the exponential form})$$

$$n = 25$$

4(iv) $10^p = 40$

$$p = \log 40 \quad (\text{Taking log})$$

$$p \log 10 = 1.6021 \quad (\text{using log table})$$

$$p = 1.6021 \quad (\log 10 = 1)$$

Q.5. Evaluate:

(i) $\log_2 \frac{1}{128}$ (ii) $\log 512$ to the base $2\sqrt{2}$

5(i) $\log_2 \frac{1}{128}$

Let $x = \log_2 \frac{1}{128}$

$$2^x = \frac{1}{128} \quad (\text{Writing in exponential form})$$

$$= \frac{1}{2^7}$$

$$2^x = 2^{-7}$$

Thus $x = -7$

5(ii) $\log 512$ to the base $2\sqrt{2}$

Let $x = \log_{2\sqrt{2}} 512$

$$(2\sqrt{2})^x = 512 \quad (\text{Writing in exponential form})$$

$$(\sqrt{2} \times 2 \times 2)^x = 512$$

$$(\sqrt{8})^x = 512$$

$$(8^{1/2})^x = 8^3$$

$$8^{x/2} = 8^3$$

MATHEMATICS FOR 9TH CLASS (UNIT # 3)

Pilot Superone Mathematics 165 Class 9th

Thus $\frac{x}{2} = 3$

Hence $x = 3 \times 2 = 6$

Q.6. Find the value of x from the following statements.

(i) $\log_2 x = 5$ (ii) $\log_8 9 = x$

(iii) $\log_{64} 8 = \frac{x}{2}$ (iv) $\log_x 64 = 2$

(v) $\log_3 x = 4$

6(i) $\log_2 x = 5$

$x = 2^5$ (Writing in exponential form)

$x = 32$

6(ii) $\log_8 9 = x$

$(81)^x = 9$ (Writing in exponential form)

$(9^2)^x = 9$

$9^{2x} = 9$

$2x = 1$

or $x = \frac{1}{2}$

6(iii) $\log_{64} 8 = \frac{x}{2}$

$(64)^{\frac{x}{2}} = 8$ (Writing in exponential form)

$8^{2 \times \frac{x}{2}} = 8$

$8^x = 8$

Thus $x = 1$

6(iv) $\log_x 64 = 2$

$x^2 = 64$ (Writing in exponential form)

$x^2 = 8^2$

Thus $x = 8$

6(v) $\log_3 x = 4$

$x = 3^4$ (Writing in exponential form)

Thus $x = 81$

MATHEMATICS FOR 9TH CLASS (UNIT # 3)

Pilot Superone Mathematics 166 Class 9th

(Laws of Logarithm)

Law (i)

$$\log_a^{(mn)} = \log_a^m + \log_a^n$$

Proof

Let $\log_a^m = x, \log_a^n = y$

$$a^x = m \quad (i)$$

and $a^y = n \quad (ii)$

$$a^x \times a^y = m \times n \quad \text{from (i) } \times \text{ (ii)}$$

$$a^{x+y} = mn$$

$$\log_a^{mn} = x + y \quad (\text{Writing log form})$$

$$= \log_a^m + \log_a^n \quad (\text{Writing values of } x, y)$$

Law (ii)

$$\log_a^{m/n} = \log_a^m - \log_a^n$$

Proof

Let $\log_a^m = x$

and $\log_a^n = y \quad (\text{Writing in exponential form})$

$$a^x = m \quad (i)$$

$$a^y = n \quad (ii) \quad \text{Dividing (i) by (ii)}$$

$$\frac{a^x}{a^y} = \frac{m}{n}$$

$$a^{x-y} = \frac{m}{n} \quad (\text{Writing in log form})$$

$$\log_a^{m/n} = x - y \quad (\text{Putting values of } x, y)$$

$$\log_a^{m/n} = \log_a^m - \log_a^n$$

Law (iii)

$$\log_a^{m^n} = n \log_a^m$$

Proof.

Let $\log_a^m = x$

MATHEMATICS FOR 9TH CLASS (UNIT # 3)

Pilot Superone Mathematics 167 Class 9th

$$\text{or } a^x = m^n$$

$$\text{Let } \log_a m = y$$

$$\text{or } a^y = m \quad (i)$$

$$a^x = m^n$$

$$= (a^y)^n \quad \text{From (i)}$$

$$a^x = a^{yn}$$

$$a^x = a^{yn}$$

$$\text{So } x = yn = ny$$

$$\log_a m^n = n \log_a m$$

Law (iv) (Change of base)

$$\log_a m = \log_b m \times \log_a b = \frac{\log_b m}{\log_b a}$$

Proof

$$\text{Let } \log_a b = x \Rightarrow a^x = b$$

$$\log_a a^x = \log_a b = x \log_a b \quad (\text{Taking log to base } a)$$

$$\log_a a^x = \log_b a \log_a b$$

$$\log_b a \times \log_a b = \log_a a = 1$$

$$\text{or } \log_a b = \frac{1}{\log_b a} \quad \text{Putting value of } \log_a b \text{ in (i)}$$

$$\log_a m = \frac{\log_b m}{\log_b a}$$

Exercise 3.3

Q.1. Write the following into sum or difference

$$(i) \log(A \times B) \qquad (ii) \log \frac{15.2}{30.5}$$

$$(iii) \log \frac{25 \times 5}{8} \qquad (iv) \log \sqrt[3]{\frac{7}{15}}$$

$$(v) \log \frac{(22)^{1/3}}{5^3} \qquad (vi) \log \frac{25 \times 47}{29}$$

Solution:-

$$\begin{aligned} I(i) \quad & \log(A \times B) \\ &= \log A + \log B \end{aligned}$$

$$\begin{aligned} I(ii) \quad & \log \frac{15.2}{30.5} \\ &= \log 15.2 - \log 30.5 \end{aligned}$$

$$\begin{aligned} I(iii) \quad & \log \frac{25 \times 5}{8} \\ &= \log 25 + \log 5 - \log 8 \end{aligned}$$

$$\begin{aligned} I(iv) \quad & \log \sqrt[3]{\frac{7}{15}} \\ &= \log \left(\frac{7}{15} \right)^{1/3} \\ &= \frac{1}{3} [\log 7 - \log 15] \end{aligned}$$

$$\begin{aligned} I(v) \quad & \log \frac{(22)^{1/3}}{5^3} \\ &= \log (22)^{1/3} - \log 5^3 \\ &= \frac{1}{3} \log 22 - 3 \log 5 \end{aligned}$$

$$\begin{aligned} I(vi) \quad & \log \frac{25 \times 47}{29} \\ &= \log 25 + \log 47 - \log 29 \end{aligned}$$

MATHEMATICS FOR 9TH CLASS (UNIT # 3)

Pilot Superone Mathematics 169 Class 9th

Q.2. Express as a single logarithm.

$$\log x - 2\log x + 3\log(x+1) - \log(x^2 - 1)$$

Solution:-

$$\begin{aligned} & \log x - 2\log x + 3\log(x+1) - \log(x^2 - 1) \\ &= \log x - \log x^2 + \log(x+1)^3 - \log(x^2 - 1) \\ &= \log \frac{x(x+1)^3}{x^2(x^2 - 1)} \\ &= \log \frac{(x+1)^3}{x(x^2 - 1)} \\ &= \log \frac{(x+1)^3}{x(x+1)(x-1)} \\ &= \log \frac{(x+1)^2}{x(x-1)} \end{aligned}$$

Q.3. Write the following in the form of a single logarithm.

(i) $\log 21 + \log 5$ (ii) $\log 25 - 2\log 3$

(iii) $2\log x - 3\log y$ (iv) $\log 5 + \log 6 - \log 2$

Solution:-

3(i) $\log 21 + \log 5$
 $= \log 21 \times 5$

3(ii) $\log 25 - 2\log 3$
 $= \log 25 - \log 3^2$
 $= \log \frac{25}{3^2}$

3(iii) $2\log x - 3\log y$
 $= \log x^2 - \log y^3$
 $= \log \frac{x^2}{y^3}$

3(iv) $\log 5 + \log 6 - \log 2$
 $= \log \frac{5 \times 6}{2}$

MATHEMATICS FOR 9TH CLASS (UNIT # 3)

Pilot Superone Mathematics 170 Class 9th

Q.4. Calculate the following:

(i) $\log_3 2 \times \log_2 81$ (ii) $\log_3 3 \times \log_3 25$

Solution:-

$$\begin{aligned} 4(i) \quad & \log_3 2 \times \log_2 81 \\ &= \frac{\log 81}{\log 3} \cdot \frac{\log 3^4}{\log 3} \\ &= \frac{4 \log 3}{\log 3} = 4 \end{aligned}$$

$$\begin{aligned} 4(ii) \quad & \log_3 3 \times \log_3 25 \\ &= \frac{\log 25}{\log 5} \cdot \frac{\log 5^2}{\log 5} \\ &= \frac{2 \log 5}{\log 5} = 2 \end{aligned}$$

Q.5. If $\log 2 = 0.3010$, $\log 3 = 0.4771$, $\log 5 = 0.6990$ then find the values of the following.

(i) $\log 32$ (ii) $\log 24$
 (iii) $\log \sqrt{3\frac{1}{3}}$ (iv) $\log \frac{8}{3}$
 (v) $\log 30$

Solution:-

$$\begin{aligned} 5(i) \quad & \log 32 \\ &= \log 2^5 \\ &= 5 \log 2 = 5(0.3010) = 1.5050 \end{aligned}$$

$$\begin{aligned} 5(ii) \quad & \log 24 \\ &= \log 8 \times 3 \\ &= \log 2^3 \times 3 = \log 2^3 + \log 3 \\ &= 3 \log 2 + \log 3 \\ &= 3(0.3010) + 0.4771 \\ &= 0.9030 + 0.4771 = 1.3801 \end{aligned}$$

$$5(iii) \quad \log \sqrt{3\frac{1}{3}}$$

MATHEMATICS FOR 9TH CLASS (UNIT # 3)

Pilot Superone Mathematics 171 Class 9th

$$= \log \left(\frac{10}{3} \right)^{1/2}$$

$$= \frac{1}{2} \log \frac{10}{3}$$

$$= \frac{1}{2} \log \frac{2 \times 5}{3}$$

$$= \frac{1}{2} [\log 2 + \log 5 - \log 3]$$

$$= \frac{1}{2} [0.3010 + 0.6990 - 0.4771]$$

$$= \frac{1}{2} (0.5229) = 0.2615$$

5(iv) $\log \frac{8}{3}$

$$= \log \frac{2^3}{3}$$

$$= \log 2^3 - \log 3$$

$$= 3 \log 2 - \log 3 = 3(0.3010) - 0.4771$$

$$= 0.9030 - 0.4771 = 0.4259$$

5(v) $\log 30$

$$= \log 2 \times 3 \times 5$$

$$= \log 2 + \log 3 + \log 5$$

$$= 0.3010 + 0.4771 + 0.6990$$

$$= 1.4771$$

MATHEMATICS FOR 9TH CLASS (UNIT # 3)

Print Superone Mathematics 172 Class 9th

Exercise 3.4

Q.1. Use log tables to find the value of:

- | | |
|---|---|
| (i) 0.8176×13.64 | (ii) $(789.5)^{1/8}$ |
| (iii) $\frac{0.678 \times 9.01}{0.0234}$ | (iv) $\sqrt[3]{2709} \times \sqrt[4]{1239}$ |
| (v) $\frac{(1.23)(0.6975)}{(0.0075)(1278)}$ | (vi) $\sqrt[3]{\frac{0.7214 \times 20.37}{60.8}}$ |
| (vii) $\frac{83 \times \sqrt[3]{92}}{127 \times \sqrt[5]{246}}$ | (viii) $\frac{(438)^3 \sqrt{0.056}}{(388)^4}$ |

I(i) 0.8176×13.64

Let $x = 0.8176 \times 13.64$

$$\log x = \log(8176 \times 13.64) \text{ (Taking log)}$$

$$= \log 8176 + \log 13.64$$

$$= \bar{1}.9125 + 1.1348$$

$$\log x = 1.0473$$

$$\text{Antilog}(\log x) = \text{Antilog } 1.0473 \text{ (Taking anti-log)}$$

$$x = 11.15$$

I(ii) $(789.5)^{1/8}$

Let $x = (789.5)^{1/8}$

$$\log x = \log (789.5)^{1/8}$$

$$= \frac{1}{8} \log 789.5$$

$$= \frac{1}{8} \{2.8974\} = 0.3622$$

$$\text{Antilog}(\log x) = \text{Antilog}(0.3622) = 2.302$$

I(iii) $\frac{0.678 \times 9.01}{0.0234}$

Let $x = \frac{0.678 \times 9.01}{0.0234}$

MATHEMATICS FOR 9TH CLASS (UNIT # 3)

Pilot Superone Mathematics 173 Class 9th

$$\begin{aligned} \log x &= \log \frac{0.678 \times 9.01}{0.0234} \quad \text{Taking log} \\ &= \log 0.678 + \log 9.01 - \log 0.0234 \\ &= \bar{1} 8312 + 0.9547 - (\bar{2} 3692) \\ &= \bar{1} 8312 + 0.9547 - \bar{2} - 3692 \\ &= -1 + 8312 + 0.9547 + 2 - 3692 = 2.4167 \\ \text{Antilog}(\log x) &= \text{Antilog } 2.4167 \quad \text{Taking anti-log} \\ x &= 261 \end{aligned}$$

$$\begin{aligned} \text{I(iv)} \quad & \sqrt[5]{2.709} \times \sqrt[7]{1.239} \\ \text{Let } x &= \sqrt[5]{2.709} \times \sqrt[7]{1.239} \\ x &= (2.709)^{1/5} \times (1.239)^{1/7} \quad \text{(Taking log)} \\ \log x &= \log (2.709)^{1/5} \times (1.239)^{1/7} \\ &= \frac{1}{5} \log 2.709 + \frac{1}{7} \log 1.239 \\ &= \frac{1}{5} (0.4328) + \frac{1}{7} (0.0931) \\ &= 0.0866 + 0.0133 \\ \log x &= 0.0999 \quad \text{(Taking anti-log)} \\ \text{Antilog}(\log x) &= \text{Antilog } 0.0999 \\ x &= 1.258 \end{aligned}$$

$$\begin{aligned} \text{I(v)} \quad & \frac{(1.23)(0.6975)}{(0.0075)(1278)} \\ \text{Let } x &= \frac{(1.23)(0.6975)}{(0.0075)(1278)} \\ \log x &= \log \frac{(1.23)(0.6975)}{(0.0075)(1278)} \quad \text{(Taking log)} \\ \log x &= \log 1.23 + \log 0.6975 - \log 0.0075 - \log 1278 \\ &= 0.0899 + \bar{1} 8435 - (\bar{3} 8751) - (3 1065) \\ &= 0.0899 - 1 + 8435 + 3 - 8751 - 3 1065 \\ 1.0482 &= -1.0482 \quad -1 + 1 \end{aligned}$$

MATHEMATICS FOR 9TH CLASS (UNIT # 3)

Pilot Superone Mathematics 174 Class 9th

$$\log x = \bar{2} \ 9518$$

$$\text{Antilog}(\log x) = \text{Antilog } \bar{2} \ 9518 \quad (\text{Taking anti-log})$$

$$x = 0895$$

$$1(\text{vi}) \quad \sqrt[3]{\frac{0.7214 \times 20.37}{60.8}}$$

$$\text{Let } x = \sqrt[3]{\frac{0.7214 \times 20.37}{60.8}}$$

$$x = \log\left(\frac{0.7214 \times 20.37}{60.8}\right)^{1/3} \quad (\text{Taking log})$$

$$\log x = \frac{1}{3} [\log 7214 + \log 20.37 - \log 60.8]$$

$$= \frac{1}{3} [1 \ 8582 + 1 \ 3090 - 1 \ 7839]$$

$$= \frac{1}{3} [-1 + 8582 + 1 \ 3090 - 1 \ 7839]$$

$$= \frac{1}{3} [-0.6167] = -0.2056$$

$$= -0.2056 - 1 + 1 \ \bar{1} \ 7944$$

$$\text{Antilog}(\log x) = \text{Antilog } \bar{1} \ 7944 \quad (\text{Taking anti-log})$$

$$x = 6229$$

$$1(\text{vii}) \quad \frac{83 \times \sqrt[3]{92}}{127 \times \sqrt[3]{246}}$$

$$\text{Let } x = \frac{83 \times \sqrt[3]{92}}{127 \times \sqrt[3]{246}}$$

$$x = \frac{83 \times (92)^{1/3}}{127 \times (246)^{1/3}} \quad (\text{Taking log})$$

$$\log x = \log 83 + \frac{1}{3} \log 92 - \log 127 - \frac{1}{3} \log$$

MATHEMATICS FOR 9TH CLASS (UNIT # 3)

Pilot Superone Mathematics 175 Class 9th

$$\approx 1\ 9191 + \frac{1}{3}(1\ 9638) \quad (2\ 1038) - \frac{1}{5}(2\ 3909)$$

$$= 1\ 9191 + 0\ 6546 \quad 2\ 1038 \quad 0\ 4782$$

$$2\ 5737 \quad 2\ 582 = 0083 = -0083 - 1 + 1$$

$$\log x \quad \bar{1}\ 9917$$

$$\text{Antilog}(\log x) = \text{Antilog } \bar{1}\ 9917 \quad (\text{Taking anti-log})$$

$$x \quad 9811$$

$$I(\text{viii}) \quad \frac{(438)^3 \sqrt{0.056}}{(388)^4}$$

$$\text{Let } x = \frac{(438)^3 \sqrt{0.056}}{(388)^4}$$

$$x = \frac{(438)^3 \times (0.056)^{\frac{1}{2}}}{(388)^4}$$

$$\log x = 3 \log 438 + \frac{1}{2} \log(0.056) - 4 \log 388 \quad (\text{Taking log})$$

$$= 3(2.6415) + \frac{1}{2}(2\ 7482) - 4(2\ 5888)$$

$$7\ 9245 + \bar{1}\ 3741 \quad 10\ 3552$$

$$= 7\ 9245 \quad 1 + 3741 \quad 10\ 3552$$

$$\log x = -3\ 0566 = -3\ 0566 - 1 + 1$$

$$\bar{4}\ 9434$$

$$\text{Antilog}(\log x) = \text{Antilog } \bar{4}\ 9434 \quad (\text{Taking anti-log})$$

$$x = 0.0008778$$

Q.2. A gas is expanding according to the law $pv^n = C$,

find C when $p = 80$, $v = 3.1$ and $n = \frac{5}{4}$

Solution:-

$$C = pv^n$$

MATHEMATICS FOR 9TH CLASS (UNIT # 3)

Pilot Superone Mathematics 176 Class 9th

Putting values of p, v, n

$$C = 80(3.1)^{3.4}$$

$$\log C = \log 80 + \frac{5}{4} \log 3.1 \quad (\text{Taking log})$$

$$= 1.9031 + \frac{5}{4} (0.4970)$$

$$= 1.9031 + \frac{2.4850}{4}$$

$$= 1.9031 + 0.6143$$

$$\log C = 2.5174$$

$$\text{Antilog}(\log C) = \text{Antilog } 2.5174 \quad (\text{Taking anti-log})$$

$$C = 329.2$$

Q.3. The formula $p = 90(5)^{-q/10}$ applies to the demand of a product, where q is the number of units and p is the price of one unit. How many units will be demanded if the price is Rs 18.00?

Solution:-

$$p = 90(5)^{-q/10}$$

$$18 = \log 90(5)^{-q/10}$$

Putting $p = 18.00$

$$\log 18 = \log 90(5)^{-q/10} \quad (\text{Taking log})$$

$$= \log 90 + \log 5^{-q/10}$$

$$\log 18 = \log 90 - \frac{q}{10} \log 5$$

$$1.2552 = 1.9542 - \frac{q}{10} (0.6990)$$

$$= 1.9542 - 0.0699q$$

$$0.699q = 1.9542 - 1.2552$$

$$0.699q = 0.6990$$

$$q = \frac{0.6990}{0.699} = 1$$

10 Units

MATHEMATICS FOR 9TH CLASS (UNIT # 3)

Pilot Superiore Mathematics 177 Class 9th

Q.4. If $A = \pi r^2$, find A , when $\pi = \frac{22}{7}$ and $r = 15$.

Solution:-

$$A = \pi r^2$$

$$\text{Putting } \pi = \frac{22}{7} \text{ and } r = 15$$

$$A = \frac{22}{7} \times (15)^2 \quad (\text{Taking log})$$

$$\begin{aligned} \log A &= \log \frac{22}{7} + 2 \log 15 \\ &= 1.3424 - 0.8451 + 2(1.1761) \\ &= 1.3424 - 0.8451 + 2.3522 \end{aligned}$$

$$\log A = 2.8494$$

$$\text{Antilog } (\log A) = \text{Antilog } 2.8494 \quad (\text{Taking anti-log})$$

$$A = 7071$$

Q.5. If $V = \frac{1}{3} \pi r^2 h$, find V , when $\pi = \frac{22}{7}$, $r = 2.5$ and $h = 4.2$

Solution:-

$$V = \frac{1}{3} \pi r^2 h$$

Putting values of h , r & π

$$V = \frac{1}{3} \times \frac{22}{7} \times (2.5)^2 \times 4.2$$

$$\log V = \log \frac{1}{3} + \log \frac{22}{7} + 2 \log 2.5 + \log 4.2 \quad (\text{Taking log})$$

$$\begin{aligned} \log V &= \log 22 - 2 \log 2.5 + \log 4.2 - \log 3 - \log 7 \\ &= 1.3424 - 2(0.3979) + 0.6232 - 0.4771 - 0.8451 \\ &= 1.3424 - 0.7958 + 0.6232 - 0.4771 - 0.8451 \end{aligned}$$

$$\log V = 1.4392$$

$$\text{Antilog } (\log V) = \text{Antilog } 1.4392 \quad (\text{Taking anti-log})$$

$$V = 27.49$$

Unit

4

**ALGEBRAIC EXPRESSIONS AND
ALGEBRAIC FORMULAS**

Types of Algebraic Expressions

- (i) Polynomial expressions
- (ii) Rational Expression
- (iii) Irrational expression

(i) Polynomial expressions.

A polynomial consists of one or more than one terms. The exponent of variable or variables is zero or positive integer. For example $12 - x$, $3x - 4y$, $x^2 - 2x + 1$, $\sqrt{3}x$ are polynomial expressions, where as $\frac{2}{x^3}$, $x + \frac{1}{x}$, $\sqrt{3}x - \frac{1}{\sqrt{3}x}$ and

$35x - \frac{6}{x}$ are not polynomials.

(ii) Rational Expression.

$\frac{p(x)}{q(x)}$ is a rational expression where as $p(x)$, $q(x)$ are polynomials and $q(x) \neq 0$.

(iii) Irrational expression.

If an algebraic expression cannot be written in the form $\frac{p(x)}{q(x)}$ where $p(x)$, $q(x)$ are polynomials and $q(x) \neq 0$ is called

an irrational expression. For example \sqrt{x} , x^2 , $\sqrt[3]{x^2y^3}$ and $\sqrt{x+1}$, $\frac{1}{\sqrt{x+2}}$ are irrational expressions.

Pilot Superone Mathematics 186 Class 9th

Exercise 4.1

Q.1 Identify whether the following algebraic expressions are polynomials (Yes or No)

- (i) $3x^2 + \frac{1}{x} - 5$ (ii) $3x^3 - 4x^2 - x\sqrt{x} + 3$
 (iii) $x^2 - 3x + \sqrt{2}$ (iv) $\frac{3x}{2x-1} + 8$

Solutions:

- 1(i)** $3x^2 + \frac{1}{x} - 5$ **No** Reason $\frac{1}{x}$
1(ii) $3x^3 - 4x^2 - x\sqrt{x} + 3$ **No** Reason \sqrt{x}
1(iii) $x^2 - 3x + \sqrt{2}$ **Yes**
1(iv) $\frac{3x}{2x-1} + 8$ **No** Reason $\frac{1}{2x-1}$

Q.2 State whether each of the following expression is a rational expression or not.

- (i) $\frac{3\sqrt{x}}{3\sqrt{x}+5}$ (ii) $\frac{x^3 - 2x^2 + \sqrt{3}}{2+3x-x^2}$
 (iii) $\frac{x^2+6x+9}{x^2-9}$ (iv) $\frac{2\sqrt{x}+3}{2\sqrt{x}-3}$

Solutions:

2(i) $\frac{3\sqrt{x}}{3\sqrt{x}+5}$

Since $3\sqrt{x}$ and $3\sqrt{x}+5$ are not polynomials therefore, the given expression is not an algebraic rational

MATHEMATICS FOR 9TH CLASS (UNIT # 4)

Pilot Superone Mathematics 187 Class 9th

$$2(ii) \quad \frac{x^3 - 2x + \sqrt{3}}{2 + 3x - x^2}$$

It is rational expressions as $x^3 - 2x + \sqrt{3}$ and $2 + 3x - x^2$ are polynomials

$$2(iii) \quad \frac{x^2 + 6x + 9}{x^2 - 9}$$

It is a rational expression. Both denominator and numerator are polynomials

$$2(iv) \quad \frac{2\sqrt{x} + 3}{2\sqrt{x} - 3}$$

It is not a rational expression $2\sqrt{x} + 3$ and $2\sqrt{x} - 3$ are not polynomials

Q.3 Reduce the following rational expressions to the lowest forms.

$$(i) \quad \frac{120x^2y^3z^4}{30x^3yz^2}$$

$$(ii) \quad \frac{8a(x+1)}{2(x^2-1)}$$

$$(iii) \quad \frac{(x+y)^2 - 4xy}{(x-y)^2}$$

(iv)

$$\frac{(x^2 - y^2)(x^2 - 2xy + y^2)}{(x - y)(x^2 + xy + y^2)}$$

$$(v) \quad \frac{(x+2)(x^2-1)}{(x+1)(x^2-4)}$$

$$(vi) \quad \frac{x^2 - 4x + 4}{2x^2 - 8}$$

$$(vii) \quad \frac{64x^3 - 64x}{(8x^2 + 8)(2x + 2)}$$

$$(viii) \quad \frac{9x^3 - (x^2 - 4)^2}{4 + 3x - x^2}$$

Solutions:

$$3(i) \quad \frac{120x^2y^3z^4}{30x^3yz^2}$$

MATHEMATICS FOR 9TH CLASS (UNIT # 4)

Pilot Superone Mathematics 188 Class 9th

$$\frac{30x + 4x^{3-1}z^{5-2}}{30x^{3-1}z^{5-2}} = \frac{4x^2}{x^2}$$

$$3(ii) \quad \frac{8a(x+1)}{2(x^2-1)} \\ = \frac{2 \times 4a(x+1)}{2(x+1)(x-1)} = \frac{4a}{x-1}$$

$$3(iii) \quad \frac{(x+y)^2 - 4xy}{(x-y)^2} \\ = \frac{x^2 + y^2 + 2xy - 4xy}{(x-y)^2} \\ = \frac{x^2 - 2xy + y^2}{(x-y)^2} = 1$$

$$3(iv) \quad \frac{(x^2 - y^2)(x^2 - 2xy + y^2)}{(x^2 + y^2)(x^2 + 2xy + y^2)} \\ = \frac{(x-y)(x+y)(x-y)^2}{(x+y)(x+y)^2} \\ = \frac{(x-y)^2}{(x+y)^2} = \left(\frac{x-y}{x+y}\right)^2$$

$$3(v) \quad \frac{(x+2)(x^2-1)}{(x+1)(x^2-4)} \\ = \frac{(x+2)(x^2-1^2)}{(x+1)(x^2-2^2)} \\ = \frac{(x+2)(x+1)(x-1)}{(x+1)(x+2)(x-2)} = \frac{x-1}{x-2}$$

$$3(vi) \quad \frac{x^2 - 4x + 4}{2x^2 - 8} \\ = \frac{x^2 - 2x - 2x + 4}{2x^2 - 8}$$

MATHEMATICS FOR 9TH CLASS (UNIT # 4)

Pilot Superone Mathematics 189 Class 9th

$$\frac{x(x-2)-2(x-2)}{2(x-2)^2}$$

$$= \frac{(x-2)(x-2)}{2(x-2)(x+2)}$$

$$= \frac{x-2}{2(x+2)}$$

3(vii) $\frac{64x^3 - 64x}{(8x^3 + 8)(2x + 2)}$

$$= \frac{64x(x^2 - 1)}{8(x^3 + 1)(2)(x + 1)}$$

$$= \frac{64x[(x^2)^2 - (1)^2]}{16(x^3 + 1)(x + 1)}$$

$$= \frac{64x(x^2 + 1)(x^2 - 1)}{16(x^3 + 1)(x + 1)}$$

$$= \frac{4x(x^2 + 1)(x^2 - 1)}{(x^3 + 1)(x + 1)}$$

$$4x(x^2 - 1)$$

3(viii) $\frac{9x^3 - (x^2 - 4)^2}{4 + 3x - x^2}$

$$= \frac{(3x)^2 - (x^2 - 4)}{4 + 3x - x^2}$$

$$= \frac{[3x + (x^2 - 4)][3x - (x^2 - 4)]}{(4 + 3x - x^2)}$$

$$= \frac{(x^2 + 3x - 4)(3x - x^2 + 4)}{(4 + 3x - x^2)}$$

$$= \frac{(x^2 + 3x - 4)(4 + 3x - x^2)}{(4 + 3x - x^2)}$$

MATHEMATICS FOR 9TH CLASS (UNIT # 4)

Pilot Superone Mathematics 190 Class 9th

$$x^2 + 3x - 4$$

Q4 Evaluate (a) $\frac{x^3y - 2z}{xz}$ for

(i) $x = 3, y = -1, z = 2$ (ii) $x = -1, y = -9, z = 4$

(b) $\frac{x^3y - 5z^4}{xyz}$ for $x = 4, y = -2, z = 1$

Solutions:

4(a)(i) $\frac{x^3y - 2z}{xz}$

Putting $x = 3, y = -1, z = 2$

$$= \frac{(3)^3(-1) - 2(-2)}{(3)(2)}$$

$$= \frac{(27)(-1) + 4}{6}$$

$$= \frac{-27 + 4}{6} = \frac{-23}{6}$$

$$= \frac{-23}{6} = -\frac{23}{6}$$

4(a)(ii) $\frac{x^3y - 2z}{xz}$

Putting $x = -1, y = -9, z = 4$

$$\frac{(-1)^3(-9) - 2(4)}{(-1)(4)}$$

$$= \frac{(-1)(-9) - 8}{-4}$$

$$= \frac{9 - 8}{-4} = -\frac{1}{4}$$

MATHEMATICS FOR 9TH CLASS (UNIT # 4)

Pilot Superone Mathematics 191 Class 9th

4(b) $\frac{x^3y^3 - 5x^3}{xy^3}$

Putting $x = 4, y = -2, z = 1$

$$= \frac{(4)^3(-2)^3 - 5(-1)^3}{(4)(-2)(-1)}$$

$$= \frac{(16)(-8) - 5(1)}{8}$$

$$= \frac{-128 - 5}{8}$$

$$= \frac{-133}{8} = -16\frac{5}{8}$$

Q.5 Perform the indicated operation and simplify.

(i) $\frac{15}{2x-3y} - \frac{4}{3y-2x}$

(ii) $\frac{1+2x}{1-2x} - \frac{1-2x}{1+2x}$

(iii) $\frac{x^2-25}{x^2-36} - \frac{x+5}{x+6}$

(iv)

$$\frac{x}{x-y} - \frac{y}{x+y} + \frac{2xy}{x^2-y^2}$$

(v) $\frac{x+2}{x^2+6x+9} - \frac{x+2}{2x^2-18}$

(vi) $\frac{1}{x-1} - \frac{1}{x+1} - \frac{2}{x^2+1} - \frac{4}{x^4-1}$

Solutions:

5(i) $\frac{15}{2x-3y} - \frac{4}{3y-2x}$

$$= \frac{15}{2x-3y} + \frac{4}{2x-3y}$$

$$= \frac{15+4}{2x-3y}$$

(N.T.S)

MATHEMATICS FOR 9TH CLASS (UNIT # 4)

Pilot Superone Mathematics 192 Class 9th

$$= \frac{19}{2x - 3y}$$

$$\begin{aligned} 5(ii) \quad & \frac{1+2x}{1-2x} - \frac{1-2x}{1+2x} \\ &= \frac{(1+2x)(1+2x) - (1-2x)(1-2x)}{(1-2x)(1+2x)} \\ &= \frac{(1+4x+4x^2) - (1-4x+4x^2)}{(1)^2 - (2x)^2} \\ &= \frac{1+4x+4x^2 - 1+4x-4x^2}{1-4x^2} \\ &= \frac{8x}{1-4x^2} \end{aligned}$$

$$\begin{aligned} 5(iii) \quad & \frac{x-25}{x-36} - \frac{x+5}{x+6} \\ &= \frac{(x-25)(x+6) - (x+5)(x-36)}{(x-36)(x+6)} \\ &= \frac{(x^2-25x+150) - (x^2-30x+180)}{(x-36)(x+6)} \\ &= \frac{x^2-25x+150-x^2+30x-180}{(x-36)(x+6)} \\ &= \frac{5x-30}{(x-36)(x+6)} \\ &= \frac{5(x-6)}{(x-36)(x+6)} \\ &= \frac{5}{x+6} \end{aligned}$$

MATHEMATICS FOR 9TH CLASS (UNIT # 4)

Pilot Superone Mathematics

193

Class 9th

$$\begin{aligned}
 S(iv) \quad & \frac{x}{x-y} - \frac{y}{x+y} - \frac{2xy}{x^2-y^2} \\
 &= \frac{x}{x-y} - \frac{y}{x+y} - \frac{2xy}{(x-y)(x+y)} \\
 &= \frac{x(x+y) - y(x-y) - 2xy}{(x-y)(x+y)} \\
 &= \frac{x^2 + xy - xy + y^2 - 2xy}{(x-y)(x+y)} \\
 &= \frac{x^2 + y^2 - 2xy}{(x-y)(x+y)} \\
 &= \frac{(x-y)^2}{(x-y)(x+y)} \\
 &= \frac{x-y}{x+y}
 \end{aligned}$$

$$\begin{aligned}
 S(v) \quad & \frac{x-2}{x^2+6x+9} - \frac{x+2}{2x^2-18} \\
 &= \frac{x-2}{x^2+3x+3x+9} - \frac{x+2}{2(x^2-9)} \\
 &= \frac{x-2}{x(x+3)+3(x+3)} - \frac{x+2}{2(x^2-3^2)} \\
 &= \frac{x-2}{(x+3)(x+3)} - \frac{x+2}{2(x+3)(x-3)} \\
 &= \frac{2(x-2)(x-3) - (x+2)(x+3)}{2(x+3)(x+3)(x-3)} \\
 &= \frac{2x^2 - 10x + 12 - x^2 - 5x - 6}{2(2x+3)^2(x-3)}
 \end{aligned}$$

Pilot Superone Mathematics 194 Class 9th

$$\begin{aligned}
 &= \frac{x^2 - 15x + 6}{2(x+3)(x-3)} \\
 5(vi) \quad &\frac{1}{x-1} - \frac{1}{x+1} - \frac{2}{x^2+1} - \frac{4}{x^4-1} \\
 &= \left(\frac{1}{x-1} - \frac{1}{x+1} \right) - \left(\frac{2}{x^2+1} + \frac{4}{(x^2)^2-1^2} \right) \\
 &= \left(\frac{x+1-x-1}{(x-1)(x+1)} \right) - \left(\frac{2}{x^2+1} + \frac{4}{(x^2+1)(x^2-1)} \right) \\
 &= \frac{2x}{x^2-1} - \left(\frac{2(x^2-1)+4}{(x^2+1)(x^2-1)} \right) \\
 &= \frac{2x}{x^2-1} - \frac{2x^2-2+4}{(x^2+1)(x^2-1)} \\
 &= \frac{2x}{x^2-1} - \frac{2x^2+2}{(x^2+1)(x^2-1)} \\
 &= \frac{2x}{x^2-1} - \frac{2(x^2+1)}{(x^2+1)(x^2-1)} \\
 &= \frac{2x}{x^2-1} - \frac{2}{x^2-1} = 0
 \end{aligned}$$

Q.6 Perform the indicated operation and simplify

$$\begin{aligned}
 (i) \quad &(x^2 - 49) \cdot \frac{5x+2}{x+7} & (ii) \quad &\frac{4x-12}{x^2-9} + \frac{18-2x^2}{x^2+6x+9} \\
 (iii) \quad &\frac{x^4-y^6}{x^3-y^3} + (x^4+x^2y^2+y^4) & (iv) \quad &\frac{x^2-1}{x^2+2x+1} \cdot \frac{x+5}{1-x}
 \end{aligned}$$

MATHEMATICS FOR 9TH CLASS (UNIT # 4)

Pilot Superone Mathematics 195

Class 9th

$$(v) \quad \frac{x^2 + xy}{y(x+y)} \cdot \frac{x^2 + xy}{y(x+y)} \div \frac{x^2 - x}{xy - 2y}$$

Solutions:

$$6(i) \quad (x^2 - 49) \cdot \frac{5x+2}{x+7}$$

$$(x^2 - 7^2) \times \frac{5x+2}{x+7}$$

$$= (x - 7)(x + 7) \times \frac{5x+2}{x+7}$$

$$= (x - 7)(5x + 2)$$

$$= 5x^2 - 33x - 14$$

$$6(ii) \quad \frac{4x-12}{x^2-9} \div \frac{18-2x^2}{x^2+6x+9}$$

$$= \frac{4(x-3)}{x^2-3^2} \div \frac{2(9-x^2)}{x^2+3x+3x+9}$$

$$= \frac{4(x-3)}{(x-3)(x+3)} \div \frac{2(3^2-x^2)}{x(x+3)+3(x+3)}$$

$$= \frac{4}{x+3} \div \frac{2(3+x)(3-x)}{(x+3)(x+3)}$$

$$= \frac{4}{x+3} \div \frac{2(3-x)}{x+3}$$

$$= \frac{4}{x+3} \times \frac{x+3}{2(3-x)}$$

$$= \frac{2}{3-x}$$

$$x^6 - y^6$$

MATHEMATICS FOR 9TH CLASS (UNIT # 4)

Pilot Superone Mathematics

196

Class 9th

$$= \frac{(x^2)^3 - (y^2)^3}{x^2 - y^2} \div (x^4 + x^2y^2 + y^4)$$

$$= \frac{(x^2 - y^2)(x^4 + x^2y^2 + y^4)}{(x^2 - y^2)} \times \frac{1}{x^4 + x^2y^2 + y^4}$$

$$= 1$$

6(iv) $\frac{x^2 - 1}{x^2 + 2x + 1} \cdot \frac{x + 5}{1 - x}$

$$= \frac{x^2 - 1^2}{x^2 + x + x + 1} \times \frac{x + 5}{-(x - 1)} \quad \text{N.T.S}$$

$$= \frac{(x - 1)(x + 1)}{x(x + 1) + 1(x + 1)} \times \frac{-(x + 5)}{(x - 1)}$$

$$= \frac{(x - 1)(x + 1)}{(x + 1)(x + 1)} \times \frac{-(x + 5)}{(x - 1)}$$

$$= \frac{(x + 5)}{x + 1}$$

6(v) $\frac{x^2 + xy}{y(x + y)} \cdot \frac{x^2 + xy}{y(x + y)} \div \frac{x^2 - x}{xy - 2y}$

$$= \frac{x(x + y)}{y(x + y)} \times \frac{x(x + y)}{y(x + y)} \times \frac{xy - 2y}{x^2 - x}$$

$$= \frac{x(x + y)}{y(x + y)} \times \frac{x(x + y)}{y(x + y)} \times \frac{y(x - 2)}{x(x - 1)}$$

$$= \frac{x(x - 2)}{y(x - 1)}$$

MATHEMATICS FOR 9TH CLASS (UNIT # 4)

Pilot Superone Mathematics

197

Class 9th

Algebraic Formulae

$$(i) \quad (a + b)^2 = a^2 + 2ab + b^2$$

$$(ii) \quad (a - b)^2 = a^2 - 2ab + b^2$$

$$(iii) \quad (a + b)^2 + (a - b)^2 = 2(a^2 + b^2)$$

$$(iv) \quad (a + b)^2 - (a - b)^2 = 4ab$$

$$(v) \quad (a + b + c)^2 = a^2 + b^2 + c^2 + 2ab + 2bc + 2ca$$

$$= a^2 + b^2 + c^2 + 2(ab + bc + ca)$$

$$(vi) \quad (a + b)^3 = a^3 + 3a^2b + 3ab^2 + b^3$$

$$= a^3 + b^3 + 3ab(a + b)$$

$$(vii) \quad (a - b)^3 = a^3 - 3a^2b + 3ab^2 - b^3$$

$$= a^3 - b^3 - 3ab(a - b)$$

$$(viii) \quad a^3 - b^3 = (a + b)(a^2 - ab + b^2)$$

$$= (a - b)(a^2 + ab + b^2)$$

Exercise 4.2

Q.1(i) If $a + b = 10$ and $a - b = 6$, then find the value of $a^2 + b^2$

I(ii) If $a + b = 5$ and $a - b = \sqrt{17}$ then find the value of ab .

1(i) If $a + b = 10$ and $a - b = 6$, then find the value of $a^2 + b^2$

Solutions:

$$\begin{aligned} a + b &= 10 && \text{Given} \\ a - b &= 6 \\ a^2 + b^2 &= ? \end{aligned}$$

We know that

$$2(a^2 + b^2) = (a + b)^2 + (a - b)^2$$

Putting value of $a + b$ and $a - b$

$$= (10)^2 + (6)^2$$

$$= 100 + 36$$

$$2(a^2 + b^2) = 136$$

$$a^2 + b^2 = \frac{136}{2} \quad (\text{Dividing by 2})$$

$$\therefore a^2 + b^2 = 68$$

I(ii) If $a + b = 5$ and $a - b = \sqrt{17}$ then find the value of ab .

Solution:

$$\begin{aligned} a + b &= 5 && \text{Given} \\ a - b &= \sqrt{17} \\ ab &= ? \end{aligned}$$

We know that

$$(a + b)^2 - (a - b)^2 = 4ab$$

Putting values of $a + b$ and $a - b$

$$(5)^2 - (\sqrt{17})^2 = 4ab$$

$$25 - 17 = 4ab$$

$$8 = 4ab$$

MATHEMATICS FOR 9TH CLASS (UNIT # 4)

Pilot Superone Mathematics 199 Class 9th

or $4ab = 8$

or $ab = \frac{8}{4}$

$ab = 2$

Q.2 If $a^2 + b^2 + c^2 = 45$ and $a + b + c = -1$, find the value of $ab + bc + ca$

Solution: $a^2 + b^2 + c^2 = 45$ Given

$a + b + c = 1$

$ab + bc + ca = ?$

$(a + b + c)^2 = (a^2 + b^2 + c^2) + 2(ab + bc + ca)$

Putting $a + b + c = -1$, $a^2 + b^2 + c^2 = 45$

$(-1)^2 = 45 + 2(ab + bc + ca)$

$1 - 45 = 2(ab + bc + ca)$

$-44 = 2(ab + bc + ca)$

$-22 = ab + bc + ca$ (Dividing by 2)

or $ab + bc + ca = -22$

Q.3 If $m + n + p = 10$ and $mn + np + mp = 27$, find the value of $m^2 + n^2 + p^2$

Solution: $m + n + p = 10$ Given

$mn + np + mp = 27$

$m^2 + n^2 + p^2 = ?$

Formula

$(m + n + p)^2 = m^2 + n^2 + p^2 + 2(mn + np + mp)$

Putting $mn + np + mp = 27$, $m + n + p = 10$

$(10)^2 = m^2 + n^2 + p^2 + 2(27)$

$100 = m^2 + n^2 + p^2 + 54$

or $m^2 + n^2 + p^2 = 100 - 54$
 $= 46$

MATHEMATICS FOR 9TH CLASS (UNIT # 4)

Pilot Superone Mathematics 200 Class 9th

Q.4 If $x^2 + y^2 + z^2 = 78$ and $xy + yz + zx = 59$, Find the value of $x + y + z$

Solution: $x^2 + y^2 + z^2 = 78$ Given

$$xy + yz + zx = 59$$

$$x + y + z = ?$$

Formula

$$(x + y + z)^2 = x^2 + y^2 + z^2 + 2(xy + yz + zx)$$

Putting $xy + yz + zx = 59$, $x^2 + y^2 + z^2 = 78$

$$(x + y + z)^2 = 78 + 2(59)$$

$$= 78 + 118$$

$$(x + y + z)^2 = 196$$

$$x + y + z = \pm\sqrt{196} \quad (\text{Taking square root})$$

$$x + y + z = \pm 14$$

Q.5 If $x + y + z = 12$ and $x^2 + y^2 + z^2 = 64$, find the value of $xy + yz + zx$

Solution: $x + y + z = 12$ Given

$$x^2 + y^2 + z^2 = 64$$

$$xy + yz + zx = ?$$

Formula

$$(x + y + z)^2 = x^2 + y^2 + z^2 + 2(xy + yz + zx)$$

Putting value of $x + y + z = 12$, $x^2 + y^2 + z^2 = 64$

$$(12)^2 = 64 + 2(xy + yz + zx)$$

$$144 = 64 + 2(xy + yz + zx)$$

$$144 - 64 = 2(xy + yz + zx)$$

$$80 = 2(xy + yz + zx)$$

Thus $2(xy + yz + zx) = 80$

$$xy + yz + zx = 40 \quad (\text{Dividing by 2})$$

MATHEMATICS FOR 9TH CLASS (UNIT # 4)

Pilot Superone Mathematics 201 Class 9th

Q.6 If $x + y = 7$ and $xy = 12$, then find the value of $x^3 + y^3$.

Solution: $x + y = 7$ Given
 $xy = 12$
 $x^3 + y^3 = ?$

Formula

$$(x + y)^3 = x^3 + y^3 + 3xy(x + y)$$

Putting value of $x + y = 7$, $xy = 12$

$$(7)^3 = x^3 + y^3 + 3(12)(7)$$

$$343 = x^3 + y^3 + 252$$

$$343 - 252 = x^3 + y^3$$

$$91 = x^3 + y^3$$

or $x^3 + y^3 = 91$

Q.7 If $3x + 4y = 11$ and $xy = 12$, then find the value of $27x^3 + 64y^3$.

Solution: $3x + 4y = 11$ Given
 $xy = 12$
 $27x^3 + 64y^3 = ?$

Formula

$$(3x + 4y)^3 = (3x)^3 + (4y)^3 + 3(3x)(4y)(3x + 4y)$$

Putting value of $3x + 4y = 11$, $xy = 12$

$$(11)^3 = 27x^3 + 64y^3 + 3 \times 3 \times 4(xy)(11)$$

$$1331 = 27x^3 + 64y^3 + 3 \times 3 \times 4 \times 12 \times 11$$

$$1331 = 27x^3 + 64y^3 + 4752$$

$$1331 - 4752 = 27x^3 + 64y^3$$

$$-3421 = 27x^3 + 64y^3$$

or $27x^3 + 64y^3 = -3421$

Q.8 If $x - y = 4$ and $xy = 21$, then find the value of $x^3 - y^3$

Solution: $x - y = 4$ Given
 $xy = 21$
 $x^3 - y^3 = ?$

MATHEMATICS FOR 9TH CLASS (UNIT # 4)

Pilot Superone Mathematics 202 Class 9th
 Formula

$$(x - y)^3 = x^3 - y^3 - 3xy(x - y)$$

Putting value of $x - y = 4$, $xy = 21$

$$(4)^3 = x^3 - y^3 - 3(21)(4)$$

$$64 = x^3 - y^3 - 252$$

$$64 + 252 = x^3 - y^3$$

$$316 = x^3 - y^3$$

or $x^3 - y^3 = 316$

Q.9 If $5x - 6y = 13$ and $xy = 6$, then find the value of $125x^3 - 216y^3$

Solution: $5x - 6y = 13$ Given
 $xy = 6$
 $125x^3 - 216y^3 = ?$

Formula

$$(5x - 6y)^3 = (5x)^3 - (6y)^3 - 3(5x)(6y)(5x - 6y)$$

Putting value of $5x - 6y = 13$, $xy = 6$

$$(13)^3 = 125x^3 - 216y^3 - 3 \times 5 \times 6 \times xy \times (13)$$

$$2197 = 125x^3 - 216y^3 - 1170 \times 6$$

$$2197 = 125x^3 - 216y^3 - 7020$$

$$2197 + 7020 = 125x^3 - 216y^3$$

$$9217 = 125x^3 - 216y^3$$

or $125x^3 - 216y^3 = 9217$

Q.10 If $x + \frac{1}{x} = 3$, then find the value of $x^3 + \frac{1}{x^3}$

Solution: $x + \frac{1}{x} = 3$ Given

$$x^3 + \frac{1}{x^3} = ?$$

Formula

MATHEMATICS FOR 9TH CLASS (UNIT # 4)

Pilot Superone Mathematics

203

Class 9th

$$\left(x + \frac{1}{x}\right)^3 = x^3 + \frac{1}{x^3} + 3(x) \left(\frac{1}{x}\right) \left(x + \frac{1}{x}\right)$$

Putting value of $x + \frac{1}{x} = 3$

$$(3)^3 = x^3 + \frac{1}{x^3} + 3(3)$$

$$27 = x^3 + \frac{1}{x^3} + 9$$

$$27 - 9 = x^3 + \frac{1}{x^3}$$

$$18 = x^3 + \frac{1}{x^3}$$

or

$$x^3 + \frac{1}{x^3} = 18$$

Q.11 If $x - \frac{1}{x} = 7$, then find the value of $x^3 - \frac{1}{x^3}$

Solution: $x - \frac{1}{x} = 7$ Given

$$x^3 - \frac{1}{x^3} = ?$$

Formula

$$\left(x - \frac{1}{x}\right)^3 = x^3 - \frac{1}{x^3} - 3(x) \left(\frac{1}{x}\right) \left(x - \frac{1}{x}\right)$$

Putting value of $x - \frac{1}{x} = 7$

$$(7)^3 = x^3 - \frac{1}{x^3} - 3(7)$$

$$343 = x^3 - \frac{1}{x^3} - 21$$

MATHEMATICS FOR 9TH CLASS (UNIT # 4)

Pilot Superone Mathematics 204 Class 9th

$$343 + 21 = x^3 - \frac{1}{x^3}$$

$$364 = x^3 - \frac{1}{x^3}$$

or
$$x^3 - \frac{1}{x^3} = 364$$

Q.12 If $\left(3x + \frac{1}{3x}\right) = 5$, then find the value of $27x^3 + \frac{1}{27x^3}$

Solution:
$$3x + \frac{1}{3x} = 5 \quad \text{Given}$$

$$\left(3x + \frac{1}{3x}\right)^3 = (5)^3 \quad \text{(Cubing)}$$

$$(3x)^3 + \left(\frac{1}{3x}\right)^3 + 3(3x)\left(\frac{1}{3x}\right)\left(3x + \frac{1}{3x}\right) = 125$$

$$27x^3 + \frac{1}{27x^3} + 3 \times (5) = 125$$

$$27x^3 + \frac{1}{27x^3} = 125 - 15$$

$$27x^3 + \frac{1}{27x^3} = 110$$

Q.13 If $\left(5x - \frac{1}{5x}\right) = 6$, then find the value of $\left(125x^3 - \frac{1}{125x^3}\right)$

Solution:
$$5x - \frac{1}{5x} = 6 \quad \text{(Given)}$$

$$\left(5x - \frac{1}{5x}\right)^3 = (6)^3 \quad \text{(Cubing)}$$

MATHEMATICS FOR 9TH CLASS (UNIT # 4)

Pilot Superone Mathematics 205 Class 9th

$$(5x)^3 - \frac{1}{(5x)^3} - 3(5x) \left(\frac{1}{5x} \right) \left(5x - \frac{1}{5x} \right) = 216$$

$$125x^3 - \frac{1}{125x^3} - 3(6) = 216$$

$$125x^3 - \frac{1}{125x^3} = 216 + 18$$

$$125x^3 - \frac{1}{125x^3} = 234$$

Q.14 Factorize:

(i) $x^3 - y^3 - x + y$

(ii) $8x^3 - \frac{1}{27y^3}$

14(i) $x^3 - y^3 - x + y$

Solution: $= x^3 - y^3 - (x - y)$
 $= (x - y)(x^2 + xy + y^2) - (x - y)$
 $= (x - y)[x^2 + xy + y^2 - 1]$

14(ii) $8x^3 - \frac{1}{27y^3}$

Solution: $= (2x)^3 - \left(\frac{1}{3y} \right)^3$
 $= \left(2x - \frac{1}{3y} \right) \left[(2x)^2 + (2x) \left(\frac{1}{3y} \right) + \left(\frac{1}{3y} \right)^2 \right]$
 $= \left(2x - \frac{1}{3y} \right) \left[4x^2 + \frac{2x}{3y} + \frac{1}{9y^2} \right]$

MATHEMATICS FOR 9TH CLASS (UNIT # 4)

Pilot Superone Mathematics 206 Class 9th

(ii) $(x^3 - y^3)(x^6 + x^3y^3 + y^6)$

(iii) $(x - y)(x + y)(x^2 + y^2)(x^2 + xy + y^2)(x^2 - xy + y^2)$
 $(x^4 - x^2y^2 + y^4)$

(iv) $(2x^2 - 1)(2x^2 + 1)(4x^4 + 2x^2 + 1)(4x^4 - 2x^2 + 1)$

Solutions:

15(i) $(x^2 - y^2)(x^4 - x^2y^2 + y^4)$

Solution: $= (x^2 + y^2)(x^4 - x^2y^2 + y^4)$

Formula $[(x + y)(x^2 - xy + y^2) = x^3 + y^3]$
 $= (x^2)^3 + (y^2)^3$
 $= x^6 + y^6$

15(ii) $(x^3 - y^3)(x^6 + x^3y^3 + y^6)$

Solution: $[(x - y)(x^2 + xy + y^2) = x^3 - y^3]$ **Formula**
 $= (x^3)^3 - (y^3)^3$
 $= x^{3 \times 3} - y^{3 \times 3}$
 $= x^9 - y^9$

15(iii) $(x - y)(x + y)(x^2 + y^2)(x^2 + xy + y^2)(x^2 - xy + y^2)$
 $(x^4 - x^2y^2 + y^4)$

Solution: $= [(x - y)(x^2 + xy + y^2)] [(x + y)(x^2 - xy + y^2)]$
 $[(x^2 + y^2)(x^4 - x^2y^2 + y^4)]$
 $= [x^3 - y^3][x^3 + y^3] [(x^2)^3 + (y^2)^3]$
 $= [(x^3 - y^3)(x^3 + y^3)] [x^6 + y^6]$
 $= [(x^3)^2 - (y^3)^2] [x^6 + y^6]$
 $= (x^6 - y^6)(x^6 + y^6)$
 $= (x^6)^2 - (y^6)^2$
 $= x^{12} - y^{12}$

15(iv) $(2x^2 - 1)(2x^2 + 1)(4x^4 + 2x^2 + 1)(4x^4 - 2x^2 + 1)$

Solution: $= [(2x^2 - 1)(4x^4 + 2x^2 + 1)] [(2x^2 + 1)(4x^4 - 2x^2 + 1)]$
 $= [(2x^2)^3 - (1)^3] [(2x^2)^3 + (1)^3]$

MATHEMATICS FOR 9TH CLASS (UNIT # 4)

Prior Superone Mathematics 207 Class 9th

SURDS

An irrational radical with rational radicand is called a surd.

Hence $\sqrt{2}, \sqrt{3}, \sqrt{\frac{3}{7}}$ are surds.

$\sqrt[n]{a}$ is surd if

- (i) a is rational
- (ii) the result $\sqrt[n]{a}$ is irrational

e.g., $\sqrt{\pi}$ and $\sqrt{3+\sqrt{17}}$ are not surds as π and $3+\sqrt{17}$ are not rational

In $\sqrt[n]{a}$ n is degree of the surd

$a - \sqrt{b}$ is called conjugate of $a + \sqrt{b}$.

If we change the denominator from its surd form to rational form, then this process is rationalization

Exercise 4.3

Q.1 Express each of the following surd in the simplest form.

(i) $\sqrt{180}$

(ii) $3\sqrt{162}$

(iii) $\frac{3}{4}\sqrt[3]{128}$

(iv) $\sqrt[3]{96x^6y^7z^8}$

1(i) $\sqrt{180}$

Solution:

$$\begin{aligned}
 &= \sqrt{2 \times 2 \times 3 \times 3 \times 5} \\
 &= \sqrt{2^2 \times 3^2 \times 5} \\
 &= 2 \times 3 \sqrt{5} \\
 &= 6\sqrt{5}
 \end{aligned}$$

1(ii) $3\sqrt{162}$

Solution:

$$\begin{aligned}
 &= 3\sqrt{9 \times 9 \times 2} \\
 &= 3\sqrt{9^2 \times 2} \\
 &= 3 \times 9 \sqrt{2} \\
 &= 27\sqrt{2}
 \end{aligned}$$

1(iii) $\frac{3}{4}\sqrt[3]{128}$

Solution:

$$\begin{aligned}
 &= \frac{3}{4}\sqrt[3]{2^3 \times 2^3 \times 2} \\
 &= \frac{3}{4} \times 2 \times 2 \times \sqrt[3]{2} \\
 &= 3\sqrt[3]{2}
 \end{aligned}$$

1(iv) $\sqrt[3]{96x^6y^7z^8}$

MATHEMATICS FOR 9TH CLASS (UNIT # 4)

Pilot Superone Mathematics 209 Class 9th

$$2xyz \sqrt[3]{3xy^2z^3}$$

Q.2 Simplify

(i) $\frac{\sqrt{18}}{\sqrt{3}\sqrt{2}}$

(ii) $\frac{\sqrt{21}\sqrt{9}}{\sqrt{63}}$

(iii) $\sqrt[3]{243x^5y^{10}z^3}$

(iv) $\sqrt[4]{125}$

(v) $\sqrt{21} \times \sqrt{7} \times \sqrt{3}$

Solutions:

2(i) $\frac{\sqrt{18}}{\sqrt{3}\sqrt{2}}$

Solution:
$$\sqrt{\frac{18}{3 \times 2}}$$

$$\sqrt{\frac{3 \times 2 \times 3}{3 \times 2}} = \sqrt{3}$$

2(ii) $\frac{\sqrt{21}\sqrt{9}}{\sqrt{63}}$

Solution:
$$\sqrt{\frac{21 \times 9}{63}}$$

$$\sqrt{\frac{21 \times 3 \times 3}{21 \times 3}} = \sqrt{3}$$

2(iii) $\sqrt[3]{243x^5y^{10}z^3}$

Solution:
$$(3^3x^5y^{10}z^3)^{\frac{1}{3}}$$

$$= 3^{3 \times \frac{1}{3}} x^{5 \times \frac{1}{3}} y^{10 \times \frac{1}{3}} z^{3 \times \frac{1}{3}}$$

$$3xy^2z^3$$

2(iv) $\sqrt[4]{125}$

MATHEMATICS FOR 9TH CLASS (UNIT # 4)

Pilot Superone Mathematics 210 Class 9th

Solution:

$$\frac{4}{5} \sqrt{5^5}$$

$$\frac{4}{5} \times 5^2 \sqrt{5}$$

$$= \frac{4}{5} \times 5 = 4$$

2(v) $\sqrt{21} \times \sqrt{7} \times \sqrt{3}$

Solution:

$$\sqrt{21 \times 7 \times 3}$$

$$= \sqrt{21 \times 21}$$

$$[(21)^2]^{1/2} = (21)^{2 \times 1/2} = 21$$

Q.3 Simplify by combining similar terms.

(i) $\sqrt{45} - 3\sqrt{20} + 4\sqrt{5}$

(ii) $4\sqrt{12} + 5\sqrt{27} - 3\sqrt{75} + \sqrt{300}$

(iii) $\sqrt{3}(2\sqrt{3} + 3\sqrt{3})$

(iv) $2(6\sqrt{5} - 3\sqrt{5})$

3(i) $\sqrt{45} - 3\sqrt{20} + 4\sqrt{5}$

Solution:

$$\sqrt{9 \times 5} - 3\sqrt{4 \times 5} + 4\sqrt{5}$$

$$= \sqrt{3^2 \times 5} - 3\sqrt{2^2 \times 5} + 4\sqrt{5}$$

$$3\sqrt{5} - 3 \times 2\sqrt{5} + 4\sqrt{5}$$

$$\sqrt{5} | 3 - 6 + 4 |$$

$$= \sqrt{5} (1) = \sqrt{5}$$

3(ii) $4\sqrt{12} + 5\sqrt{27} - 3\sqrt{75} + \sqrt{300}$

Solution:

$$4\sqrt{4 \times 3} + 5\sqrt{9 \times 3} - 3\sqrt{25 \times 3} + \sqrt{100 \times 3}$$

$$4\sqrt{2^2 \times 3} + 5\sqrt{3^2 \times 3} - 3\sqrt{5^2 \times 3} + \sqrt{10^2 \times 3}$$

MATHEMATICS FOR 9TH CLASS (UNIT # 4)

Pilot Superone Mathematics 211 _____ Class 9th

$$8\sqrt{3} - 15\sqrt{3} - 15\sqrt{3} + 10\sqrt{3}$$

$$\sqrt{3}[8 + 15 - 15 - 10]$$

$$\sqrt{3}(18 - 18)\sqrt{3}$$

$$3(\text{iii}) \quad \sqrt{3}(2\sqrt{3} + 3\sqrt{3})$$

$$\text{Solution:} \quad (\sqrt{3})(\sqrt{3})(2 + 3)$$

$$3 \times 5 = 15 \quad (\sqrt{3})^2 (5)$$

$$3(\text{iv}) \quad 2(6\sqrt{5} - 3\sqrt{5})$$

$$\text{Solution:} \quad 2(\sqrt{5})(6 - 3)$$

$$2(\sqrt{5})(3) = 6\sqrt{5}$$

Q.4 Simplify:

$$(i) \quad (3 + \sqrt{3})(3 - \sqrt{3})$$

$$(ii) \quad (\sqrt{5} + \sqrt{3})^2$$

$$(iii) \quad (\sqrt{5} + \sqrt{3})(\sqrt{3} - \sqrt{3}) \quad (iv) \quad \left(\sqrt{2} + \frac{1}{\sqrt{3}}\right)\left(\sqrt{2} - \frac{1}{\sqrt{3}}\right)$$

$$(v) \quad (\sqrt{x} + \sqrt{y})(\sqrt{x} - \sqrt{y})(x + y)(x^2 + y^2)$$

$$4(i) \quad (3 + \sqrt{3})(3 - \sqrt{3})$$

$$\text{Solution:} \quad (3)^2 - (\sqrt{3})^2$$

$$9 - 3 = 6$$

$$4(ii) \quad (\sqrt{5} + \sqrt{3})^2$$

$$\text{Solution:} \quad (\sqrt{5})^2 + (\sqrt{3})^2 + 2\sqrt{5} \times \sqrt{3}$$

$$5 + 3 + 2\sqrt{5} \times 3$$

$$8 + 2\sqrt{15}$$

$$4(iii) \quad (\sqrt{5} + \sqrt{3})(\sqrt{5} - \sqrt{3})$$

$$\text{Solution:} \quad (\sqrt{5})^2 - (\sqrt{3})^2$$

MATHEMATICS FOR 9TH CLASS (UNIT # 4)

Pilot Superone Mathematics 212 Class 9th

4(iv) $\left(\sqrt{2} + \frac{1}{\sqrt{3}} \right) \left(\sqrt{2} - \frac{1}{\sqrt{3}} \right)$

Solution: $\left(\sqrt{2} \right)^2 - \left(\frac{1}{\sqrt{3}} \right)^2$

$$2 - \frac{1}{3}$$

$$\frac{6 - 1}{3} = \frac{5}{3}$$

4(v) $(\sqrt{x} + \sqrt{y})(\sqrt{x} - \sqrt{y})(x + y)(x^2 + y^2)$

Solution: $(\sqrt{x} + \sqrt{y})(\sqrt{x} - \sqrt{y})(x + y)(x^2 + y^2)$

$$[(\sqrt{x})^2 - (\sqrt{y})^2](x + y)(x^2 + y^2)$$

$$(x - y)(x + y)(x^2 + y^2)$$

$$[(x - y)(x + y)](x^2 + y^2)$$

$$(x^2 - y^2)(x^2 + y^2)$$

$$(x^2)^2 - (y^2)^2 = x^4 - y^4$$

Rationalization of Surds Definitions

(i) A surd which contains a single term is called a monomial surd.

(ii) A surd which contains sum of two monomial surds or sum of a monomial surd and a rational number is called a binomial surd.

(iii) The conjugate of $x + \sqrt{y}$ is $x - \sqrt{y}$

Pilot Superone Mathematics 213

Class 9th

Exercise 4.4

Q.1 Rationalize the denominator of the following:

(i) $\frac{3}{4\sqrt{3}}$

(ii) $\frac{14}{\sqrt{98}}$

(iii) $\frac{6}{\sqrt{8}\sqrt{27}}$

(iv) $\frac{1}{3+2\sqrt{5}}$

(v) $\frac{15}{\sqrt{3}-4}$

(vi) $\frac{2}{\sqrt{5}-\sqrt{3}}$

(vii) $\frac{\sqrt{3}-1}{\sqrt{3}+1}$

(viii) $\frac{\sqrt{3}+\sqrt{3}}{\sqrt{3}-\sqrt{3}}$

1(i) $\frac{3}{4\sqrt{3}}$

Solution:
$$\frac{3}{4\sqrt{3}} \times \frac{\sqrt{3}}{\sqrt{3}} = \frac{3\sqrt{3}}{4 \times 3} = \frac{1}{4}\sqrt{3}$$

1(ii) $\frac{14}{\sqrt{98}}$

Solution:
$$\frac{14}{\sqrt{98}} \times \frac{\sqrt{98}}{\sqrt{98}} = \frac{14\sqrt{98}}{98}$$

$$\frac{\sqrt{98}}{7} = \frac{\sqrt{7 \times 7 \times 2}}{7} = \frac{7\sqrt{2}}{7}$$

1(iii) $\frac{6}{\sqrt{8}\sqrt{27}}$

MATHEMATICS FOR 9TH CLASS (UNIT # 4)

Pilot Superone Mathematics 214 Class 9th

Solution:
$$\frac{6\sqrt{8}\sqrt{27}}{\sqrt{8}\sqrt{27}} \times \frac{\sqrt{8}\sqrt{27}}{\sqrt{8}\sqrt{27}}$$

$$\frac{6 \times \sqrt{8} \times \sqrt{27}}{8 \times 27}$$

$$\frac{6 \times \sqrt{2^3} \times 3 \times \sqrt{3^3} \times 3}{8 \times 27}$$

$$\frac{6 \times 2\sqrt{2} \times 3\sqrt{3}}{8 \times 27}$$

$$\frac{6 \times 2 \times 3 \times 2 \times \sqrt{3}}{8 \times 27}$$

$$\frac{1}{6} \times \sqrt{2 \times 3} \quad \frac{1}{6} \times \sqrt{6} = \frac{\sqrt{6}}{6}$$

1(iv) $3 + \frac{1}{2\sqrt{5}}$

Solution:
$$\frac{1}{3 + \frac{1}{2\sqrt{5}}} \times \frac{3 - \frac{1}{2\sqrt{5}}}{3 - \frac{1}{2\sqrt{5}}}$$

$$\frac{3 - \frac{1}{2\sqrt{5}}}{9 - \left(\frac{1}{2\sqrt{5}}\right)^2}$$

$$\frac{3 - \frac{1}{2\sqrt{5}}}{9 - \frac{1}{20}}$$

$$\frac{1}{11} \left(3 - \frac{1}{2\sqrt{5}}\right)$$

1(v) $\frac{15}{\sqrt{31} - 4}$

$$\frac{15}{\sqrt{31} - 4} \times \frac{(\sqrt{31} + 4)}{(\sqrt{31} + 4)}$$

MATHEMATICS FOR 9TH CLASS (UNIT # 4)

Pilot Superone Mathemat.cs 215 Class 9th

$$\frac{15(\sqrt{31} + 4)}{(\sqrt{31})^2 - (4)^2}$$

$$= \frac{15(\sqrt{31} + 4)}{31 - 16}$$

$$\frac{15(\sqrt{31} + 4)}{15} \quad \sqrt{31} + 4$$

1(vi) $\frac{2}{\sqrt{5} - \sqrt{3}}$

Solution:

$$\frac{2}{\sqrt{5} - \sqrt{3}} \times \left(\frac{\sqrt{5} + \sqrt{3}}{\sqrt{5} + \sqrt{3}} \right)$$

$$\frac{2(\sqrt{5} + \sqrt{3})}{(\sqrt{5})^2 - (\sqrt{3})^2}$$

$$\frac{2(\sqrt{5} + \sqrt{3})}{5 - 3}$$

$$= \frac{2(\sqrt{5} + \sqrt{3})}{2} = \sqrt{5} + \sqrt{3}$$

1(vii) $\frac{\sqrt{3} - 1}{\sqrt{3} + 1}$

Solution:

$$\frac{\sqrt{3} - 1}{\sqrt{3} + 1} \times \left(\frac{\sqrt{3} - 1}{\sqrt{3} - 1} \right)$$

$$= \frac{(\sqrt{3} - 1)^2}{(\sqrt{3})^2 - (1)^2}$$

MATHEMATICS FOR 9TH CLASS (UNIT # 4)

Pilot Superone Mathematics 216 Class 9

$$\frac{(\sqrt{3})^2 + (1) - 2\sqrt{3}(1)}{3 - 1}$$

$$\frac{3 + 1 - 2\sqrt{3}}{2}$$

$$\frac{4 - 2\sqrt{3}}{2}$$

$$\frac{2(2 - \sqrt{3})}{2} = 2 - \sqrt{3}$$

I (vii) $\frac{\sqrt{5}}{\sqrt{5} - \sqrt{3}}$

Solution: $\frac{(\sqrt{5} + \sqrt{3})}{(\sqrt{5} - \sqrt{3})} \times \frac{(\sqrt{5} + \sqrt{3})}{(\sqrt{5} + \sqrt{3})}$

$$\frac{(\sqrt{5} + \sqrt{3})^2}{(\sqrt{5})^2 - (\sqrt{3})^2}$$

$$\frac{(\sqrt{5})^2 + (\sqrt{3})^2 + 2\sqrt{5}\sqrt{3}}{5 - 3}$$

$$\frac{5 + 3 + 2\sqrt{15}}{2} = \frac{8 + 2\sqrt{15}}{2}$$

$$\frac{2(4 + \sqrt{15})}{2} = 4 + \sqrt{15}$$

Q.2 Find the conjugate of $x + \sqrt{y}$

(i) $3 + \sqrt{7}$

(ii) $4 - \sqrt{5}$

(iii) $2 + \sqrt{3}$

(iv) $2 + \sqrt{5}$

MATHEMATICS FOR 9TH CLASS (UNIT # 4)

Pilot Superone Mathematics 217 Class 9th

(vii) $7 - \sqrt{6}$ (viii) $9 + \sqrt{2}$

2(i) $3 + \sqrt{7}$

Solution: Conjugate $3 - \sqrt{7}$

2(ii) $4 - \sqrt{5}$

Solution: Conjugate $4 + \sqrt{5}$

2(iii) $2 + \sqrt{3}$

Solution: Conjugate $2 - \sqrt{3}$

2(iv) $2 + \sqrt{5}$

Solution: Conjugate $2 - \sqrt{5}$

2(v) $5 + \sqrt{7}$

Solution: Conjugate $5 - \sqrt{7}$

2(vi) $4 - \sqrt{15}$

Solution: Conjugate $4 + \sqrt{15}$

2(vii) $7 - \sqrt{6}$

Solution: Conjugate $7 + \sqrt{6}$

2(viii) $9 + \sqrt{2}$

Solution: Conjugate $9 - \sqrt{2}$

Q.3 (i) If $x = 2 - \sqrt{3}$, find $\frac{1}{x}$

(ii) If $x = 4 - \sqrt{17}$, find $\frac{1}{x}$

(iii) If $x = \sqrt{3} + 2$, find $x + \frac{1}{x}$

3(i) $x = 2 - \sqrt{3}$

MATHEMATICS FOR 9TH CLASS (UNIT # 4)

Pilot Superone Mathematics 218 Class 9th

Solution: $\frac{1}{x} = \frac{1}{2 - \sqrt{3}}$

$$\frac{1}{2 - \sqrt{3}} \cdot \frac{(2 + \sqrt{3})}{(2 + \sqrt{3})}$$

$$= \frac{2 + \sqrt{3}}{(2)^2 - (\sqrt{3})^2}$$

$$= \frac{2 + \sqrt{3}}{4 - 3} = 2 + \sqrt{3}$$

3(ii) $x = 4 - \sqrt{17}$

Solution: $\frac{1}{x} = \frac{1}{4 - \sqrt{17}}$

$$\frac{1}{4 - \sqrt{17}} \cdot \frac{(4 + \sqrt{17})}{(4 + \sqrt{17})}$$

$$= \frac{4 + \sqrt{17}}{(4)^2 - (\sqrt{17})^2}$$

$$\frac{4 + \sqrt{17}}{16 - 17}$$

$$\frac{4 + \sqrt{17}}{-1} = -4 - \sqrt{17}$$

3(iii) $x = \sqrt{3} + 2$

Solution: $\frac{1}{x} = \frac{1}{\sqrt{3} + 2}$

$$\frac{1}{\sqrt{3} + 2} \cdot \frac{(\sqrt{3} - 2)}{(\sqrt{3} - 2)}$$

MATHEMATICS FOR 9TH CLASS (UNIT # 4)

Pilot Superone Mathematics 210 Class 9th

$$\frac{\sqrt{3}-2}{(\sqrt{3})^2 - (2)^2}$$

$$= \frac{\sqrt{3}-2}{3-4}$$

$$= \frac{\sqrt{3}-2}{-1} = \sqrt{3}+2$$

Now $x = \sqrt{3}+2$

And $\frac{1}{x} = \frac{1}{\sqrt{3}+2}$

$$x + \frac{1}{x} = (\sqrt{3}+2) + \frac{1}{\sqrt{3}+2}$$

$$= \sqrt{3}+2 + \frac{\sqrt{3}-2}{(\sqrt{3}+2)(\sqrt{3}-2)}$$

Q.4 Simplify:

(i) $\frac{1+\sqrt{2}}{\sqrt{5}+\sqrt{3}} + \frac{1-\sqrt{2}}{\sqrt{5}-\sqrt{3}}$

(ii) $\frac{1}{2+\sqrt{3}} + \frac{2}{\sqrt{5}-\sqrt{3}} + \frac{1}{2+\sqrt{3}}$

(iii) $\frac{2}{\sqrt{5}+\sqrt{3}} + \frac{1}{\sqrt{3}+\sqrt{2}} + \frac{3}{\sqrt{5}+\sqrt{2}}$

9 (i) $\frac{1+\sqrt{2}}{\sqrt{5}+\sqrt{3}} + \frac{1-\sqrt{2}}{\sqrt{5}-\sqrt{3}}$

Solution:

$$\frac{(1+\sqrt{2})(\sqrt{5}-\sqrt{3})}{(\sqrt{5}+\sqrt{3})(\sqrt{5}-\sqrt{3})} + \frac{(1-\sqrt{2})(\sqrt{5}+\sqrt{3})}{(\sqrt{5}-\sqrt{3})(\sqrt{5}+\sqrt{3})}$$

MATHEMATICS FOR 9TH CLASS (UNIT # 4)

Pilot Superone Mathematics 220 Class 9th

$$\frac{(1+\sqrt{2})(\sqrt{5}-\sqrt{3})}{(\sqrt{5})-(\sqrt{3})} + \frac{(1-\sqrt{2})(\sqrt{5}+\sqrt{3})}{(\sqrt{5})+(\sqrt{3})}$$

$$\frac{(1+\sqrt{2})(\sqrt{5}-\sqrt{3})}{5-3} + \frac{(1-\sqrt{2})(\sqrt{5}+\sqrt{3})}{5+3} \quad (1)$$

$$\frac{\sqrt{5}-\sqrt{3}+\sqrt{2}\times 5-\sqrt{2}\times 3}{2} + \frac{\sqrt{5}+\sqrt{3}-\sqrt{2}\times 5+\sqrt{2}\times 3}{2}$$

$$\frac{\sqrt{5}-\sqrt{3}+\sqrt{10}-\sqrt{6}+\sqrt{5}+\sqrt{3}-\sqrt{10}+\sqrt{6}}{2}$$

$$\frac{2\sqrt{5}-2\sqrt{6}}{2}$$

$$2(\sqrt{5}-\sqrt{6}) \quad \sqrt{5}-\sqrt{6}$$

$$4(ii) \quad \frac{2}{\sqrt{5}+\sqrt{3}} + \frac{1}{\sqrt{3}+\sqrt{2}} - \frac{3}{\sqrt{5}+\sqrt{2}}$$

Solution:

$$\frac{1(2-\sqrt{3})}{(2+\sqrt{3})(2-\sqrt{3})} + \frac{2(\sqrt{5}+\sqrt{3})}{(\sqrt{5}-\sqrt{3})(\sqrt{5}+\sqrt{3})} + \frac{1(2-\sqrt{5})}{(2+\sqrt{5})(2-\sqrt{5})}$$

$$\frac{2-\sqrt{3}}{(2+\sqrt{3})(2-\sqrt{3})} + \frac{2(\sqrt{5}+\sqrt{3})}{(\sqrt{5})-(\sqrt{3})} + \frac{2-\sqrt{5}}{(2+\sqrt{5})(2-\sqrt{5})}$$

$$\frac{2-\sqrt{3}}{4-3} + \frac{2(\sqrt{5}+\sqrt{3})}{(\sqrt{5})-(\sqrt{3})} + \frac{2-\sqrt{5}}{(2)^2-(\sqrt{5})^2}$$

$$2-\sqrt{3} + \frac{2(\sqrt{5}+\sqrt{3})}{\sqrt{5}-\sqrt{3}} + \frac{2-\sqrt{5}}{2-5}$$

MATHEMATICS FOR 9TH CLASS (UNIT # 4)

Pilot Superiore Mathematics 221 Class 9th

$$\frac{2 - \sqrt{3}}{1} + \frac{2(\sqrt{5} + \sqrt{3})}{2} + \frac{2 - \sqrt{5}}{-1}$$

$$\frac{2 - \sqrt{3} + \sqrt{5} + \sqrt{3} - 2 + \sqrt{5}}{2\sqrt{5}}$$

4(iii) $\frac{2}{\sqrt{5} + \sqrt{3}} + \frac{1}{\sqrt{3} + \sqrt{2}} - \frac{3}{\sqrt{5} + \sqrt{2}}$

Solution: A $\frac{2}{\sqrt{5} + \sqrt{3}}$, B $\frac{1}{\sqrt{3} + \sqrt{2}}$, C $\frac{3}{\sqrt{5} + \sqrt{2}}$

A $\frac{2(\sqrt{5} - \sqrt{3})}{(\sqrt{5} + \sqrt{3})(\sqrt{5} - \sqrt{3})}$

$$\frac{2(\sqrt{5} - \sqrt{3})}{(\sqrt{5})^2 - (\sqrt{3})^2}$$

$$\frac{2(\sqrt{5} - \sqrt{3})}{5 - 3}$$

$$\frac{2(\sqrt{5} - \sqrt{3})}{2} = \sqrt{5} - \sqrt{3}$$

B $\frac{1}{\sqrt{3} + \sqrt{2}}$

$$\frac{1(\sqrt{3} - \sqrt{2})}{(\sqrt{3} + \sqrt{2})(\sqrt{3} - \sqrt{2})}$$

$$\frac{1(\sqrt{3} - \sqrt{2})}{(\sqrt{3})^2 - (\sqrt{2})^2}$$

MATHEMATICS FOR 9TH CLASS (UNIT # 4)

Pilot Superone Mathematics 222 Class 9th

$$= \frac{\sqrt{3} - \sqrt{2}}{3 - 2} = \sqrt{3} - \sqrt{2}$$

$$C \quad \frac{3(\sqrt{5} - \sqrt{2})}{(\sqrt{5} + \sqrt{2})(\sqrt{5} - \sqrt{2})}$$

$$\frac{3(\sqrt{5} - \sqrt{2})}{(\sqrt{5})^2 - (\sqrt{2})^2}$$

$$= \frac{3(\sqrt{5} - \sqrt{2})}{5 - 2}$$

$$\frac{3(\sqrt{5} - \sqrt{2})}{3} = \sqrt{5} - \sqrt{2}$$

$$A + B - C$$

$$(\sqrt{5} - \sqrt{3}) + (\sqrt{3} - \sqrt{2}) - (\sqrt{5} - \sqrt{2})$$

$$= \sqrt{5} - \sqrt{3} + \sqrt{3} - \sqrt{2} - \sqrt{5} + \sqrt{2}$$

$$0$$

Q.5, (i) If $x = 2 + \sqrt{3}$, then find the value of $x - \frac{1}{x}$ and

$$\left(x - \frac{1}{x}\right)^2$$

(ii) If $x = \frac{\sqrt{5} - \sqrt{2}}{\sqrt{5} + \sqrt{2}}$, then find the value of $x + \frac{1}{x}$,

$$x + \frac{1}{x}, \text{ and } x^2 + \frac{1}{x^2}$$

MATHEMATICS FOR 9TH CLASS (UNIT # 4)

Pilot Superone Mathematics 223 Class 9th

5(i) If $x = 2 + \sqrt{3}$, then find the value of $x + \frac{1}{x}$ and $\left(x - \frac{1}{x}\right)$

Solution:

$$x = 2 + \sqrt{3}$$

$$\frac{1}{x} = \frac{1}{2 + \sqrt{3}}$$

$$\frac{1(2 - \sqrt{3})}{(2 + \sqrt{3})(2 - \sqrt{3})}$$

$$\frac{2 - \sqrt{3}}{(2)^2 - (\sqrt{3})^2}$$

$$= \frac{2 - \sqrt{3}}{4 - 3} = 2 - \sqrt{3}$$

Now $x + \frac{1}{x} = (2 + \sqrt{3}) + (2 - \sqrt{3})$

$$= 2 + \sqrt{3} + 2 - \sqrt{3}$$

Thus $x + \frac{1}{x} = 2\sqrt{3}$

$$\left(x - \frac{1}{x}\right)^2 = (2\sqrt{3})^2 \quad \text{(Squaring)}$$

$$4 \times 3 = 12$$

5(ii) If $x = \frac{\sqrt{5} - \sqrt{2}}{\sqrt{5} + \sqrt{2}}$, then find the value of $x + \frac{1}{x}$, $x - \frac{1}{x}$ and $x^2 + \frac{1}{x^2}$

Solution:

$$x = \frac{\sqrt{5} - \sqrt{2}}{\sqrt{5} + \sqrt{2}}$$

MATHEMATICS FOR 9TH CLASS (UNIT # 4)

Pilot Superone Mathematics 224 Class 9th

$$\begin{aligned} & \frac{1}{x} \cdot \frac{\sqrt{5} + \sqrt{2}}{\sqrt{5} - \sqrt{2}} \\ \text{Now } & \frac{1}{x} \cdot \frac{\sqrt{5} + \sqrt{2}}{\sqrt{5} + \sqrt{2}} \cdot \frac{\sqrt{5} + \sqrt{2}}{\sqrt{5} - \sqrt{2}} \\ & \frac{(\sqrt{5} - \sqrt{2}) + (\sqrt{5} + \sqrt{2})}{(\sqrt{5} + \sqrt{2})(\sqrt{5} - \sqrt{2})} \\ & \frac{5 + 2 - 2\sqrt{5 \times 2} + 5 + 2 + 2\sqrt{5 \times 2}}{(\sqrt{5})^2 - (\sqrt{2})^2} \\ & \frac{7 - 2\sqrt{10} + 7 + 2\sqrt{10}}{5 - 2} \\ & \frac{14}{3 - 2} = \frac{14}{3} \end{aligned}$$

$$x^2 + \frac{1}{x^2} = ?$$

$$\left(x + \frac{1}{x}\right)^2 = \left(\frac{14}{3}\right)^2$$

$$x^2 + \frac{1}{x^2} + 2 = \frac{196}{9}$$

$$x^2 + \frac{1}{x^2} = \frac{196}{9} - 2$$

$$\frac{196}{9} - \frac{18}{9} = \frac{178}{9}$$

$$x^2 + \frac{1}{x^2} = ?$$

$$\text{Now } \left(x + \frac{1}{x}\right)^2 = \left(\frac{14}{3}\right)^2$$

MATHEMATICS FOR 9TH CLASS (UNIT # 4)

Pilot Superone Mathematics 225 Class 9th

$$x^3 + \frac{1}{x^3} + 3x \times \frac{1}{x} \left(x + \frac{1}{x} \right) = \frac{27+4}{27}$$

$$x^3 + \frac{1}{x^3} + 3 \left(\frac{14}{3} \right) = \frac{27+4}{27}$$

$$x^3 + \frac{1}{x^3} + 14 = \frac{27+4}{27}$$

$$x^3 + \frac{1}{x^3} = \frac{27+4}{27} - 14$$

$$= \frac{27+4}{27} - \frac{378}{27} = \frac{236}{27}$$

Q.6 Determine the rational number a and b if

$$\frac{\sqrt{3}-1}{\sqrt{3}+1} + \frac{\sqrt{3}+1}{\sqrt{3}-1} = a + b\sqrt{3}$$

$$\frac{\sqrt{3}-1}{\sqrt{3}+1} + \frac{\sqrt{3}+1}{\sqrt{3}-1} = a + b\sqrt{3}$$

$$\frac{(\sqrt{3}-1)(\sqrt{3}-1) + (\sqrt{3}+1)(\sqrt{3}+1)}{(\sqrt{3}+1)(\sqrt{3}-1)} = a + b\sqrt{3}$$

$$\frac{(\sqrt{3}-1)^2 + (\sqrt{3}+1)^2}{(\sqrt{3})^2 - (1)^2} = a + b\sqrt{3}$$

$$\frac{(3+1-2\sqrt{3}) + (3+1+2\sqrt{3})}{3-1} = a + b\sqrt{3}$$

$$\frac{8}{2} = a + b\sqrt{3}$$

$$\Rightarrow 4 + (0)\sqrt{3} = a + b\sqrt{3}$$

**Unit
5**

FACTORIZATION

The process of writing a polynomial as a product of two or more than two polynomials is called factorization. Every new polynomial is called a factor of the given polynomial.

For example In $ab + bc = b(a + c)$, $b(a + c)$ is the factorization of $ab + bc$ and b , $(a + c)$ are its factors.

Exercise 5.1

Factorize:

- Q.1. (i) $2abc - 4abx + 2abd$
 (ii) $9xy - 12x^2y + 18y^2$
 (iii) $3x^2y - 3x + 9xy^2$
 (iv) $5ab^2c^3 - 10a^2b^3c - 20a^3bc^2$
 (v) $3x^2y(x - 3y) - 7x^2y^2(x - 3y)$
 (vi) $2xy^2(x^2 + 5) + 8xy^2(x^2 + 5)$

Solutions:-

- 1(i) $2abc - 4abx + 2abd$
 $2ab(c - 2x + d)$
 1(ii) $9xy - 12x^2y + 18y^2$
 $= 3y(3x - 4x^2 + 6y)$
 1(iii) $3x^2y - 3x + 9xy^2$
 $= 3x(xy + 1 - 3y^2)$
 1(iv) $5ab^2c^3 - 10a^2b^3c - 20a^3bc^2$
 $5abc(bc^2 - 2ab^2 - 4a^2c)$
 1(v) $3x^2y(x - 3y) - 7x^2y^2(x - 3y)$
 $x^2y(x - 3y)(3x - 7y)$
 1(vi) $2xy^2(x^2 + 5) + 8xy^2(x^2 + 5)$
 $2xy^2(x^2 + 5)(1 + 4)$

MATHEMATICS FOR 9TH CLASS (UNIT # 5)

Pilot Superone Mathematics 238 Class 9th

Q.2. Factorize:

(i) $5ax - 3ay - 5bx + 3by$

(ii) $3xy + 2y - 12x - 8$

(iii) $x^3 + 3xy^2 - 2x^2y - 6y^3$

(iv) $(x^2 - y^2)z + (y^2 - z^2)x$

Solution:-

2(i) $5ax - 3ay - 5bx + 3by$
 $(5ax - 3ay) - (5bx - 3by)$
 $a(5x - 3y) - b(5x - 3y)$
 $(5x - 3y)(a - b)$

2(ii) $3xy + 2y - 12x - 8$
 $(3xy + 2y) - (12x + 8)$
 $y(3x + 2) - 4(3x + 2)$
 $(3x + 2)(y - 4)$

2(iii) $x^3 + 3xy^2 - 2x^2y - 6y^3$
 $(x^3 + 3xy^2) - (2x^2y + 6y^3)$
 $x(x^2 + 3y^2) - 2y(x^2 + 3y^2)$
 $(x^2 + 3y^2)(x - 2y)$

2(iv) $(x^2 - y^2)z + (y^2 - z^2)x$
 $x^2z - y^2z + y^2x - z^2x$
 $x^2z + y^2x - y^2z - z^2x$
 $x(xz + y^2) - z(y^2 + xz)$
 $(xz - y^2)(x - z)$

Q.3. Factorize:

(i) $144a^2 + 24a + 1$

(ii) $\frac{a^2}{b^2} - 2 + \frac{b^2}{a^2}$

(iii) $(x + y)^2 - 14z(x + y) + 49z^2$

(iv) $12x^2 - 36x + 27$

Solutions:-

3(i) $144a^2 + 24a + 1$
 $= 144a^2 - 12a + 12a + 1$

MATHEMATICS FOR 9TH CLASS (UNIT # 5)

Pilot Superone Mathematics 239 Class 9th

$$\begin{aligned} & (144a^2 + 12a + 1)(12a + 1) \\ & 12a(12a + 1) + 1(12a + 1) \\ & (12a + 1)(12a + 1) = (12a + 1)^2 \\ 3(ii) \quad & \frac{a^2}{b^2} - 2 + \frac{b^2}{a^2} \end{aligned}$$

$$\begin{aligned} & \left(\frac{a}{b}\right)^2 - 2\left(\frac{a}{b}\right)\left(\frac{b}{a}\right) + \left(\frac{b}{a}\right)^2 \\ & \left(\frac{a}{b} - \frac{b}{a}\right)^2 \\ & \frac{a^2}{b^2} - 2 + \frac{b^2}{a^2} \\ & \frac{a^2}{b^2} - 2 \times \frac{a}{b} \times \frac{b}{a} + \frac{b^2}{a^2} \\ & \frac{a^2}{b^2} - \frac{a}{b} \times \frac{b}{a} - \frac{a}{b} \times \frac{b}{a} + \frac{b^2}{a^2} \\ & \left(\frac{a^2}{b^2} - \frac{a}{b} \times \frac{b}{a}\right) - \left(\frac{a}{b} \times \frac{b}{a} - \frac{b^2}{a^2}\right) \\ & \frac{a}{b}\left(\frac{a}{b} - \frac{b}{a}\right) - \frac{b}{a}\left(\frac{a}{b} - \frac{b}{a}\right) \\ & \left(\frac{a}{b} - \frac{b}{a}\right)\left(\frac{a}{b} - \frac{b}{a}\right) \\ & \left(\frac{a}{b} - \frac{b}{a}\right)^2 \end{aligned}$$

$$\begin{aligned} 3(iii) \quad & (x + y)^2 - 14z(x + y) + 49z^2 \\ & (x + y)^2 - 7z(x + y) - 7z(x + y) + 49z^2 \\ & [(x + y)^2 - 7z(x + y)] - [7z(x + y) - 49z^2] \\ & (x + y)(x + y - 7z) - (7z)(x + y - 7z) \\ & (x + y - 7z)(x + y - 7z) \\ & (x + y - 7z)^2 \end{aligned}$$

$$\begin{aligned} 3(iv) \quad & 12x^2 - 36x - 27 \\ & 3(4x^2 - 12x - 9) \\ & 3(2x(2x - 3) - 3(2x - 3)) \\ & = 3(2x - 3)(2x - 3) \end{aligned}$$

MATHEMATICS FOR 9TH CLASS (UNIT # 5)

Pilot Superone Mathematicist 240 Class 9th

$$3(2x - 3)^2$$

Q.4. Factorize:

(i) $3x^2 - 75y^2$

(ii) $x(x - 1) - y(y - 1)$

(iii) $128a^{100} - 242a^{99}$

(iv) $3x - 243x^3$

Solution:-

4(i) $3x^2 - 75y^2$

$$3(x^2 - 25y^2)$$

$$3[(x)^2 - (5y)^2]$$

$$3(x + 5y)(x - 5y)$$

4(ii) $x(x - 1) - y(y - 1)$

$$x^2 - x - y^2 + y$$

$$x^2 - y^2 - x + y$$

$$(x^2 - y^2) - (x - y)$$

$$(x + y)(x - y) - 1(x - y)$$

$$(x + y)(x - y - 1)$$

$$(x + y)(x - y - 1)$$

4(iii) $128a^{100} - 242a^{99}$

$$2a(64a^{99} - 121a^{98})$$

$$2a[(8a)^2 - (11a)^2]$$

$$2a(8a + 11a)(8a - 11a)$$

4(iv) $3x - 243x^3$

$$3x(1 - 81x^2)$$

$$= 3x[(1)^2 - (9x)^2]$$

$$= 3x(1 + 9x)(1 - 9x)$$

Q.5. Factorize:

(i) $x^2 - y^2 - 6y - 9$

(ii) $x^2 - a^2 + 2a - 1$

(iii) $4x^2 - y^2 - 2y - 1$

(iv) $x^2 - y^2 - 4x - 2y +$

(v) $25x^2 - 10x + 1 - 36z^2$

(vi) $x^2 - y^2 - 4xz + 4z^2$

MATHEMATICS FOR 9TH CLASS (UNIT # 5)

Prior Supreme Mathematics 241 Class 9th

Solutions:-

$$\begin{aligned} 3(i) \quad & x^2 - y^2 - 6y + 9 \\ & = x^2 - (y^2 + 6y + 9) \\ & = (x)^2 - \{y^2 + 3y + 3y + 9\} \\ & = (x)^2 - \{y(y + 3) + 3(y + 3)\} \\ & = (x)^2 - (y + 3)(y + 3) \\ & = (x)^2 - (y + 3)^2 \\ & = [x + (y + 3)][x - (y + 3)] \end{aligned}$$

$$\begin{aligned} 3(ii) \quad & x^2 - a^2 + 2a - 1 \\ & = x^2 - (a^2 - 2a + 1) \\ & = (x)^2 - [a^2 - a - a + 1] \\ & = (x)^2 - [a(a - 1) - 1(a - 1)] \\ & = (x)^2 - (a - 1)(a - 1) \\ & = (x)^2 - (a - 1)^2 \\ & = (x + a - 1)(x - a + 1) \end{aligned}$$

$$\begin{aligned} 3(iii) \quad & 4x^2 - y^2 - 2y + 1 \\ & = (2x)^2 - (y^2 + 2y + 1) \\ & = (2x)^2 - [y^2 + y + y + 1] \\ & = (2x)^2 - [y(y + 1) + 1(y + 1)] \\ & = (2x)^2 - (y + 1)(y + 1) \\ & = (2x)^2 - (y + 1)^2 \\ & = (2x + y + 1)(2x - y - 1) \end{aligned}$$

$$\begin{aligned} 3(iv) \quad & x^2 - y^2 - 4x - 2y + 3 \\ & \text{Re-arranging} \\ & x^2 - 4x + 4 - y^2 - 2y + 1 \quad NTS \\ & = (x^2 - 4x + 4) - (y^2 + 2y + 1) \\ & = [x^2 - 2x - 2x + 4] - [y^2 + y - y + 1] \\ & = [x(x - 2) - 2(x - 2)] - [y(y + 1) - 1(y + 1)] \\ & = [(x - 2)(x - 2)] - [(y + 1)(y + 1)] \end{aligned}$$

www.dn

MATHEMATICS FOR 9TH CLASS (UNIT # 5)

Pilot Superone Mathematics 242 **Class 9th**

$$\begin{aligned} &= (x-2)^2 - (y-1)^2 \\ &= (x-2+y+1)(x-2-y-1) \\ &= (x+y-1)(x-y-3) \end{aligned}$$

$$\begin{aligned} 5(v) \quad &25x^2 - 10x + 1 - 36z^2 \\ &= (25x^2 - 10x + 1) - (6z)^2 \\ &= [25x^2 - 5x - 5x + 1] - (6z)^2 \\ &= [5x(5x-1) - 1(5x-1)] - (6z)^2 \\ &= (5x-1)(5x-1) - (6z)^2 \\ &= (5x-1)^2 - (6z)^2 \\ &= (5x-1+6z)(5x-1-6z) \end{aligned}$$

$$\begin{aligned} 5(vi) \quad &x^2 - y^2 - 4xz + 4z^2 \\ &= (x^2 - 4xz + 4z^2) - (y^2) \\ &= [x^2 - 2xz - 2xz + 4z^2] - (y^2) \\ &= [x(x-2z) - 2z(x-2z)] - y^2 \\ &= (x-2z)(x-2z) - y^2 \\ &= (x-2z)^2 - (y)^2 \\ &= (x-2z+y)(x-2z-y) \\ &= (x+y-2z)(x-y-2z) \end{aligned}$$

www.don

MATHEMATICS FOR 9TH CLASS (UNIT # 5)

Pilot Superone Mathematics 243 Class 9th

Exercise 5.2

Q.1. Factorize:

(i) $x^4 + \frac{1}{x^2} - 3$

(ii) $3x^4 + 12y^4$

(iii) $a^4 + 3a^2b^2 + 4b^4$

(iv) $4x^4 + 81$

(v) $x^4 + x^2 + 25$

(vi) $x^4 + 4x^2 + 16$

Solutions:-

1(i) $x^4 + \frac{1}{x^2} - 3$

$$x^4 + \frac{1}{x^2} - 2 + 1$$

$$= \left[(x^2)^2 - 2 + \left(\frac{1}{x^2}\right)^2 \right] + 1$$

$$= \left(x^2 - \frac{1}{x^2} \right)^2 - (1)^2$$

$$= \left(x^2 - \frac{1}{x^2} + 1 \right) \left(x^2 - \frac{1}{x^2} - 1 \right)$$

1(ii) $3x^4 + 12y^4$
 $= 3(x^4 + 4y^4)$

Completing the square

$$= 3\{(x^2)^2 + (2y^2)^2 + 4x^2y^2 - 4x^2y^2\}$$

$$= 3\{(x^2)^2 + 2(x^2)(2y^2) + (2y^2)^2 - (2xy)^2\}$$

$$= 3\{(x^2 + 2y^2)^2 - (2xy)^2\}$$

$$= 3(x^2 + 2y^2 + 2xy)(x^2 + 2y^2 - 2xy)$$

$$= 3(x^2 + 2xy + 2y^2)(x^2 - 2xy + 2y^2)$$

1(iii) $a^4 + 3a^2b^2 + 4b^4$

Completing the square

$$= (a^2)^2 + 1(a^2)(2b^2) + (2b^2)^2 - a^2b^2$$

$$= (a^2 + 2b^2)^2 - (ab)^2$$

$$= (a^2 + 2b^2 + ab)(a^2 + 2b^2 - ab)$$

MATHEMATICS FOR 9TH CLASS (UNIT # 5)

Pilot Superior Mathematics 244 Class 9th

$$\begin{aligned}
 1(iv) \quad & 4x^2 + 81 \\
 & \text{Completing the square} \\
 & = (2x^2)^2 + (9)^2 \\
 & = (2x^2)^2 + (9)^2 + 2(2x^2)(9) - 36x^2 \\
 & = (2x^2 + 9 - 6x)(2x^2 + 9 + 6x) \\
 & = (2x^2 - 6x + 9)(2x^2 + 6x + 9)
 \end{aligned}$$

$$\begin{aligned}
 1(v) \quad & x^2 + x^2 + 25 \\
 & = (x^2)^2 + (5)^2 + x^2 \\
 & \text{Completing the square} \\
 & = (x^2)^2 + (5)^2 + 2(x^2)(5) - 9x^2 \\
 & = (x^2 + 5)^2 - (3x)^2 \\
 & = (x^2 + 5 + 3x)(x^2 + 5 - 3x) \\
 & = (x^2 + 3x + 5)(x^2 - 3x + 5)
 \end{aligned}$$

$$\begin{aligned}
 1(vi) \quad & x^2 + 4x^2 + 16 \\
 & = (x^2)^2 + (4)^2 + 2(x^2)(4) - 4x^2 \\
 & \text{Completing the square} \\
 & = (x^2)^2 + (4)^2 + 2(x^2)(4) - 4x^2 \\
 & = (x^2 + 4)^2 - (2x)^2 \\
 & = (x^2 + 4 + 2x)(x^2 + 4 - 2x) \\
 & = (x^2 + 2x + 4)(x^2 - 2x + 4)
 \end{aligned}$$

Q.2. Factorize:

$$(i) \quad x^2 + 14x + 48$$

$$(iii) \quad x^2 - 11x - 42$$

$$(ii) \quad x^2 - 21x + 108$$

$$(iv) \quad x^2 + x - 132$$

Solutions:-

$$\begin{aligned}
 2(i) \quad & x^2 + 14x + 48 \\
 & = x^2 + 8x + 6x + 48 \\
 & = x(x + 8) + 6(x + 8) \\
 & = (x + 8)(x + 6)
 \end{aligned}$$

$$\begin{aligned}
 2(ii) \quad & x^2 - 21x + 108 \\
 & = x^2 - 12x - 9x + 108 \\
 & = x(x - 12) - 9(x - 12) \\
 & = (x - 12)(x - 9)
 \end{aligned}$$

$$2(iii) \quad x^2 - 11x - 42$$

MATHEMATICS FOR 9TH CLASS (UNIT # 5)

Pilot Superior Mathematics 245 **Class 9th**

$$= x^2 - 11x + 42$$

$$= x^2 - 14x + 3x + 42$$

$$= x(x - 14) + 3(x - 14)$$

$$= (x - 14)(x + 3)$$

2(iv) $x^2 + x - 132$

$$= x^2 + 12x - 11x - 132$$

$$= x(x + 12) - 11(x + 12)$$

$$= (x + 12)(x - 11)$$

Q.3. Factorize:

(i) $4x^2 + 12x + 5$	(ii) $30x^2 + 7x - 15$
(iii) $24x^2 - 65x + 21$	(iv) $5x^2 - 16x - 21$
(v) $4x^2 - 17xy + 13y^2$	(vi) $3x^2 - 38xy + 13y^2$
(vii) $5x^2 + 33xy - 14y^2$	
(viii) $\left(5x - \frac{1}{x}\right)^2 + 4\left(5x - \frac{1}{x}\right) + 4, x \neq 0$	

Solutions:-

3(i) $4x^2 + 12x + 5$

$$= 4x^2 + 10x + 2x + 5$$

$$= 2x(2x + 5) + 1(2x + 5)$$

$$= (2x + 5)(2x + 1)$$

3(ii) $30x^2 + 7x - 15$

$$= 30x^2 + 25x - 18x - 15$$

$$= 5x(6x + 5) - 3(6x + 5)$$

$$= (6x + 5)(5x - 3)$$

3(iii) $24x^2 - 65x + 21$

$$= 24x^2 - 56x + 9x + 21$$

$$= 8x(3x - 7) + 3(3x - 7)$$

$$= (3x - 7)(8x + 3)$$

3(iv) $5x^2 - 16x - 21$

$$= 5x^2 - 21x + 5x - 21$$

$$= x(5x - 21) + 1(5x - 21)$$

$$= (5x - 21)(x + 1)$$

MATHEMATICS FOR 9TH CLASS (UNIT # 5)

Pilot Superong Mathematics 246 Class 9th

$$\begin{aligned} 3(v) \quad & 4x^2 - 17xy + 4y^2 \\ & = 4x^2 - 16xy - xy + 4y^2 \\ & = 4x(x - 4y) - y(x - 4y) \\ & = (x - 4y)(4x - y) \end{aligned}$$

$$\begin{aligned} 3(vi) \quad & 3x^2 - 38xy + 13y^2 \\ & = 3x^2 - xy - 39xy + 13y^2 \\ & = x(3x + y) - 13y(3x + y) \\ & = (3x + y)(x - 13y) \end{aligned}$$

$$\begin{aligned} 3(vii) \quad & 5x^2 + 33xy + 4y^2 \\ & 5x^2 + 35xy - 2xy + 4y^2 \\ & 5x(x + 7y) - 2y(x + 7y) \\ & = (x + 7y)(5x - 2y) \end{aligned}$$

$$\begin{aligned} 3(viii) \quad & \left(5x - \frac{1}{x}\right)^2 + 4\left(5x - \frac{1}{x}\right) + 4, x \neq 0 \\ & \left(5x - \frac{1}{x}\right)^2 + 2\left(5x - \frac{1}{x}\right) + 2\left(5x - \frac{1}{x}\right) + 4 \\ & = \left[\left(5x - \frac{1}{x}\right)^2 + 2\left(5x - \frac{1}{x}\right)\right] + 2\left(5x - \frac{1}{x}\right) + 4 \\ & = \left(5x - \frac{1}{x}\right)\left(5x - \frac{1}{x} + 2\right) + 2\left[5x - \frac{1}{x} + 2\right] \\ & = \left(5x - \frac{1}{x} + 2\right)\left(5x - \frac{1}{x} + 2\right) \end{aligned}$$

Q4. Factorize:

- $(x^2 + 5x + 4)(x^2 + 5x + 6) - 3$
- $(x^2 - 4x)(x^2 - 4x - 1) - 20$
- $(x + 2)(x + 3)(x + 4)(x + 5) - 15$
- $(x + 4)(x - 5)(x + 6)(x - 7) - 504$
- $(x + 1)(x + 2)(x + 3)(x + 6) - 3x^2$

Solution:-

$$\begin{aligned} 4(i) \quad & (x^2 + 5x + 4)(x^2 + 5x + 6) - 3 \\ \text{Let } & x^2 + 5x = y \\ & (x^2 + 5x + 4)(x^2 + 5x + 6) - 3 \end{aligned}$$

MATHEMATICS FOR 9TH CLASS (UNIT # 5)

Pilot Supercone Mathematics 247 Class 9th

$$= (y + 4)(y + 6) - 3$$

$$= y^2 + 10y + 24 - 3$$

$$= y^2 + 10y + 21$$

$$= y^2 + 7y + 3y + 21$$

$$= y(y + 7) + 3(y + 7)$$

$$= (y + 7)(y + 3)$$

$$\text{Putting } y = x^2 + 5x$$

$$= (x^2 + 5x + 7)(x^2 + 5x + 3)$$

$$4(ii) \quad (x^2 - 4x)(x^2 - 4x - 1) - 20$$

$$\text{Let } x^2 - 4x = y$$

$$= (x^2 - 4x)(x^2 - 4x - 1) - 20 \quad (\text{given expression})$$

$$= (y)(y - 1) - 20$$

$$= y^2 - y - 20$$

$$= y^2 - 5y + 4y - 20$$

$$= y(y - 5) + 4(y - 5)$$

$$\text{Putting } y = x^2 - 4x$$

$$= (x^2 - 4x - 5)(x^2 - 4x + 4)$$

$$= (x^2 - 5x + x - 5)[x^2 - 2x - 2x + 4]$$

$$= [x(x - 5) + 1(x - 5)][x(x - 2) - 2(x - 2)]$$

$$= (x - 5)(x + 1)(x - 2)(x - 2)$$

$$= (x - 5)(x + 1)(x - 2)^2$$

$$4(iii) \quad (x + 2)(x + 3)(x + 4)(x + 5) - 15$$

$$\text{Here } 2 + 5 = 3 + 4$$

$$(x + 2)(x + 5)(x + 3)(x + 4) - 15$$

$$= (x^2 + 7x + 10)(x^2 + 7x + 12) - 15$$

$$\text{Let } x^2 + 7x = y$$

$$(x^2 + 7x + 10)(x^2 + 7x + 12) - 15$$

$$= (y + 10)(y + 12) - 15$$

$$= y^2 + 22y + 120 - 15$$

$$= y^2 + 22y + 105$$

$$= y^2 + 15y + 7y + 105$$

$$= y(y + 15) + 7(y + 15)$$

$$= (y + 15)(y + 7)$$

MATHEMATICS FOR 9TH CLASS (UNIT # 5)

Pilot Superone Mathematics 248 Class 9th

$$\begin{aligned}
 & \text{Putting } y = x^2 + 7x \\
 & = (x^2 + 7x + 15)(x^2 + 7x + 7) \\
 4(iv) \quad & (x + 4)(x - 5)(x + 6)(x - 7) \quad 504 \\
 & \text{Here } +4 \quad -5 = 6 - 7 \\
 & (x^2 - x - 20)(x^2 - x - 42) \quad 504 \\
 \text{Let } & x^2 - x = y \\
 & (y - 20)(y - 42) \quad 504 \text{ (given expression)} \\
 & = (y - 20)(y - 42) - 504 \\
 & = y^2 - 62y + 840 - 504 \\
 & = y^2 - 62y + 336 \\
 & = y^2 - 56y - 6y + 336 \\
 & = y(y - 56) - 6(y - 56) \\
 & = (y - 56)(y - 6) \\
 & \text{Putting } y = x^2 - x \\
 & = (x^2 - x - 56)(x^2 - x - 6) \\
 & = (x^2 - 8x + 7x - 56)(x^2 - 3x + 2x - 6) \\
 & = [x(x - 8) + 7(x - 8)][x(x - 3) + 2(x - 3)] \\
 & = (x - 8)(x + 7)(x - 3)(x + 2) \\
 4(v) \quad & (x + 1)(x + 2)(x + 3)(x + 6) \quad 3x^2 \\
 & 1 \times 6 = 2 \times 3 \\
 & (x + 1)(x + 6)(x + 2)(x + 3) \quad 3x^2 \\
 & (x^2 + 7x + 6)(x^2 + 5x + 6) \quad 3x^2 \\
 & (x^2 + 6 + 7x)(x^2 + 6 + 5x) \quad 3x^2 \\
 \text{Let } & x^2 + 6 = y \\
 & = (y + 7x)(y + 5x) - 3x^2 \quad \text{(Expression)} \\
 & = y^2 + 12xy + 35x^2 - 3x^2 \\
 & = y^2 + 12xy + 32x^2 \\
 & = y^2 + 8xy + 4xy + 32x^2 \\
 & = y(y + 8x) + 4x(y + 8x) \\
 & = (y + 8x)(y + 4x) \\
 & \text{Putting } y = x^2 + 6 \\
 & = (x^2 + 6 + 8x)(x^2 + 6 + 4x) \\
 & = (x^2 + 8x + 6)(x^2 + 4x + 6)
 \end{aligned}$$

www.dn

MATHEMATICS FOR 9TH CLASS (UNIT # 5)

Pilot Superone Mathematics 249

Class 9th**Q.5. Factorize:**

(i) $x^3 + 48x - 12x^2 - 64$

(ii) $8x^3 + 60x^2 + 150x + 125$

(iii) $x^3 - 18x^2 + 108x - 216$

(iv) $8x^3 - 125y^3 - 60x^2y + 150xy^2$

Solutions:-

$$\begin{aligned}
 5(i) \quad & x^3 + 48x - 12x^2 - 64 \\
 & = x^3 - 64 - 12x^2 + 48x \\
 & = (x)^3 - (4)^3 - 3(x^2)(4) + 3(x)(4^2) \\
 & = (x - 4)^3 \{ (a - b)^3 = a^3 - 3a^2b + 3ab^2 - b^3 \} \text{ (Formula)}
 \end{aligned}$$

$$\begin{aligned}
 5(ii) \quad & 8x^3 + 60x^2 + 150x + 125 \\
 & = 8x^3 + 125 + 60x^2 + 150x \\
 & = (2x)^3 + (5)^3 + 3(2x)^2(5) + 3(2x)(5)^2 \\
 & = (2x + 5)^3
 \end{aligned}$$

$$\begin{aligned}
 5(iii) \quad & x^3 - 18x^2 + 108x - 216 \\
 & = x^3 - 216 - 18x^2 + 108x \\
 & = (x)^3 - (6)^3 - 3(x)^2(6) + 3(x)(6)^2 \\
 & = (x - 6)^3
 \end{aligned}$$

$$\begin{aligned}
 5(iv) \quad & 8x^3 - 125y^3 - 60x^2y + 150xy^2 \\
 & = (2x)^3 - (5y)^3 - 3(2x)^2(5y) + 3(2x)(5y)^2 \\
 & = (2x - 5y)^3
 \end{aligned}$$

Q.6. Factorize:

(i) $27 + 8x^3$

(ii) $125x^3 - 216y^3$

(iii) $64x^3 - 27y^3$

(iv) $8x^3 + 125y^3$

Solutions:-

$$\begin{aligned}
 6(i) \quad & 27 + 8x^3 \\
 & = (3)^3 + (2x)^3 \\
 & a^3 + b^3 = (a + b)(a^2 - ab + b^2) \quad \text{(Formula)} \\
 & = (3 + 2x) \{ (3)^2 - (3)(2x) + (2x)^2 \}
 \end{aligned}$$

MATHEMATICS FOR 9TH CLASS (UNIT # 5)

Pilot Superone Mathematics 250 Class 9th

$$= (3 + 2x)(9 - 6x + 4x^2)$$

$$\begin{aligned} 6(ii) \quad & 125x^3 - 216y^3 \\ & a^3 - b^3 = (a - b)(a^2 + ab + b^2) \quad \text{(Formula)} \\ & (5x)^3 - (6y)^3 \\ & (5x - 6y) \{ (5x)^2 + (5x)(6y) + (6y)^2 \} \\ & = (5x - 6y)(25x^2 + 30xy + 36y^2) \end{aligned}$$

$$\begin{aligned} 6(iii) \quad & 64x^3 + 27y^3 \\ & (4x)^3 + (3y)^3 \\ & a^3 + b^3 = (a + b)(a^2 - ab + b^2) \quad \text{(Formula)} \\ & = (4x + 3y) \{ (4x)^2 - (4x)(3y) + (3y)^2 \} \\ & = (4x + 3y)(16x^2 - 12xy + 9y^2) \end{aligned}$$

$$\begin{aligned} 6(iv) \quad & 8x^3 - 125y^3 \\ & = (2x)^3 - (5y)^3 \\ & a^3 - b^3 = (a - b)(a^2 + ab + b^2) \quad \text{(Formula)} \\ & = (2x - 5y) \{ (2x)^2 + (2x)(5y) + (5y)^2 \} \\ & = (2x - 5y)(4x^2 + 10xy + 25y^2) \end{aligned}$$

Remainder Theorem

If a polynomial $p(x)$ is divided by a linear divisor $(x - a)$ then the remainder is $p(a)$

www.doc

MATHEMATICS FOR 9TH CLASS (UNIT # 5)

Prior Superong Mathematics 251 Class 9th

Exercise 5.3

Q.1. Use the remainder theorem to find the remainder when

- (i) $3x^3 - 10x^2 + 13x - 6$ is divided by $(x-2)$
 (ii) $4x^3 - 4x + 3$ is divided by $(2x - 1)$
 (iii) $6x^4 + 2x^3 - x + 2$ is divided by $(x + 2)$
 (iv) $(2x - 1)^3 + 6(3 + 4x)^2 - 10$ is divided by $(2x + 1)$
 (v) $x^3 - 3x^2 + 4x - 14$ is divided by $(x + 2)$

Solutions:-

1(i) $p(x) = 3x^3 - 10x^2 + 13x - 6$

From $x - 2 = 0$ $x = 2$

$$p(2) = 3(2)^3 - 10(2)^2 + 13(2) - 6$$

$$= 24 - 40 + 26 - 6$$

$= 4$ (R)

1(ii) $p(x) = 4x^3 - 4x + 3$

From $2x - 1 = 0$ $x = \frac{1}{2}$

$$p\left(\frac{1}{2}\right) = 4\left(\frac{1}{2}\right)^3 - 4\left(\frac{1}{2}\right) + 3$$

$$= 4 \times \frac{1}{8} - 2 + 3$$

$$= \frac{1}{2} - 2 + 3 = \frac{1}{2} + 1 = \frac{3}{2}$$

1(iii) $p(x) = 6x^4 + 2x^3 - x + 2$

From $x + 2 = 0$ $x = -2$

$$p(-2) = 6(-2)^4 + 2(-2)^3 - (-2) + 2$$

$$= 6(16) + 2(-8) + 2 + 2$$

$$= 96 - 16 + 2 + 2$$

$$= 84$$
 (R)

1(v) $p(x) = (2x - 1)^3 + 6(3 + 4x)^2 - 10$

From $2x + 1 = 0$, $x = -\frac{1}{2}$

MATHEMATICS FOR 9TH CLASS (UNIT # 5)

1. Ekst Superone Mathematics 252 Class 9th

$$\begin{aligned} p\left(-\frac{1}{2}\right) &= \left[2\left(-\frac{1}{2}\right) - 1\right]^3 + 6\left[3 + 4\left(-\frac{1}{2}\right)\right]^2 - 10 \\ &= [-1 - 1]^3 + 6[3 - 2]^2 - 10 \\ &= (-2)^3 + 6(1)^2 - 10 \\ &= -8 + 6 - 10 \\ &= -12 \text{ (R)} \end{aligned}$$

$$\begin{aligned} \text{1(v)} \quad p(x) &= x^3 - 3x^2 + 4x - 14 \\ \text{From } x + 2 &= 0, x = -2 \\ p(-2) &= (-2)^3 - 3(-2)^2 + 4(-2) - 14 \\ &= -8 - 12 - 8 - 14 \\ &= -42 \text{ (R)} \end{aligned}$$

- Q.2. (i) If $(x + 2)$ is a factor of $3x^2 - 4kx - 4k^2$ then find the value(s) of k .
 (ii) If $(x - 1)$ is a factor of $x^3 - kx^2 + 11x - 6$, then find the value of k .

Solutions:-

$$\begin{aligned} \text{2(i)} \quad p(x) &= 3x^2 - 4kx - 4k^2 \\ \text{From } x + 2 &= 0, x = -2 \\ p(-2) &= 3(-2)^2 - 4k(-2) - 4k^2 \\ &= 3(4) + 8k - 4k^2 \\ &= 12 + 8k - 4k^2 \text{ (R)} \end{aligned}$$

∵ If $(x + 2)$ is a factor then $R = 0$

$$\begin{aligned} \text{Thus } 12 + 8k - 4k^2 &= 0 && \text{divided by 4} \\ 3 + 2k - k^2 &= 0 \\ 3 + 3k - k - k^2 &= 0 \\ 3(1 + k) - k(1 + k) &= 0 \\ (1 + k)(3 - k) &= 0 \end{aligned}$$

$$\begin{aligned} \text{From } 1 + k &= 0 \\ k &= -1 \\ \text{and } 3 - k &= 0 \\ k &= 3 \\ \text{give } k &= 3, -1 \end{aligned}$$

MATHEMATICS FOR 9TH CLASS (UNIT # 5)

Pilot Superone Mathematics 253 Class 9th

2(ii) $p(x) = x^3 - kx^2 + 11x - 6$

From $x - 1 = 0$ $x = 1$

$p(1) = (1)^3 - k(1)^2 + 11(1) - 6$

$= 1 - k + 11 - 6$

$= 6 - k \quad (R)$

If $x - 1$ is factor then $R = 0$

From

$6 - k = 0$

$k = 6$

Q.3. Without actual long division determine whether

(i) $(x - 2)$ and $(x - 3)$ are factors of $p(x) = x^3 - 12x^2 + 44x - 48$

(ii) $(x - 2)$, $(x + 3)$ and $(x - 1)$ are factors of $q(x) = x^3 + 2x^2 - 5x - 6$

3(i) $p(x) = x^3 - 12x^2 + 44x - 48$

From $x - 2 = 0$, $x = 2$

$p(2) = (2)^3 - 12(2)^2 + 44(2) - 48$

$= 8 - 48 + 88 - 48$

$= 96 - 96 = 0 \quad (R)$

$R = 0$ therefore, $x - 2$ is a factor of the polynomial

$p(x) = x^3 - 12x^2 + 44x - 48$

From $(x - 3)$, $x = 3$, $x - 3 = 0$

$p(3) = (3)^3 - 12(3)^2 + 44(3) - 48$

$= 27 - 108 + 132 - 48$

$R = 159 - 156 = 3 \neq 0$

$R \neq 0$, therefore, $x - 3$ is NOT factor

(ii) $p(x) = x^3 + 2x^2 - 5x - 6$

From $x - 2 = 0$, $x = 2$

$p(2) = (2)^3 + 2(2)^2 - 5(2) - 6$

$= 8 + 8 - 10 - 6$

$= 16 - 16 = 0 \quad (R)$

$R = 0$, therefore, $x - 2$ is factor

$p(x) = x^3 + 2x^2 - 5x - 6$

from $x + 3 = 0$, $x = -3$

MATHEMATICS FOR 9TH CLASS (UNIT # 5)

Pilot Supreme Mathematics 254 Class 9th

$$\begin{aligned} p(-3) &= (-3)^3 + 2(-3)^2 - 5(-3) - 6 \\ &= -27 + 18 + 15 - 6 \\ &= 33 - 33 = 0 \quad (R) \end{aligned}$$

∴ $R = 0$, $x + 3$ is factor

$$p(x) = x^3 + 2x^2 - 5x - 6$$

$$\text{From } (x - 4) = 0, x = 4$$

$$\begin{aligned} p(4) &= (4)^3 + 2(4)^2 - 5(4) - 6 \\ &= 64 + 32 - 20 - 6 \\ &= 96 - 26 \neq 0 \quad (R) \end{aligned}$$

$R \neq 0$, therefore, $x - 4$ is not factor

Q.4. For what value of m is the polynomial $p(x) = 4x^3 - 7x^2 + 6x - 3m$ exactly divisible by $x + 2$?

Solution:-

$$p(x) = 4x^3 - 7x^2 + 6x - 3m$$

$$\text{From } x + 2 = 0, x = -2$$

$$\begin{aligned} p(-2) &= 4(-2)^3 - 7(-2)^2 + 6(-2) - 3m \\ &= -32 - 28 - 12 - 3m \\ &= -72 - 3m \quad (R) \end{aligned}$$

If $x + 2$ is factor then $R = 0$

$$-72 - 3m = 0$$

$$\text{or } 24 + m = 0$$

$$\text{thus } m = -24$$

Q.5. Determine the value of k if $p(x) = kx^3 + 4x^2 + 3x - 4$ and $q(x) = x^3 - 4x + k$ leaves the same remainder when divided by $(x - 3)$.

$$p(x) = kx^3 + 4x^2 + 3x - 4$$

$$\text{From } x - 3 = 0, x = 3$$

$$\begin{aligned} p(3) &= k(3)^3 + 4(3)^2 + 3(3) - 4 \\ &= 27k + 36 + 9 - 4 \\ &= 27k + 41 \quad (R_1) \end{aligned}$$

$$\text{Now } q(x) = x^3 - 4x + k$$

$$\text{and } q(3) = (3)^3 - 4(3) + k$$

MATHEMATICS FOR 9TH CLASS (UNIT # 5)

Pilot Superone Mathematics 255 Class 9th

$$= 27 - 12 + k$$

$$= 15 + k$$

(R₂)

Now $R_1 = R_2$

$$27k + 41 = 15 + k$$

$$27k - k = 15 - 41$$

$$26k = -26$$

thus $k = -1$

Q.6. The remainder after dividing the polynomial $p(x) = x^3 + 9x^2 + 7$ by $(x + 1)$ is $2b$. Calculate the value of a and b if this expression leaves a remainder of $(b + 5)$ on being divided by $(x - 2)$.

Solution:-

$$p(x) = x^3 + ax^2 + 7$$

From $x + 1 = 0$, $x = -1$

$$p(-1) = (-1)^3 + a(-1)^2 + 7$$

$$= -1 + a + 7$$

$$= a + 6$$

$$a + 6 = 2b \quad (1^{st} \text{ Condition})$$

Now $p(x) = x^3 + ax^2 + 7$

From $(x - 2) = 0$, $x = 2$

and $p(2) = (2)^3 + a(2)^2 + 7$

$$= 8 + 4a + 7$$

$$4a + 15$$

$$4a + 15 = b + 5 \quad (2^{nd} \text{ Condition})$$

$$4a - b = 5 - 15$$

$$4a - b = -10 \quad (B)$$

$$a - 2b = 6 \quad (i) \text{ From A, B}$$

$$8a - 2b = 20 \quad (ii) \text{ mult.plying B by 2}$$

$$7a = -14$$

subtracting (i) from (ii)

$$a = \frac{14}{7}$$

$$a = 2$$

$$a - 2b = -6 \quad \text{From A}$$

MATHEMATICS FOR 9TH CLASS (UNIT # 5)

Pilot Superone Mathematics 256 Class 9th

$$-2 - 2b = -6 \quad (\text{Putting } a = -6)$$

$$-2b = -6 + 2$$

$$2b = -4$$

$$b = -2$$

Thus $a = -6, b = -2$

Q.7. The polynomial $x^3 + lx^2 + mx + 24$ has a factor $(x + 4)$ and it leaves a remainder of 36 when divided by $(x - 2)$. Find the values of l and m .

Solution:-

$$p(x) = x^3 + lx^2 + mx + 24$$

$$\text{From } x + 4 = 0, x = -4$$

$$p(-4) = (-4)^3 + l(-4)^2 + m(-4) + 24$$

$$= -64 + 16l - 4m + 24$$

$$= 16l - 4m - 40$$

$$16l - 4m - 40 = 0 \quad (1^{\text{st}} \text{ Condition})$$

$$\text{or } 4l - m - 10 = 0$$

$$\text{or } 4l - m = 10 \quad (A)$$

$$\text{Now } p(x) = x^3 + lx^2 + mx + 24$$

$$\text{From } x - 2 = 0, x = 2$$

$$p(2) = (2)^3 + l(2)^2 + m(2) + 24$$

$$= 8 + 4l + 2m + 24$$

$$= 4l + 2m + 32$$

$$4l + 2m + 32 = 36 \quad (2^{\text{nd}} \text{ Condition})$$

$$\text{or } 4l + 2m = 36 - 32$$

$$\text{or } 4l + 2m = 4$$

$$\text{or } 2l + m = 2 \quad (B)$$

$$4l - m = 10 \quad \text{From (A)}$$

$$6l = 12 \quad \text{From A + B}$$

$$l = 2$$

Putting $l = 2$ in A

$$4l - m = 10$$

$$4(2) - m = 10$$

$$8 - m = 10$$

MATHEMATICS FOR 9TH CLASS (UNIT # 5)

Pilot Superone Mathematics 257 Class 9th

$$-m = 10 - 8$$

$$m = 2$$

$$m = 2$$

thus $l = 2, m = 2$

Q.8. The expression $lx^3 + mx^2 - 4$ leaves remainder of -3 and 12 when divided by $(x - 1)$ and $(x + 2)$ respectively. Calculate the values of l and m .

Solution.

$$p(x) = lx^3 + mx^2 - 4$$

$$\text{From } x - 1 = 0, x = 1$$

$$p(1) = l(1)^3 + m(1)^2 - 4$$

$$= l + m - 4$$

$$l + m - 4 = -3 \quad (1^{\text{st}} \text{ Condition})$$

$$l + m = -3 + 4$$

$$l + m = 1 \quad (A)$$

$$p(x) = lx^3 + mx^2 - 4$$

$$\text{From } x + 2 = 0, x = -2$$

$$p(-2) = l(-2)^3 + m(-2)^2 - 4$$

$$= -8l + 4m - 4$$

$$-8l + 4m - 4 = 12 \quad (2^{\text{nd}} \text{ Condition})$$

$$-8l + 4m = 12 + 4$$

$$-8l + 4m = 16$$

$$2l - m = -4 \quad (B)$$

$$l - m = -1 \quad \text{From A}$$

$$3l = -3 \quad \text{From A + B}$$

$$l = -1$$

$$l + m = 1 \quad (\text{From A})$$

$$\text{Putting } l = -1$$

$$-1 + m = 1$$

$$m = 1 + 1$$

$$m = 2$$

$$\text{Thus } l = -1, m = 2$$

MATHEMATICS FOR 9TH CLASS (UNIT # 5)

Pilot Supreme Mathematics 258 Class 9th

Q.9. The expression $ax^3 - 9x^2 + bx + 3a$ is exactly divisible by $x^2 - 5x + 6$. Find the values of a and b .

Solution:-

$$\begin{aligned} \text{Factorizing: } x^2 - 5x + 6 \\ &= x^2 - 2x - 3x + 6 \\ &= x(x - 2) - 3(x - 2) \\ &= (x - 2)(x - 3) \end{aligned}$$

If $x^2 - 5x + 6$ divides the expression $ax^3 - 9x^2 + bx + 3a$ then $x - 2$ and $x - 3$ will also divide it.

$$\begin{aligned} p(x) &= ax^3 - 9x^2 + bx + 3a \\ \text{When } x - 2 &= 0; x = 2 \\ p(2) &= a(2)^3 - 9(2)^2 + b(2) + 3a \\ &= 8a - 36 + 2b + 3a \\ &= 11a + 2b - 36 \end{aligned}$$

$$11a + 2b - 36 = 0 \quad (1^{\text{st}} \text{ Condition})$$

$$11a + 2b = 36 \quad (A)$$

$$p(x) = ax^3 - 9x^2 + bx + 3a$$

$$\begin{aligned} p(3) &= a(3)^3 - 9(3)^2 + b(3) + 3a \\ &= 27a - 81 + 3b + 3a \\ &= 30a + 3b - 81 \end{aligned}$$

$$30a + 3b - 81 = 0 \quad (2^{\text{nd}} \text{ Condition})$$

$$30a + 3b = 81$$

$$10a + b = 27 \quad (B) \text{ Dividing by 3}$$

$$11a + 2b = 36 \quad A$$

$$10a + b = 27 \quad B$$

$$20a + 2b = 54 \quad (i) \text{ Multiplying } B \text{ by } 2$$

$$a \downarrow 11a + 2b = 36 \quad (ii) \text{ Subtracting (i) from (ii)}$$

$$9a = 18$$

$$a = 2$$

$$10a + b = 27 \quad \text{From } B$$

$$10(2) + b = 27 \quad \text{Putting } a = 2$$

$$20 + b = 27$$

$$b = 27 - 20$$

$$\text{Thus } a = 2, b = 7$$

MATHEMATICS FOR 9TH CLASS (UNIT # 5)

Pilot Superone Mathematics

259

Class 9th

Exercise 5.4

Factorize each of the following cubic polynomials by factor theorem.

- | | |
|---------------------------|---------------------------|
| 1. $x^3 - 2x^2 - x + 2$ | 2. $x^3 - x^2 - 22x + 40$ |
| 3. $x^3 - 6x^2 + 3x + 10$ | 4. $x^3 + x^2 - 10x + 8$ |
| 5. $x^3 - 2x^2 - 5x + 6$ | 6. $x^3 + 5x^2 - 2x - 24$ |
| 7. $3x^3 - x^2 - 12x + 4$ | 8. $2x^3 + x^2 - 2x - 1$ |

Solutions:-

1. $P(x) = x^3 - 2x^2 - x + 2$

$p = 2$ and possible factors of 2 are $\pm 1 \pm 2$

$q = 1$ and possible factor of 1 are 1 possible factors of $P(x)$ will be from

$$\frac{p}{q} = \pm 1, \pm 2$$

$$P(x) = x^3 - 2x^2 - x + 2$$

$$P(1) = (1)^3 - 2(1)^2 - (1) + 2 \\ = 1 - 2 - 1 + 2 = 0 \quad \text{Remainder}$$

$x - 1$ is one factor

$$P(-1) = (-1)^3 - 2(-1)^2 - (-1) + 2 \\ = -1 - 2 + 1 + 2 = 0 \quad \text{(Remainder)}$$

$x + 1$ is another factor

$$P(2) = (2)^3 - 2(2)^2 - (2) + 2 \\ = 8 - 8 - 2 + 2 = 0 \quad \text{(R)}$$

$x - 2$ is third factor

Thus

$$x^3 - 2x^2 - x + 2 = (x - 1)(x + 1)(x - 2)$$

2.

$$P(x) = x^3 - x^2 - 22x + 40$$

$p = 40$, possible factors of 40 are $\pm 1, \pm 2, \pm 4, \pm 5, \pm 8, \pm 10, \pm 20$

$q = 1$, possible factor of 1 are ± 1 , $P(x) = 0$: possible factors of $P(x)$ will be from

MATHEMATICS FOR 9TH CLASS (UNIT # 5)

Pilot Superintend Mathematics 260 Class 9th

$$\frac{P}{q} = \pm 1, \pm 2, \pm 4, \pm 5, \pm 8, \pm 10, \pm 20$$

$$P(x) = x^3 - x^2 - 22x + 40$$

Put $x = 2$

$$P(2) = (2)^3 - (2)^2 - 22(2) + 40 \\ = 8 - 4 - 44 + 40 = 0 \quad (R)$$

$x = 2$ is factor of $P(x)$

Put $x = 4$

$$P(4) = (4)^3 - (4)^2 - 22(4) + 40 \\ = 64 - 16 - 88 + 40 \\ = 104 - 104 = 0 \quad (R)$$

$x = 4$ is also factor of $P(x)$

Put $x = 5$

$$P(5) = (5)^3 - (5)^2 - 22(5) + 40 \\ = 125 - 25 - 110 + 40 \\ = 165 - 135 \neq 0 \quad (R)$$

$(x - 5)$ is not factor of $P(x)$

Put $x = -5$

$$P(-5) = (-5)^3 - (-5)^2 - 22(-5) + 40 \\ = -125 - 25 + 110 + 40 \\ = -150 + 150 = 0 \quad (R)$$

$x + 5$ is also factor of $P(x)$

Thus

$$x^3 - x^2 - 22x + 40 = (x - 2)(x - 4)(x + 5)$$

$$P(x) = x^3 - 6x^2 + 3x + 40$$

$P = 10$, possible factor of 10 are $\pm 1, \pm 2, \pm 5, \pm 10$

$q = 1$ factor of 1 are ± 1 possible factors of $P(x)$ can be found from

$$\frac{P}{q} = \pm 1, \pm 2, \pm 5, \pm 10$$

$$P(x) = x^3 - 6x^2 + 3x + 10$$

Put $x = 1$

$$P(1) = (1)^3 - (1)^2 - 6(1) + 10$$

MATHEMATICS FOR 9TH CLASS (UNIT # 5)

Pilot Superong Mathematics 261 Class 9th

$$1^3 - 6 \cdot 1 + 3 \cdot 10$$

$$= 1 - 6 + 30 \neq 0 \quad (R)$$

$\therefore 1$ is not factor

Put $x = -1$

$$P(x) = (-1)^3 - 6(-1)^2 + 3(-1) + 10$$

$$= -1 - 6 - 3 + 10 = 0 \quad (R)$$

$(x + 1)$ is factor

$$P(x) = x^3 - 6x^2 + 3x + 10$$

Put $x = 2$

$$P(2) = (2)^3 - 6(2)^2 + 3(2) + 10$$

$$= 8 - 24 + 6 + 10$$

$$= 8 - 24 + 16 = 0 \quad (R)$$

$x - 2$ factor

Put $x = 5$

$$P(5) = (5)^3 - 6(5)^2 + 3(5) + 10$$

$$= 125 - 150 + 15 + 10$$

$$= 125 - 150 + 25 = 0 \quad (R)$$

$x - 5$ also factor

Thus

$$x^3 - 6x^2 + 3x + 10 = (x + 1)(x - 2)(x - 5)$$

$$P(x) = x^3 + x^2 - 10x + 8$$

$P = 8$, factor of 8 are $\pm 1, \pm 2, \pm 4, \pm 8$

$q = 1$ factor of 1 are ± 1 possible factors can be found from

$$\frac{p}{q} = \pm 1, \pm 2, \pm 4, \pm 8$$

$$P(x) = x^3 + x^2 - 10x + 8$$

Put $x = 1$

MATHEMATICS FOR 9TH CLASS (UNIT # 5)

Pilot Superone Mathematics 262 Class 9th

$$P(1) = (1)^3 + (1)^2 - 10(1) + 8$$

$$= 1 + 1 - 10 + 8 = 0 \quad (R)$$

$x - 1$ is factor of $P(x)$

$$P(x) = x^3 + x^2 - 10x + 8$$

Put $x = 2$

$$P(2) = (2)^3 + (2)^2 - 10(2) + 8$$

$$= 8 + 4 - 20 + 8$$

$$= 20 - 20 = 0 \quad (R)$$

$x - 2$ is also factor

$$P(x) = x^3 + x^2 - 10x + 8$$

Put $x = -4$

$$P(-4) = (-4)^3 + (-4)^2 - 10(-4) + 8$$

$$= -64 + 16 - 40 + 8$$

$$= -64 + 64 = 0 \quad (R)$$

$x + 4$ is factor

Thus

$$x^3 + x^2 - 10x + 8 = (x - 1)(x - 2)(x + 4)$$

5.

$$P(x) = x^3 - 2x^2 - 5x + 6$$

$P = 6$, factor of 6 are $\pm 1, \pm 2, \pm 3, \pm 6$

$q = 1$, factor of 1 are ± 1 members of factors of $P(x)$
 can be found from

$$\frac{p}{q} = \pm 1, \pm 2, \pm 3, \pm 6$$

$$P(x) = x^3 - 2x^2 - 5x + 6$$

Put $x = 1$

$$P(1) = (1)^3 - 2(1)^2 - 5(1) + 6$$

$$= 1 - 2 - 5 + 6$$

$$= 7 - 7 = 0 \quad \checkmark$$

MATHEMATICS FOR 9TH CLASS (UNIT # 5)

Pilot Superone Mathematics 263

Class 9th

$x - 1$ is factor of $P(x)$

$$P(x) = x^3 - 2x^2 - 5x + 6$$

Put $x = 2$

$$P(2) = (+2)^3 - 2(+2)^2 - 5(+2) + 6$$

$$= +8 - 8 - 10 + 6$$

$$= 14 - 18 = -4 \neq 0 \quad (R)$$

$x - 2$ is not factor of $P(x)$

$$P(x) = x^3 - 2x^2 - 5x + 6$$

Put $x = 3$

$$P(3) = (3)^3 - 2(3)^2 - 5(3) + 6$$

$$= 27 - 18 - 15 + 6$$

$$= 33 - 33 = 0 \quad (R)$$

$x - 3$ is also factor of $P(x)$

Thus

$$P(x) = x^3 - 2x^2 - 5x + 6$$

Put $x = -2$

$$P(-2) = (-2)^3 - 2(-2)^2 - 5(-2) + 6$$

$$= -8 - 8 + 10 + 6$$

$$= -16 + 16 = 0 \quad (R)$$

$x + 2$ is one factor of $P(x)$

Thus

$$x^3 - 2x^2 - 5x + 6 = (x - 1)(x - 3)(x + 2)$$

$$P(x) = x^3 + 5x^2 - 2x - 24$$

$P = -24$, factor of 6 are $\pm 1, \pm 2, \pm 3, \pm 4, \pm 6, \pm 8, \pm 12, \pm 24$

$q = 1$, factor of q are ± 1 possible factors of $P(x)$ will be found from

MATHEMATICS FOR 9TH CLASS (UNIT # 5)

P. of Superior Mathematics 364 Class 9th

$$\frac{P}{q} = \pm 1, \pm 2, \pm 3, \pm 4, \pm 6, \pm 8, \pm 12, \pm 24$$

$$P(x) = x^3 + 5x^2 - 2x - 24$$

Put $x = 2$

$$\begin{aligned} P(2) &= (2)^3 + 5(2)^2 - 2(2) - 24 \\ &= 8 + 20 - 4 - 24 \\ &= 28 - 28 = 0 \end{aligned} \quad (R)$$

$(x - 2)$ is factor of $P(x)$

$$P(x) = x^3 + 5x^2 - 2x - 24$$

Put $x = -3$

$$\begin{aligned} P(-3) &= (-3)^3 + 5(-3)^2 - 2(-3) - 24 \\ &= -27 + 45 + 6 - 24 \\ &= 51 - 51 = 0 \end{aligned} \quad (R)$$

$x + 3$ is factor of $P(x)$

$$P(x) = x^3 + 5x^2 - 2x - 24$$

Put $x = 4$

$$\begin{aligned} P(4) &= (4)^3 + 5(4)^2 - 2(4) - 24 \\ &= 64 + 80 - 8 - 24 \\ &= 144 - 32 \neq 0 \end{aligned} \quad (R)$$

$x - 4$ is not factor of $P(x)$

Put $x = -4$

$$\begin{aligned} P(-4) &= (-4)^3 + 5(-4)^2 - 2(-4) - 24 \\ &= -64 + 80 + 8 - 24 \\ &= 18 - 18 = 0 \end{aligned} \quad (R)$$

$x + 4$ is factor of $P(x)$

$$x^3 + 5x^2 - 2x - 24 = (x - 2)(x^2 + 7x + 12)$$

$$P(x) = 3x^2 - x^2 - 12x + 12$$

MATHEMATICS FOR 9TH CLASS (UNIT # 5)

Pilot Supreme Mathematics 265 Class 9th

P 4 factor of 4 are $\pm 1, \pm 2, \pm 4$

q 3 factor of 3 are $\pm 1, \pm 3$ possible factors of $P(x)$

can be found from $\frac{P}{q}$

$$\frac{P}{q} = \pm 1, \pm 2, \pm 3, \pm 4, \pm \frac{1}{3}, \pm \frac{2}{3}, \pm \frac{4}{3}$$

$$P(x) = 3x^3 - x^2 - 12x + 4$$

Put $x = 2$

$$\begin{aligned} P(2) &= 3(2)^3 - (2)^2 - 12(2) + 4 \\ &= 24 - 4 - 24 + 4 \\ &= 28 - 28 = 0 \end{aligned} \quad (R)$$

$x - 2$ is factor of $P(x)$

Put $x = -2$ in $P(x)$

$$\begin{aligned} P(-2) &= 3(-2)^3 - (-2)^2 - 12(-2) + 4 \\ &= 24 - 4 + 24 + 4 \\ &= -28 + 28 = 0 \end{aligned} \quad (R)$$

$x + 2$ is factor of $P(x)$

Put $x = \frac{1}{2}$

$$\begin{aligned} P\left(\frac{1}{2}\right) &= 3\left(\frac{1}{2}\right)^3 - \left(\frac{1}{2}\right)^2 - 12\left(\frac{1}{2}\right) + 4 \\ &= \frac{3}{8} - \frac{1}{4} - 6 + 4 \neq 0 \end{aligned} \quad (R)$$

$2x - 1$ is not factor of $P(x)$

Put $x = \frac{1}{3}$

$$\begin{aligned} P\left(\frac{1}{3}\right) &= 3\left(\frac{1}{3}\right)^3 - \left(\frac{1}{3}\right)^2 - 12\left(\frac{1}{3}\right) + 4 \\ &= 3 \times \frac{1}{27} - \frac{1}{9} - 4 + 4 = 0 \end{aligned} \quad (R)$$

$x = \frac{1}{3}$ gives $3x - 1 = 0$

MATHEMATICS FOR 9TH CLASS (UNIT # 5)

Pilot Superone Mathematics 266

Class 9th

$3x - 1$ is factor of $P(x)$

Thus

$$3x^3 - x^2 - 12x + 4 = (x - 2)(x + 2)(3x - 1)$$

8. $P(x) = 2x^3 + x^2 - 2x - 1$

$P = -1$, factor of -1 are ± 1

$q = 2$, factor of 2 are $\pm 1, \pm 2$ possible factors of $P(x)$

will be found from $\frac{P}{q}$

$$\frac{P}{q} = \pm 1, \pm \frac{1}{2}$$

$$P(x) = 2x^3 + x^2 - 2x - 1$$

Put $x = 1$

$$\begin{aligned} P(1) &= 2(1)^3 + (1)^2 - 2(1) - 1 \\ &= 2 + 1 - 2 - 1 = 0 \end{aligned} \quad (R)$$

$x + 1$ is factor of $P(x)$

Put $x = -1$

$$\begin{aligned} P(-1) &= 2(-1)^3 + (-1)^2 - 2(-1) - 1 \\ &= -2 + 1 + 2 - 1 = 0 \end{aligned} \quad (R)$$

$x + 1$ is factor of $P(x)$

Put $x = -\frac{1}{2}$

$$\begin{aligned} P(-\frac{1}{2}) &= 2(-\frac{1}{2})^3 + (-\frac{1}{2})^2 - 2(-\frac{1}{2}) - 1 \\ &= -2 \times \frac{1}{8} + \frac{1}{4} + 1 - 1 \\ &= -\frac{1}{4} + \frac{1}{4} + 1 - 1 = 0 \end{aligned} \quad (R)$$

From $x = -\frac{1}{2}$, $2x + 1$ is factor of $P(x)$

Thus

$$2x^3 - x^2 - 2x - 1 = (x - 1)(x + 1)(2x + 1)$$

**Unit
6**

ALGEBRAIC MANIPULATION

**Highest Common Factor (H.C.F) and Least
Common Multiple (L.C.M) of Algebraic Expressions**

Highest Common Factor (H.C.F):

If two or more algebraic expressions are given then their common factor of highest power is called the H.C.F. of the expressions.

Least Common Multiple (L.C.M.)

If an algebraic expression $p(x)$ is exactly divisible by two or more expressions, then $p(x)$ is called the Common Multiple of the given expressions.

The Least Common Multiple (L.C.M) is the product of common factor of the given expressions.

Square Root:

Square root of an algebraic expression $p(x)$ will be another algebraic expression $q(x)$ if $[q(x)] \times [q(x)] = p(x)$

Exercise 6.1

Q.1 Find the H.C.F. of the following expressions.

(i) $39x^7y^3z, 91x^5y^6z^7$

(ii) $102xy^2z, 85x^2yz, 187xyz^2$

1(i) $39x^7y^3z, 91x^5y^6z^7$

Solution:

$$39x^7y^3z = 3 \times 13 x^5 x^2 y^3 z$$

$$91x^5y^6z^7 = 7 \times 13 x^5 y^3 y^3 z z^6$$

$$\text{H.C.F.} = 13x^5y^3z$$

1(ii) $102xy^2z, 85x^2yz, 187xyz^2$

Solution:

$$102xy^2z = 2 \times 3 \times 17 xyyz$$

$$85x^2yz = 5 \times 17 xxyz$$

MATHEMATICS FOR 9TH CLASS (UNIT # 6)

Pilot Superone Mathematics 272 Class 9th

$$187xyz^2 = 11 \times 17xyz$$

$$\text{H.C.F.} = 17xyz$$

Q.2 Find the H.C.F. of the following expressions by factorization.

(i) $x^2 + 5x + 6$, $x^2 - 4x - 12$

(ii) $x^3 - 27$, $x^2 + 6x - 27$, $2x^2 - 18$

(iii) $x^3 - 2x^2 + x$, $x^2 + 2x - 3$, $x^2 + 3x - 4$

(iv) $18(x^3 - 9x^2 + 8x)$, $24(x^2 - 3x + 2)$

(v) $36(3x^4 + 5x^3 - 2x^2)$, $54(27x^4 - x)$

2(i) $x^2 + 5x + 6$, $x^2 - 4x - 12$

Solution. $x^2 + 5x + 6 = x^2 + 2x + 3x + 6$
 $= x(x + 2) + 3(x + 2)$
 $= (x + 2)(x + 3)$

and $x^2 - 4x - 12 = x^2 - 6x + 2x - 12$
 $= x(x - 6) + 2(x - 6)$
 $= (x - 6)(x + 2)$

Now $x^2 + 5x + 6 = (x + 2)(x + 3)$
 $x^2 - 4x - 12 = (x - 6)(x + 2)$
 $\text{H.C.F.} = x + 2$

2(ii) $x^3 - 27$, $x^2 + 6x - 27$, $2x^2 - 18$

Solution: $x^3 - 27 = (x)^3 - (3)^3$
 $= (x - 3)[(x)^2 + (x)(3) + (3)^2]$
 $= (x - 3)(x^2 + 3x + 9)$

$x^2 + 6x - 27 = x^2 + 9x - 3x - 27$
 $= x(x + 9) - 3(x + 9)$
 $= (x - 3)(x + 9)$

$2x^2 - 18 = 2(x^2 - 9)$
 $= 2[(x)^2 - (3)^2]$
 $= 2(x - 3)(x + 3)$

MATHEMATICS FOR 9TH CLASS (UNIT # 6)

Pilot Superone Mathematics 273 Class 9th

Now $x^3 - 27 = (x - 3)(x^2 + 3x + 9)$

$x^2 + 6x - 27 = (x - 3)(x + 9)$

$2x^2 - 18 = 2(x - 3)(x + 3)$

H.C.F. = $(x - 3)$

2(iii) $x^3 - 2x^2 + x$, $x^2 + 2x - 3$, $x^2 + 3x - 4$

Solution. $x^3 - 2x^2 + x = x(x^2 - 2x + 1)$
 $= x(x^2 - x - x + 1)$
 $= x[x(x - 1) - 1(x - 1)]$
 $= x(x - 1)(x - 1)$

$x^2 + 2x - 3 = x^2 + 3x - x - 3$
 $= x(x + 3) - 1(x + 3)$
 $= (x - 1)(x + 3)$

$x^2 + 3x - 4 = x^2 + 4x - x - 4$
 $= x(x + 4) - 1(x + 4)$
 $= (x - 1)(x + 4)$

Now $x^3 - 2x^2 + x = x(x - 1)(x - 1)$

$x^2 + 2x - 3 = (x - 1)(x + 3)$

$x^2 + 3x - 4 = (x - 1)(x + 4)$

H.C.F. = $x - 1$

2(iv) $18(x^3 - 9x^2 + 8x)$, $24(x^2 - 3x + 2)$

Solution: $18(x^3 - 9x^2 + 8x) = 2 \times 3 \times 3x(x^2 - 9x + 8)$
 $= 2 \times 3 \times 3x[x^2 - x - 8x + 8]$
 $= 2 \times 3 \times 3x[x(x - 1) - 8(x - 1)]$
 $= 2 \times 3 \times 3x(x - 1)(x - 8)$

$24(x^2 - 3x + 2) = 2 \times 2 \times 2 \times 3[x^2 - x - 2x + 2]$
 $= 2 \times 2 \times 2 \times 3[x(x - 1) - 2(x - 1)]$
 $= 2 \times 2 \times 2 \times 3(x - 1)(x - 2)$

Now $18(x^3 - 9x^2 + 8x) = 2 \times 3 \times 3x(x - 1)(x - 8)$

$24(x^2 - 3x + 2) = 2 \times 2 \times 2 \times 3(x - 1)(x - 2)$

H.C.F. = $2 \times 3(x - 1)$

= $6(x - 1)$

MATHEMATICS FOR 9TH CLASS (UNIT # 6)

Pilot Superone Mathematics 274 Class 9th

2(v) $36(3x^4 + 5x^3 - 2x^2)$, $54(27x^4 - x)$

Solution $36(3x^4 + 5x^3 - 2x^2) = 4 \times 9 \times 2(3x^2 + 5x - 2)$
 $= 4 \times 9 \times 2(3x^2 + 6x - x - 2)$
 $= 4 \times 9 \times 2(3x(x + 2) - 1(x + 2))$
 $= 2 \times 2 \times 3 \times 3 \times 2(x + 2)(3x - 1)$
 $54(27x^4 - x) = 6 \times 9 \times (27x^3 - 1)$
 $= 6 \times 9 \times ((3x)^3 - (1)^3)$
 $= 2 \times 3 \times 3 \times 3 \times (3x - 1)(9x^2 + 3x + 1)$

Now $36(3x^4 + 5x^3 - 2x^2) = 2 \times 2 \times 3 \times 3 \times 2(x + 2)(3x - 1)$
 $54(27x^4 - x) = 2 \times 3 \times 3 \times 3 \times (3x - 1)(9x^2 + 3x + 1)$
 H.C.F. $= 2 \times 3 \times 3(3x - 1)$
 $= 18x(3x - 1)$

Q.3 Find the H.C.F. of the following by division method.

(i) $x^3 + 3x^2 - 16x + 12$, $x^3 + x^2 - 10x + 8$

(ii) $x^4 + x^3 - 2x^2 + x - 3$, $5x^3 + 3x^2 - 17x + 6$

(iii) $2x^5 - 4x^4 - 6x$, $x^5 + x^4 - 3x^3 - 3x^2$

3(i) $x^3 + 3x^2 - 16x + 12$, $x^3 + x^2 - 10x + 8$

Solution:

$$\begin{array}{r}
 x^3 + x^2 - 10x + 8 \overline{) x^3 + 3x^2 - 16x + 12} \\
 \underline{+x^3 + x^2 + 10x + 8} \\
 2x^2 - 6x + 4 \\
 2 \overline{) 2x^2 - 6x + 4} \\
 \underline{x^3 + x^2 - 10x + 8} \quad x+4 \\
 \underline{+x^3 \quad 3x^2 + 2x + 8} \\
 4x^2 - 12x + 8 \\
 \underline{4x^2 - 12x + 8} \\
 0
 \end{array}$$

H.C.F. $= x^2 - 3x + 2$

MATHEMATICS FOR 9TH CLASS (UNIT # 6)

Pilot Superone Mathematics 275 Class 9th
 3(ii) $x^4 + x^3 - 2x^2 + x - 3$, $5x^3 + 3x^2 - 17x + 6$

Solution.

$$\begin{array}{r} 5x^3 + 3x^2 - 17x + 6 \quad | \quad x^4 + x^3 - 2x^2 + x - 3 \\ \underline{+ 5x^4 + 5x^3 - 10x^2 + 5x - 15} \\ 10x^3 + 8x^2 - 16x + 21 \\ \underline{+ 5x^4 + 3x^3 - 17x^2 + 6x} \\ 2x^3 + 7x^2 - x - 5 \\ \underline{+ 10x^3 + 35x^2 - 5x - 75} \\ 12x^3 + 42x^2 - 6x - 80 \\ \underline{+ 10x^3 + 6x^2 - 34x + 12} \\ 22x^3 + 48x^2 - 40x - 68 \\ 22 \quad | \quad 22x^3 + 48x^2 - 40x - 68 \\ x^3 + x^2 - 3x - 3 \quad | \quad 5x^3 + 3x^2 - 17x + 6 \quad | \quad 5x - 2 \\ \underline{+ 5x^3 + 5x^2 - 15x - 6} \\ 2x^2 - 2x + 12 \\ \underline{+ 2x^2 + 2x - 6} \\ 4x + 6 \end{array}$$

H.C.F = $x^2 + x - 3$

3(iii) $2x^5 - 4x^4 - 6x$, $x^5 + x^4 - 3x^3 - 3x^2$
 Solution: $2x^5 - 4x^4 - 6x = x(2x^4 - 4x^3 - 6)$
 $x^5 + x^4 - 3x^3 - 3x^2 = x^2(x^3 + x^2 - 3x - 3)$

$$\begin{array}{r} 2x - 6 \\ x^3 + x^2 - 3x - 3 \quad | \quad 2x^4 - 4x^3 - 6 \\ \underline{+ 2x^4 + 2x^3 - 6x^2 + 6x} \\ - 6x^3 + 6x^2 + 6x - 6 \\ \underline{+ 6x^3 + 6x^2 + 18x + 18} \\ 12x^2 + 12x - 24 \\ 12 \quad | \quad 12x^2 + 12x - 24 \\ x^3 + x^2 - 3x - 3 \quad | \quad x + 2 \\ \underline{+ x^3 + x^2 - 2x - 3} \\ 2x^2 - x - 3 \\ \underline{+ 2x^2 + 2x + 4} \\ x + 1 \quad | \quad x^2 - x - 2 \quad | \quad 5x - 2 \\ \underline{+ x^2 + x} \end{array}$$

MATHEMATICS FOR 9TH CLASS (UNIT # 6)

Pilot Superone Mathematics 276 Class 9th

$$\begin{array}{r} -2x \quad 2 \\ -2x \quad -2 \\ \hline \times \end{array}$$

H.C.F. of x and x^2 is x

$$\text{H.C.F.} = x(x+1)$$

Q.4 Find the L.C.M. of the following expression by factorization.

(i) $39x^7y^3z$, $91x^5y^6z^7$

(ii) $102xy^2z$, $85x^2yz$, $187xyz^2$

4(i) $39x^7y^3z$, $91x^5y^6z^7$

Solution: $39x^7y^3z = 3 \times 13x^7y^3z = 3 \times 13x^5x^2y^3z$

$$91x^5y^6z^7 = 7 \times 13x^5y^6z^7 = 7 \times 13x^5y^3y^3z^6z$$

$$\begin{aligned} \text{L.C.M.} &= (\text{Common factor}) \times (\text{non-common factor}) \\ &= (13x^5y^3z)(3 \times 7x^2y^3z^6) \\ &= 13 \times 3 \times 7x^{5+2}y^{3+3}z^{1+6} \\ &= 273x^7y^6z^7 \end{aligned}$$

4(ii) $102xy^2z$, $85x^2yz$, $187xyz^2$

Solution: $102xy^2z = 2 \times 3 \times 17xyyz$

$$85x^2yz = 5 \times 17 \times x \times y \times z$$

$$187xyz^2 = 11 \times 17xyz^2$$

$$\begin{aligned} \text{L.C.M.} &= (\text{Common factor}) \times (\text{non-common factor}) \\ &= (17xyz)(2 \times 5 \times 11xyz) \\ &= 17 \times 2 \times 3 \times 5 \times 11x^2y^2z^2 \\ &= 5610x^2y^2z^2 \end{aligned}$$

Q.5 Find the L.C.M. of the following expression by factorization.

(i) $x^2 - 25x + 100$, $x^2 - x - 20$

(ii) $x^2 + 4x + 4$, $x^2 - 4$, $2x^2 + x - 6$

(iii) $2(x^4 - y^4)$, $3(x^3 + 2x^2 - xy^2 - 2y^3)$

(iv) $4(x^4 - 1)$, $6(x^3 - x^2 - x + 1)$

MATHEMATICS FOR 9TH CLASS (UNIT # 6)

Pilot Superone Mathematics 277 Class 9th

5(i) $x^2 - 25x + 100, x^2 - x - 20$

Solution: $x^2 - 25x + 100 = x^2 - 5x - 20x + 100$
 $= x(x - 5) - 20(x - 5)$
 $= (x - 5)(x - 20)$
 $x^2 - x - 20 = x^2 - 5x + 4x - 20$
 $= x(x - 5) + 4(x - 5)$
 $= (x - 5)(x + 4)$

Now $x^2 - 25x + 100 = (x - 5)(x - 20)$

and $x^2 - x - 20 = (x - 5)(x + 4)$

L.C.M. $= (x - 5)(x - 20)(x + 4)$

5(ii) $x^2 + 4x + 4, x^2 - 4, 2x^2 + x - 6$

Solution: $x^2 + 4x + 4 = x^2 + 2x + 2x + 4$
 $= x(x + 2) + 2(x + 2)$
 $= (x + 2)(x + 2)$

$x^2 - 4 = (x)^2 - (2)^2$
 $= (x - 2)(x + 2)$

$2x^2 + x - 6 = 2x^2 + 4x - 3x - 6$
 $= 2x(x + 2) - 3(x + 2)$
 $= (x + 2)(2x - 3)$

Now $x^2 + 4x + 4 = (x + 2)(x + 2)$

$x^2 - 4 = (x - 2)(x + 2)$

$2x^2 + x - 6 = (x + 2)(2x - 3)$

L.C.M. $= (x + 2)(x + 2)(x - 2)(2x - 3)$
 $= (x + 2)^2(x - 2)(2x - 3)$

5(iii) $2(x^4 - y^4), 3(x^3 + 2x^2 - xy^2 - 2y^3)$

Solution: $2(x^4 - y^4) = 2[(x^2)^2 - (y^2)^2]$
 $= 2(x^2 + y^2)(x^2 - y^2)$
 $= 2(x^2 + y^2)(x + y)(x - y)$
 $3(x^3 + 2x^2 - xy^2 - 2y^3) = 3[(x^3 + 2x^2y) - (xy^2 + 2y^3)]$
 $= 3[x^2(x + 2y) - y^2(x + 2y)]$

MATHEMATICS FOR 9TH CLASS (UNIT # 6)

Piot Superone Mathematics 278 Class 9th

$$\begin{aligned}
 &= 3(x+2y)(x^2-y^2) \\
 &= 3(x+2y)(x+y)(x-y) \\
 2(x^4-y^4) &= 2(x^2+y^2)(x+y)(x-y) \\
 3(x^3+2x^2-xy^2-2y^3) &= 3(x+2y)(x+y)(x-y) \\
 \text{L.C.M.} &= 2 \times 3(x+y)(x-y)(x+2y)(x^2+y^2) \\
 &= 6[(x+y)(x-y)](x+2y)(x^2+y^2) \\
 &= 6(x^2-y^2)(x^2+y^2)(x+2y) \\
 &= 6[(x^2-y^2)(x^2+y^2)](x+2y) \\
 &= 6(x^4-y^4)(x+2y)
 \end{aligned}$$

5(iv) $4(x^4-1), 6(x^3-x^2-x+1)$

Solution

$$\begin{aligned}
 4(x^4-1) &= 2 \times 2[(x^2)^2-(1)^2] \\
 &= 2 \times 2(x^2+1)(x^2-1) \\
 &= 2 \times 2(x^2+1)(x+1)(x-1) \\
 6(x^3-x^2-x+1) &= 2 \times 3[x^3-x^2-x+1] \\
 &= 2 \times 3[x^2(x-1)-1(x-1)] \\
 &= 2 \times 3(x-1)(x^2-1) \\
 4(x^4-1) &= 2 \times 2(x^2+1)(x+1)(x-1) \\
 6(x^3-x^2-x+1) &= 2 \times 3(x-1)(x+1)(x-1)(x^2+1) \\
 \text{L.C.M.} &= 2 \times 2 \times 3(x-1)(x+1)(x-1)(x^2+1) \\
 &= 12(x-1)(x+1)(x-1)(x^2+1) \\
 &= 12(x-1)(x^2-1)(x^2+1) \\
 &= 12(x-1)(x^2)^2+(1)^2 \\
 &= 12(x-1)(x^4-1)
 \end{aligned}$$

Q.6 For what value of k is $(x+4)$ the H.C.F. of $x^2+x-(2k+2)$ and $2x^2+kx-12$?

Solution: $(x+4)$ will divide completely $x^2+x-(2k+2)$ and $2x^2+kx-12$.

$$p(x) = x^2 + x - 2k - 2$$

put $x = -4$

$$p(-4) = (-4)^2 + (-4) - 2k - 2$$

MATHEMATICS FOR 9TH CLASS (UNIT # 6)

Pilot Superone Mathematics 279 Class 9th

$$16 - 4 - 2k - 2$$

$$10 - 2k = R$$

R must be zero

$$10 - 2k = 0$$

$$2k = 10$$

$$k = 5$$

$$q(x) = 2x^2 + kx - 12$$

Put $x = -4$

$$q(-4) = 2(-4)^2 - 4k - 12$$

$$= 32 - 4k - 12$$

$$= 20 - 4k = R$$

R must be zero

$$20 - 4k = 0$$

$$-4k = -20$$

$$k = 5$$

Q.7 If $(x + 3)(x - 2)$ is the H.C.F. of

$$p(x) = (x + 3)(2x^2 - 3x + k) \text{ and}$$

$$q(x) = (x - 2)(3x^2 + 7x - 1), \text{ find } k \text{ and } l.$$

Solution: $(x - 2)(x + 3)$ will divide $p(x) = (x + 3)(2x^2 - 3x + k)$

We put $x = 2$ in it.

$$p(2) = (2 + 3)[2 \times 2^2 - 3(2) + k]$$

$$= 5(8 - 6 + k)$$

$$= 5(2 + k) = R$$

R must be zero

$$5(2 + k) = 0$$

$$2 + k = 0$$

$$k = -2$$

$$q(x) = (x - 2)(3x^2 + 7x - 1)$$

$$(x - 2)(x + 3) \text{ will divide } q(x) = (x - 2)(3x^2 + 7x - 1)$$

Completely We put $x = -3$ in it

MATHEMATICS FOR 9TH CLASS (UNIT # 6)

Pilot Superone Mathematics 280 Class 9th

$$\begin{aligned} q(-3) &= (-3 - 2)[3(-3)^2 + 7(-3) - 1] \\ &= (-5)(27 - 21 - 1) \\ &= (-5)(6 - 1) = R \end{aligned}$$

R must be zero

$$(-5)(6 - 1) = 0$$

$$6 - 1 = 0$$

$$1 = 6$$

Q.8 The L.C.M. and H.C.F of two polynomials $p(x)$ and $2(x^4 - 1)$ and $(x + 1)(x^2 + 1)$ respectively.

If $p(x) = x^3 + x^2 + x + 1$, find $q(x)$ are

Solution: Formula $p(x) \times q(x) = \text{L.C.M.} \times \text{H.C.F}$

$$p(x) \times q(x) = 2(x^4 - 1)(x + 1)(x^2 + 1)$$

$$(x^3 + x^2 + x + 1) \times q(x) = 2(x^4 - 1)(x + 1)(x^2 + 1)$$

$$q(x) = \frac{2(x^4 - 1)(x + 1)(x^2 + 1)}{x^3 + x^2 + x + 1}$$

$$= \frac{2(x^4 - 1)(x + 1)(x^2 + 1)}{x^2(x + 1) + 1(x + 1)}$$

$$= \frac{2(x^4 - 1)(x + 1)(x^2 + 1)}{(x + 1)(x^2 + 1)}$$

$$qx = 2(x^4 - 1)$$

Q.9 Let $p(x) = 10(x^2 - 9)(x^2 - 3x + 2)$ and

$q(x) = 10x(x + 3)(x - 1)^2$. If the H.C.F. of $p(x)$, $q(x)$ is $10(x + 3)(x - 1)$, find their L.C.M.

Solution: Formula $p(x) \times q(x) = \text{L.C.M.} \times \text{H.C.F}$

Putting values in it.

$$10(x + 3)(x - 1) \times q(x) = 10(x^2 - 9)(x^2 - 3x + 2)(10)(x)(x + 3)(x - 1)^2$$

$$q(x) = \frac{10(x^2 - 9)(x^2 - 3x + 2)(10)(x)(x + 3)(x - 1)^2}{10(x + 3)(x - 1)}$$

MATHEMATICS FOR 9TH CLASS (UNIT # 6)

$$\begin{aligned}
 & \text{Pilot Superone Mathematics} \quad 281 \quad \text{Class 9th} \\
 & = (x^2 - 9)(x^2 - 3x + 2)(10)(x)(x - 1) \\
 & = 10x(x^2 - 9)(x - 1)(x^2 - 3x + 2) \\
 & = 10x(x^2 - 9)(x - 1)(x^2 - 2x - x + 2) \\
 & = 10x(x^2 - 9)(x - 1)x(x - 2) - 1(x - 2) \\
 & = 10x(x^2 - 9)(x - 1)(x - 2) \\
 & = 10x(x^2 - 9)(x - 1)^2(x - 2)
 \end{aligned}$$

Q.10 Let the product of L.C.M. and H.C.F. of two polynomials be $(x + 3)^2(x - 2)(x + 5)$. If one polynomial is $(x + 3)(x - 2)$ and the second polynomial is $x^2 + kx + 15$, find the value of k .

Solution. (L.C.M.) \times (H.C.F.) $= (x + 3)^2(x - 2)(x + 5)$

$$p(x) = (x + 3)(x - 2)$$

$$q(x) = x^2 + kx + 15$$

$$p(x) \times q(x) = \text{H.C.F.} \times \text{L.C.M.}$$

Putting value in it

$$(x^2 + kx + 15)(x + 3)(x - 2) = (x + 3)^2(x - 2)(x + 5)$$

$$x^2 + kx + 15 = \frac{(x + 3)^2(x - 2)(x + 5)}{(x + 3)(x - 2)}$$

$$x^2 + kx + 15 = (x + 3)(x + 5)$$

$$x^2 + kx + 15 = x^2 + 8x + 15$$

$$kx = x^2 + 8x + 15 - x^2 - 15$$

$$kx = 8x$$

$$\therefore k = 8$$

MATHEMATICS FOR 9TH CLASS (UNIT # 6)

Pilot Superone Mathematics 282 Class 9th

Q. 11 Waqas wishes to distribute 128 bananas and also 176 apples among a certain number of children. Find the highest number of children who can get the fruit in this way.

Solution: Number of children will be H.C.F. of 176 and 128

$$\begin{array}{r}
 \underline{1} \\
 128 \overline{) 176} \\
 \underline{128} 2 \\
 48 128 \\
 \underline{96} 1 \\
 32 \overline{) 48} \\
 \underline{32} 2 \\
 16 \overline{) 32} \\
 \underline{32} \\
 0
 \end{array}$$

Number of children = 16

MATHEMATICS FOR 9TH CLASS (UNIT # 6)

Pilot Superone Mathematics 284 Class 9th

$$Q.2 \quad \left[\frac{x+1}{x-1} - \frac{x-1}{x+1} - \frac{4x}{x^2+1} \right] + \frac{4x}{x^4-1}$$

$$\begin{aligned} \text{Solution:} &= \left(\frac{x+1}{x-1} - \frac{x-1}{x+1} \right) - \frac{4x}{x^2+1} + \frac{4x}{x^4-1} \\ &= \frac{(x+1)^2 - (x-1)^2}{(x-1)(x+1)} - \frac{4x}{x^2+1} + \frac{4x}{x^4-1} \\ &= \frac{x^2+2x+1-x^2+2x-1}{x^2-1} - \frac{4x}{x^2+1} + \frac{4x}{x^4-1} \\ &= \left[\frac{4x}{x^2-1} - \frac{4x}{x^2+1} \right] + \frac{4x}{x^4-1} \\ &= (4x) \left[\frac{1}{x^2-1} - \frac{1}{x^2+1} \right] + \frac{4x}{x^4-1} \\ &= (4x) \left[\frac{x^2+1-x^2+1}{(x^2-1)(x^2+1)} \right] + \frac{4x}{x^4-1} \\ &= (4x) \left[\frac{2}{x^4-1} \right] + \frac{4x}{x^4-1} \\ &\swarrow (4x) \left[\frac{2}{x^4-1} + \frac{1}{x^4-1} \right] \\ &= (4x) \left[\frac{2+1}{(x^4-1)} \right] \\ &= \frac{12x}{x^4-1} \end{aligned}$$

$$Q.3 \quad \frac{1}{x^2-8x+15} + \frac{1}{x^2-4x+3} - \frac{2}{x^2-6x+5}$$

Solution: =

$$\frac{1}{x^2-3x-5x+15} + \frac{1}{x^2-x-3x+3} - \frac{2}{x^2-x-5x+5}$$

MATHEMATICS FOR 9TH CLASS (UNIT # 6)

$$\begin{aligned}
 & \text{Pilot Superone Mathematics} \quad 285 \quad \text{Class 9th} \\
 & = \frac{1}{x(x-3)-5(x-3)} + \frac{1}{x(x-1)-3(x-1)} - \frac{2}{x(x-1)-5(x-1)} \\
 & = \frac{1}{(x-3)(x-5)} + \frac{1}{(x-1)(x-3)} - \frac{2}{(x-1)(x-5)} \\
 & = \frac{(x-1)+(x-5)-2(x-3)}{(x-3)(x-5)(x-1)} \\
 & = \frac{x-1+x-5-2x+6}{(x-3)(x-5)(x-1)} \\
 & = \frac{2x-2x+6-6}{(x-3)(x-5)(x-1)} \\
 & = 0
 \end{aligned}$$

$$\text{Q.4} \quad \frac{(x+2)(x+3)}{x^2-9} + \frac{(x+2)(2x^2-32)}{(x-4)(x^2-x-6)}$$

$$\begin{aligned}
 \text{Solution:} \quad & = \frac{(x+2)(x+3)}{(x)^2-(3)^2} + \frac{(x+2)(2)(x^2-16)}{(x-4)(x^2-3x+2x-6)} \\
 & = \frac{(x+2)(x+3)}{(x-3)(x+3)} + \frac{2(x+2)(x^2-4^2)}{(x-4)[x(x-3)+2(x-3)]} \\
 & = \frac{(x+2)}{(x-3)} + \frac{2(x+2)(x-4)(x+4)}{(x-4)(x-3)(x+2)} \\
 & = \frac{x+2}{x-3} + \frac{2(x+4)}{x-3} \\
 & = \frac{x+2+2x+8}{x-3} \\
 & = \frac{3x+10}{x-3}
 \end{aligned}$$

MATHEMATICS FOR 9TH CLASS (UNIT # 6)

Pilot Superone Mathematics 286 Class 9th

$$Q.5 \quad \frac{x+3}{2x^2+9x+9} + \frac{1}{2(2x-3)} - \frac{4x}{4x^2-9}$$

$$\begin{aligned} \text{Solution: } &= \frac{x+3}{2x^2+6x+3x+9} + \frac{1}{2(2x-3)} - \frac{4x}{(2x)^2-(3)^2} \\ &= \frac{x+3}{2x(x+3)+3(x+3)} + \frac{1}{2(2x-3)} - \frac{4x}{(2x+3)(2x-3)} \\ &= \frac{x+3}{(2x+3)(x+3)} + \frac{1}{2(2x-3)} - \frac{4x}{(2x+3)(2x-3)} \\ &= \frac{1}{(2x+3)} + \frac{1}{2(2x-3)} - \frac{4x}{(2x+3)(2x-3)} \\ &= \frac{2(2x-3) + (2x+3) - (4x)(2)}{2(2x+3)(2x-3)} \\ &= \frac{4x-6+2x+3-8x}{2(2x+3)(2x-3)} \\ &= \frac{2x-3}{2(2x+3)(2x-3)} \\ &= \frac{-(2x+3)}{2(2x+3)(2x-3)} \\ &= \frac{-1}{2(2x-3)} \end{aligned}$$

$$Q.6 \quad A - \frac{1}{A} \text{ if } A = \frac{a+1}{a-1}$$

$$\begin{aligned} \text{Solution: } \quad A - \frac{1}{A} &= \frac{a+1}{a-1} - \frac{1}{\frac{a+1}{a-1}} \\ &= \frac{a+1}{a-1} - \frac{a-1}{a+1} \end{aligned}$$

MATHEMATICS FOR 9TH CLASS (UNIT # 6)

Pilot Superone Mathematics 287

Class 9th

$$\begin{aligned} & \frac{(a+1)^2 - (a-1)^2}{(a-1)(a+1)} \\ & \frac{(a^2 + 2a + 1) - (a^2 - 2a + 1)}{(a-1)(a+1)} \\ & = \frac{a^2 + 2a + 1 - a^2 + 2a - 1}{(a-1)(a+1)} \\ & = \frac{4a}{a^2 - 1} \end{aligned}$$

Q7 $\left[\frac{x-1}{x-2} + \frac{2}{2-x} \right] - \left[\frac{x+1}{x+2} + \frac{4}{4-x^2} \right]$

Solution: $= \left[\frac{x-1}{x-2} - \frac{2}{x-2} \right] - \left[\frac{x+1}{x+2} - \frac{4}{x^2-4} \right]$ N.T.S

$$\begin{aligned} & = \left[\frac{x-1-2}{x-2} \right] - \left[\frac{x+1}{x+2} - \frac{4}{(x+2)(x-2)} \right] \\ & = \left[\frac{x-3}{x-2} \right] - \left[\frac{(x+1)(x-2)(4)}{(x+2)(x-2)} \right] \\ & = \frac{x-3}{x-2} - \frac{x^2 - x - 6}{(x+2)(x-2)} \\ & = \frac{x-3}{x-2} - \frac{x^2 - 3x + 2x - 6}{(x+2)(x-2)} \\ & = \frac{x-3}{x-2} - \frac{x(x-3) + 2(x-3)}{(x+2)(x-2)} \\ & = \frac{x-3}{x-2} - \frac{(x-3)(x+2)}{(x+2)(x-2)} \end{aligned}$$

MATHEMATICS FOR 9TH CLASS (UNIT # 6)

Pilot Superone Mathematics 288 Class 9th

$$= \frac{x-3}{x-2} - \frac{x-3}{x-2}$$

$$= 0$$

Q.8 What rational expression should be subtracted from

$$\frac{2x^2 + 2x - 7}{x^2 + x - 6} \text{ to get } \frac{x-1}{x-2} ?$$

Let A be the required expression.

$$\frac{2x^2 + 2x - 7}{x^2 + x - 6} - A = \frac{x-1}{x-2}$$

$$\begin{aligned} \text{Solution: Thus } A &= \frac{2x^2 + 2x - 7}{x^2 + x - 6} - \frac{x-1}{x-2} \\ &= \frac{2x^2 + 2x - 7}{x^2 + 3x - 2x - 6} - \frac{x-1}{x-2} \\ &= \frac{2x^2 + 2x - 7}{x(x+3) - 2(x+3)} - \frac{x-1}{x-2} \\ &= \frac{2x^2 + 2x - 7}{(x+3)(x-2)} - \frac{x-1}{x-2} \\ &= \frac{2x^2 + 2x - 7 - (x-1)(x+3)}{(x+3)(x-2)} \\ &= \frac{2x^2 + 2x - 7 - (x^2 + 2x - 3)}{(x+3)(x-2)} \\ &= \frac{2x^2 + 2x - 7 - x^2 - 2x + 3}{(x+3)(x-2)} \\ &= \frac{x^2 - 4}{(x+3)(x-2)} \\ &= \frac{(x)^2 - (2)^2}{(x+3)(x-2)} \end{aligned}$$

MATHEMATICS FOR 9TH CLASS (UNIT # 6)

Pilot Superone Mathematics 289 Class 9th

$$= \frac{(x+2)(x-2)}{(x+3)(x-2)}$$

$$= \frac{x+2}{x+3}$$

Perform the indicated operations and simplify to the lowest

forms. $\frac{x^2+x-6}{x^2-x-6} \times \frac{x^2-4}{x^2-9}$

Q.9 Simplify to the lowest form

$$\frac{x^2+x-6}{x^2-x-6} \times \frac{x^2-2^2}{x^2-3^2}$$

Solution.

$$= \frac{x^2+3x-2x-6}{x^2-3x+2x-6} \times \frac{x^2-2^2}{x^2-3^2}$$

$$= \frac{x(x+3)-2(x+3)}{x(x-3)+2(x-3)} \times \frac{(x+2)(x-2)}{(x+3)(x-3)}$$

$$= \frac{(x+3)(x-2)}{(x-3)(x+2)} \times \frac{(x+2)(x-2)}{(x+3)(x-3)}$$

$$= \frac{(x-2)(x-2)}{(x-3)(x-3)}$$

$$= \frac{(x-2)^2}{(x-3)^2}$$

Q.10 $\frac{x^3-8}{x^2-4} + \frac{x^2+6x+8}{x^2-2x+1}$

$$\frac{x^3-2^3}{x^2-2^2} \times \frac{x^2+4x+2x+8}{x^2-x-x+1}$$

Solution:

$$= \frac{x^3-2^3}{x^2-2^2} + \frac{x(x+4)+2(x+4)}{x(x-1)-1(x-1)}$$

$$= \frac{(x-2)(x^2+2x+4)}{(x-2)(x+2)} + \frac{x(x+4)+2(x+4)}{x(x-1)-1(x-1)}$$

MATHEMATICS FOR 9TH CLASS (UNIT # 6)

Pilot Superone Mathematics 290 Class 9th

$$= \frac{x^2 + 2x + 4}{x + 2} + \frac{(x + 4)(x + 2)}{(x - 1)(x - 1)}$$

$$= \frac{(x^2 + 2x + 4)(x + 4)}{(x - 1)(x - 1)}$$

$$= \frac{(x + 4)(x^2 + 2x + 4)}{(x - 1)^2}$$

Q.11. $\frac{x^3 - 8x}{2x^2 + 5x - 3} \times \frac{2x - 1}{x^2 + 2x + 4} \times \frac{x + 3}{x^2 - 2x}$

Solution:

$$= \frac{x(x^2 - 8)}{2x^2 + 6x - 3} \times \frac{2x - 1}{x^2 + 2x + 4} \times \frac{x + 3}{x(x - 2)}$$

$$= \frac{x(x^2 - 2^3)}{2x(x + 3) - 1(x + 3)} \times \frac{2x - 1}{x^2 + 2x + 4} \times \frac{x + 3}{x(x - 2)}$$

$$= \frac{x(x - 2)(x^2 + 2x + 4)}{(x + 3)(2x - 1)} \times \frac{2x - 1}{x^2 + 2x + 4} \times \frac{x + 3}{x(x - 2)}$$

$$= \frac{x(x - 2)(x^2 + 2x + 4)(2x - 1)(x + 3)}{x(x - 2)(x^2 + 2x + 4)(2x - 1)(x + 3)}$$

$$= 1$$

Q.12 $\frac{2y^2 + 7y - 4}{3y^2 - 13y + 4} + \frac{2y^2 - 1}{6y^2 + y - 1}$

$$\frac{2y^2 + 7y - 4}{3y^2 - 13y + 4} + \frac{4y^2 - 1}{6y^2 + y - 1}$$

Solution:

$$= \frac{2y^2 + 8y - y - 4}{3y^2 - 12y - y + 4} + \frac{(2y)^2 - (1)^2}{6y^2 + 3y - 2y - 1}$$

$$= \frac{2y(y + 4) - 1(y + 4)}{3y(y - 4) - 1(y - 4)} + \frac{(2y - 1)(2y + 1)}{3y(2y + 1) - 1(2y + 1)}$$

$$= \frac{(y + 4)(2y - 1)}{(y - 4)(3y - 1)} + \frac{(2y - 1)(2y + 1)}{(2y + 1)(3y - 1)}$$

MATHEMATICS FOR 9TH CLASS (UNIT # 6)

Pilot Superone Mathematics 291 Class 9th

$$= \frac{(y+4)(2y-1)}{(y-4)(3y-1)} \times \frac{2y-1}{3y-1}$$

$$= \frac{(y+4)(2y-1)}{(y-4)(3y-1)} \times \frac{3y-1}{2y-1}$$

$$= \frac{y+4}{y-4}$$

Q.13 $\left[\frac{x^2+y^2}{x^2-y^2} - \frac{x^2-y^2}{x^2+y^2} \right] \div \left[\frac{x+y}{x-y} - \frac{x-y}{x+y} \right]$

Solution. = $\left[\frac{(x^2+y^2)^2 - (x^2-y^2)^2}{(x^2-y^2)(x^2+y^2)} \right] \div \left[\frac{(x+y)^2 - (x-y)^2}{(x-y)(x+y)} \right]$

$$\frac{(x^4+y^4+2x^2y^2) - (x^4+y^4-2x^2y^2)}{(x^2-y^2)(x^2+y^2)} \div \frac{x^2+y^2+2xy - (x^2+y^2-2xy)}{(x-y)(x+y)}$$

$$\frac{x^4+y^4+2x^2y^2-x^4-y^4-2x^2y^2}{(x^2-y^2)(x^2+y^2)} \div \frac{x^2+y^2+2xy-x^2-y^2+2xy}{x^2-y^2}$$

$$= \frac{4x^2y^2}{(x^2-y^2)(x^2+y^2)} \div \frac{4xy}{x^2-y^2}$$

$$= \frac{4x^2y^2}{(x^2-y^2)(x^2+y^2)} \times \frac{x^2-y^2}{4xy}$$

$$= \frac{xy}{x^2+y^2}$$

Exercise 6.3

Q.1 Use factorization to find the square root of the following expressions.

(i) $4x^2 - 12xy + 9y^2$

(ii) $x^2 - 1 + \frac{1}{4x^2}, (x \neq 0)$

(iii) $\frac{1}{16}x^2 - \frac{1}{12}xy + \frac{1}{36}y^2$

(iv) $4(a+b)^2 - 12(a^2 - b^2) + 9(a-b)^2$

(v) $\frac{4x^6 - 12x^3y^3 + 9y^6}{9x^4 + 24x^2y^2 + 16y^4}, (x \neq 0)$

(vi) $\left(x + \frac{1}{x}\right)^2 - 4\left(x - \frac{1}{x}\right), (x \neq 0)$

(vii) $\left(x^2 + \frac{1}{x^2}\right)^2 - 4\left(x + \frac{1}{x}\right)^2 + 12, (x \neq 0)$

(viii) $(x^2 + 3x + 2)(x^2 + 4x + 3)(x^2 + 5x + 6)$

(ix) $(x^2 + 8x + 7)(2x^2 - x - 3)(2x^2 + 11x - 21)$

1(i) $4x^2 - 12xy + 9y^2$

Solution: $4x^2 - 12xy + 9y^2 = 4x^2 - 6xy - 6xy + 9y^2$
 $= 2x(2x - 3y) - 3y(2x - 3y)$
 $= (2x - 3y)(2x - 3y)$
 $= (2x - 3y)^2$

$\sqrt{4x^2 - 12xy + 9y^2} = \pm(2x - 3y)$

1(ii) $x^2 - 1 + \frac{1}{4x^2}, (x \neq 0)$

Solution: $x^2 - 1 + \frac{1}{4x^2} = (x)^2 - 1 + \left(\frac{1}{2x}\right)^2$

MATHEMATICS FOR 9TH CLASS (UNIT # 6)

Pilot Superone Mathematics 293 Class 9th

$$= (x)^2 - 2(x)\left(\frac{1}{2x}\right) + \left(\frac{1}{2x}\right)^2$$

$$= \left(x - \frac{1}{2x}\right)^2$$

$$\sqrt{x^2 - 1 + \frac{1}{4x^2}} = \pm \left(x - \frac{1}{2x}\right)$$

1(iii) $\frac{1}{16}x^2 - \frac{1}{12}xy + \frac{1}{36y^2}$

Solution: $\frac{1}{16}x^2 - \frac{1}{12}xy + \frac{1}{36y^2} = \left(\frac{1}{4}x\right)^2 - \frac{1}{12}xy + \left(\frac{1}{6y}\right)^2$

$$= \left(\frac{1}{4}x\right)^2 - 2\left(\frac{1}{4}x\right)\left(\frac{1}{6y}\right) + \left(\frac{1}{6y}\right)^2$$

$$= \left(\frac{1}{4}x - \frac{1}{6y}\right)^2$$

$$\sqrt{\frac{1}{16}x^2 - \frac{1}{12}xy + \frac{1}{36y^2}} = \pm \left(\frac{x}{4} - \frac{1}{6y}\right)$$

1(iv) $4(a+b)^2 - 12(a^2 - b^2) + 9(a-b)^2$

Solution: $= [2(a+b)]^2 - 12(a+b)(a-b) + [3(a-b)]^2$

$$= [2(a+b)]^2 - (2)(2)(a+b)(3)(a-b) + [3(a-b)]^2$$

$$= [2(a+b) - 3(a-b)]^2$$

$$= (2a + 2b - 3a + 3b)^2$$

$$= (5b - a)^2$$

$$\sqrt{4(a+b)^2 - 12(a^2 - b^2) + 9(a-b)^2} = \pm(5b - a)$$

1(v) $\frac{4x^4 - 12x^3y^3 + 9y^6}{9x^4 + 24x^3y^3 + 16y^6}, (x \neq 0)$

MATHEMATICS FOR 9TH CLASS (UNIT # 6)

Pilot Superone Mathematics 294 Class 9th

$$\begin{aligned}\text{Solution.} \quad &= \frac{(2x^1)^2 - 2(2x^1)(3y^1) + (3y^1)^2}{(3x^1)^2 + 2(3x^1)(4y^1) + (4y^1)^2} \\ &= \frac{(2x^1 - 3y^1)^2}{(3x^1 + 4y^1)^2}\end{aligned}$$

$$\frac{4x^6 - 12x^1y^3 + 9y^6}{9x^4 + 24x^1y^2 + 16y^4} = \left(\frac{2x^1 - 3y^1}{3x^1 + 4y^1} \right)^2$$

$$\sqrt{\frac{4x^6 - 12x^1y^3 + 9y^6}{9x^4 + 24x^1y^2 + 16y^4}} = \pm \frac{2x^1 - 3y^1}{3x^1 + 4y^1}$$

$$I(vf) \quad \left(x + \frac{1}{x} \right)^2 - 4 \left(x - \frac{1}{x} \right), (x \neq 0)$$

$$\text{Solution: Let } x + \frac{1}{x} = a$$

$$\begin{aligned}\text{Now } \left(x + \frac{1}{x} \right)^2 &= \left(x - \frac{1}{x} \right)^2 + 4 \\ &= a^2 + 4\end{aligned}$$

$$= (a^2 + 4) - 4a \quad \text{[Given expression]}$$

$$= a^2 - 4a + 4$$

$$= (a - 2)^2$$

$$= \left[\left(x + \frac{1}{x} \right) - 2 \right]^2$$

$$\left(x + \frac{1}{x} \right)^2 - 4 \left(x - \frac{1}{x} \right) = \left[\left(x - \frac{1}{x} \right) - 2 \right]^2$$

$$\sqrt{\left(x + \frac{1}{x} \right)^2 - 4 \left(x - \frac{1}{x} \right)} = \pm \left[\left(x - \frac{1}{x} \right) - 2 \right]$$

MATHEMATICS FOR 9TH CLASS (UNIT # 6)

Pilot Superone Mathematics 295

Class 9th

$$I(vi) \left(x^2 + \frac{1}{x^2} \right)^2 - 4 \left(x + \frac{1}{x} \right)^2 + 12$$

Solution: Let $x + \frac{1}{x} = a$ (A)

$$\left(x + \frac{1}{x} \right)^2 = a^2 \quad (\text{Squaring})$$

$$x^2 + \frac{1}{x^2} + 2 = a^2$$

$$x^2 + \frac{1}{x^2} = a^2 - 2 \quad (B)$$

$$\left(x^2 + \frac{1}{x^2} \right)^2 - 4 \left(x + \frac{1}{x} \right)^2 + 12 = (a^2 - 2)^2 - 4a^2 + 12 \quad (\text{From A, B})$$

$$= a^4 - 4a^2 + 4 - 4a^2 + 12$$

$$= a^4 - 8a^2 + 16$$

$$= (a^2)^2 - 2(a^2)(4) + (4)^2$$

$$= (a^2 - 4)^2$$

$$\sqrt{\left(x^2 + \frac{1}{x^2} \right)^2 - 4 \left(x + \frac{1}{x} \right)^2 + 12} = \sqrt{(a^2 - 4)^2} \quad (\text{Taking square root})$$

$$= \pm (a^2 - 4)$$

$$= \pm \left[\left(x + \frac{1}{x} \right)^2 - 4 \right] \quad (\text{Putting value of } a)$$

a)

$$= + \left[x^2 + \frac{1}{x^2} + 2 - 4 \right] \quad \pm \left[x^2 + \frac{1}{x^2} - 2 \right]$$

$$I(viii) (x^2 + 3x + 2)(x^2 + 4x + 3)(x^2 + 5x + 6)$$

Solution: $= [x^2 + 2x + x + 2][x^2 + x + 3x + 3][x^2 + 2x + 3x + 6]$
 $= [x(x+2) + 1(x+2)][x(x+1) + 2(x+1)][x(x+2) + 3(x+2)]$

MATHEMATICS FOR 9TH CLASS (UNIT # 6)

Pilot Superone Mathematics 296 Class 9th

$$= (x + 2)(x + 1)(x + 1)(x + 3)(x + 2)(x + 3)$$

$$= (x + 1)^2 (x + 2)^2 (x + 3)^2$$

Taking square root

$$\sqrt{(x + 1)^2 (x + 2)^2 (x + 3)^2} = (x + 1)(x + 2)(x + 3)$$

1(ix) $(x^2 + 8x + 7)(2x^2 - x - 3)(2x^2 + 11x - 21)$

Solution: $= \{x^2 + x + 7x + 7\}[2x^2 - 3x + 2x - 3][2x^2 + 14x - 3x - 21]$

$$= \{x(x + 1) + 7(x + 1)\}[x(2x - 3) + 1(2x - 3)][2x(x + 7) -$$

$$(x + 7)]$$

$$= (x + 1)(x + 7)(2x - 3)(x + 1)(x + 7)(2x - 3)$$

$$= (x + 1)^2 (x + 7)^2 (2x - 3)^2$$

Taking square root

$$\sqrt{(x + 1)^2 (x + 7)^2 (2x - 3)^2} = (x + 1)(x + 7)(2x - 3)$$

Q.2 Use division method to find the square root of the following expression.

(i) $4x^2 + 12xy + 9y^2 + 16x + 24y + 16$

(ii) $x^4 - 10x^3 + 37x^2 - 60x + 36$

(iii) $9x^4 - 6x^3 + 7x^2 - 2x + 1$

(iv) $4 + 25x^2 - 12x - 24x^3 + 16x^4$

(v) $\frac{x^2}{y^2} - 10\frac{x}{y} + 27 - \frac{10y}{x} + \frac{y^2}{x^2}, (x \neq 0, y \neq 0)$

2(i) $4x^2 + 12xy + 9y^2 + 16x + 24y + 16$

Solution.

	$2x + 3y + 4$
$2x$	$4x^2 + 12xy + 9y^2 + 16x + 24y + 16$
	$\underline{-4x^2}$
$4x + 3y$	$12xy + 9y^2$
	$\underline{-12xy - 9y^2}$
$4x + 6y + 4$	$16x + 24y + 16$
	$\underline{-16x - 24y - 16}$
	0

Square root = $2x + 3y + 4$

MATHEMATICS FOR 9TH CLASS (UNIT # 6)

Pilot Superone Mathematics 297

Class 9th

2(ii) $x^4 - 10x^3 + 37x^2 - 60x + 36$

Solution.

$$\begin{array}{r} x^2 - 5x + 6 \\ x^2 \overline{) x^4 - 10x^3 + 37x^2 - 60x + 36} \\ \underline{\pm 4x^4} \\ 10x^3 + 37x^2 \\ \underline{- 10x^3 \pm 25x^2} \\ 12x^2 - 60x + 36 \\ \underline{\pm 12x^2 - 60x \pm 36} \\ 0 \end{array}$$

Square root = $x^2 - 5x + 6$

2(iii) $9x^4 - 6x^3 + 7x^2 - 2x + 1$

Solution:

$$\begin{array}{r} 3x^2 - x + 1 \\ 3x^2 \overline{) 9x^4 - 6x^3 + 7x^2 - 2x + 1} \\ \underline{\pm 9x^4} \\ 6x^3 + 7x^2 \\ \underline{6x^3 \pm x^2} \\ 6x^2 - 2x + 1 \\ \underline{\pm 6x^2 - 2x \pm 1} \\ 0 \end{array}$$

Square root = $3x^2 - x + 1$

2(iv) $4 + 25x^2 - 12x - 24x^3 + 16x^4$

Solution.

$$\begin{array}{r} 4x^2 - 3x + 2 \\ 4x^2 \overline{) 16x^4 - 24x^3 + 25x^2 - 12x + 4} \\ \underline{\pm 16x^4} \\ 24x^3 + 25x^2 \\ \underline{- 24x^3 \pm 9x^2} \\ 16x^2 - 12x + 4 \\ \underline{\pm 16x^2 - 12x \pm 4} \\ 0 \end{array}$$

Square root = $4x^2 - 3x + 2$

2(v) $\frac{x^4}{y^2} - 10\frac{x}{y} + 27 - \frac{10y}{x} + \frac{y^2}{x^2}, (x \neq 0, y \neq 0)$

Solution: Writing in descending order

MATHEMATICS FOR 9TH CLASS (UNIT # 6)

Pilot Superone Mathematics 298 Class 9th

$$\begin{array}{r}
 \frac{x}{y} - 5 + \frac{y}{x} \\
 \hline
 \frac{x}{y} \quad \frac{x^2}{y^2} - \frac{10x}{y} + 27 \cdot \frac{10y}{x} + \frac{y^2}{x^2} \\
 \frac{x^2}{y^2} \\
 \hline
 \frac{2x}{y} - 5 \quad \frac{10x}{y} + 27 \\
 \frac{10x}{y} + 25 \\
 + \quad y \\
 \hline
 \frac{2x}{y} - 10 + \frac{y}{x} \quad 2 \cdot \frac{10y}{x} + \frac{y^2}{x^2} \\
 \quad \quad \quad 2 \cdot \frac{10x}{x} + \frac{y^2}{x^2} \\
 \quad \quad \quad - \quad + \quad - \\
 \quad \quad \quad \quad x
 \end{array}$$

Square root $\frac{x}{y} - 5 + \frac{y}{x}$

Q.3 Find the value of k for which the following expressions will become a perfect square.

(i) $4x^4 - 12x^3 + 37x^2 - 42x + k$

(ii) $x^4 - 4x^3 + 10x^2 - kx + 9$

3(i) $4x^4 - 12x^3 + 37x^2 - 42x + k$

Solut on: Taking square root

$$\begin{array}{r}
 2x^2 \quad 3x + 7 \\
 \hline
 4x^4 \quad 12x^3 + 37x^2 \quad 42x + k \\
 \pm 4x^4 \\
 \hline
 \quad - 12x^3 + 37x^2 \\
 \quad \quad 12x^2 \pm 9x^2 \\
 \hline
 4x^2 \quad 6x + 7 \quad \quad 28x^2 \quad 42x + k \\
 \quad \quad \quad \pm 28x^2 + 42x + 49 \\
 \quad \quad \quad \quad k - 49
 \end{array}$$

In case of perfect square, remainder is zero

Therefore $k - 49 = 0$

MATHEMATICS FOR 9TH CLASS (UNIT # 6)

Pilot Superone Mathematics 299 Class 9th
 and $k = 49$

3(ii) $x^4 - 4x^3 + 10x^2 - kx + 9$

Solution: Taking square root

$$\begin{array}{r}
 x^2 \quad 2x + 3 \\
 \hline
 x^4 - 4x^3 + 10x^2 - kx + 9 \\
 \pm x^4 \\
 \hline
 4x^3 + 10x^2 \\
 \pm 4x^3 \pm 4x^2 \\
 \hline
 6x^2 - kx + 9 \\
 \pm 6x^2 \pm 12x \pm 9 \\
 \hline
 kx + 12x
 \end{array}$$

In case of perfect square

$$-kx + 12x = 0$$

$$x(k - 12) = 0$$

$$k - 12 = 0$$

$$k = 12$$

Q.4 Find the value of l and m for which the following expressions will become perfect square.

(i) $x^4 + 4x^3 + 16x^2 + lx + m$

(ii) $49x^4 - 70x^3 + 109x^2 - lx - m$

4(i) $x^4 + 4x^3 + 16x^2 - lx + m$

Solution: Taking square root

$$\begin{array}{r}
 x^2 + 2x + 6 \\
 \hline
 x^4 + 4x^3 + 16x^2 + lx + m \\
 \pm x^4 \\
 \hline
 4x^3 + 16x^2 \\
 \pm 4x^3 \pm 4x^2 \\
 \hline
 12x^2 + lx + m \\
 \pm 12x^2 \pm 24x \pm 36 \\
 \hline
 (lx - 24x) + (m - 36)
 \end{array}$$

In case of perfect square, remainder is zero

MATHEMATICS FOR 9TH CLASS (UNIT # 6)

Pilot Superone Mathematics 300 Class 9th

$$1x - 24x = 0$$

$$x(1 - 24) = 0$$

$$1 - 24 = 0 \quad x = 0$$

$$1 = 24$$

and $m - 36 = 0$

$$m = 36$$

4(ii) $49x^4 - 70x^3 + 109x^2 - 1x - m$

Solution: Taking square root

$$\begin{array}{r}
 7x^2 \quad \begin{array}{l} x^2 + 2x + 6 \\ 49x^4 - 70x^3 + 109x^2 - 1x - m \\ \pm 49x^4 \\ \hline -70x^3 + 109x^2 \\ 70x^3 \pm 25x^2 \\ \hline 84x^2 + 1x - m \\ \pm 84x^2 - 60x + 36 \\ \hline (1x - 60x) + (-m - 36) \end{array} \\
 14x^2 - 5x \\
 14x^2 - 10x + 6
 \end{array}$$

In case of perfect square, remainder is zero

$$1x + 60x = 0$$

$$x(1 + 60) = 0$$

$$1 + 60 = 0 \quad x = 0$$

$$1 = 60$$

and $-m - 36 = 0$

$$m = 36$$

$$m = 36$$

Q.5 To make the expression $9x^4 - 12x^3 + 22x^2 - 13x + 12$, a perfect square

(i) What should be added to it?

(ii) What should be subtracted from it?

(iii) What should be the value of x?

Solution: Taking square root

$$3x^2 - 2x + 3$$

MATHEMATICS FOR 9TH CLASS (UNIT # 6)

Pilot Superone Mathematics 301 3x ² 14x ² - 5x 14x ² 10x+6	<table style="width: 100%; border-collapse: collapse;"> <tr> <td style="border-bottom: 1px solid black; padding: 5px;"> $9x^4 - 12x^3 + 22x^2 - 13x + 12$ $\pm 9x^4$ </td> </tr> <tr> <td style="border-bottom: 1px solid black; padding: 5px;"> $12x^3 + 22x^2$ $\pm 12x^3 \pm 4x^2$ </td> </tr> <tr> <td style="border-bottom: 1px solid black; padding: 5px;"> $18x^2 - 13x + 12$ $\pm 18x^2 - 12x \pm 9$ </td> </tr> <tr> <td style="padding: 5px;"> $x + 3 \text{ (R)}$ </td> </tr> </table>	$9x^4 - 12x^3 + 22x^2 - 13x + 12$ $\pm 9x^4$	$12x^3 + 22x^2$ $\pm 12x^3 \pm 4x^2$	$18x^2 - 13x + 12$ $\pm 18x^2 - 12x \pm 9$	$x + 3 \text{ (R)}$	Class 9 th
$9x^4 - 12x^3 + 22x^2 - 13x + 12$ $\pm 9x^4$						
$12x^3 + 22x^2$ $\pm 12x^3 \pm 4x^2$						
$18x^2 - 13x + 12$ $\pm 18x^2 - 12x \pm 9$						
$x + 3 \text{ (R)}$						

- (i) What should be subtracted, remainder should be subtracted i.e. $x + 3$
- (ii) What should be added, Negative of remainder should be added i.e. $x - 3$
- (iii) To find x, put remainder equal to zero.
 $x - 3 = 0$
 $x = 3$

**Unit
7**

**LINEAR EQUATIONS
AND INEQUALITIES**

Linear Equations

A linear equation is one unknown variable x is an equation of the form $ax + b = 0$, where $a, b \in \mathbb{R}$ and $a \neq 0$

Radical Equations

When the variable in an equation occurs under a radical the equation is called a radical equation

Exercise 7.1

Q.1 Solve the following equations.

(i) $\frac{2}{3}x - \frac{1}{2}x = x + \frac{1}{6}$ (ii) $x - \frac{3}{3} = \frac{x-2}{2} - 1$

(iii) $\frac{1}{2}\left(x - \frac{1}{6}\right) + \frac{2}{3} = \frac{5}{6} + \frac{1}{3}\left(\frac{1}{2} - 3x\right)$

(iv) $x + \frac{1}{3} = 2\left(x - \frac{2}{3}\right) - 6x$

(v) $\frac{5(x-3)}{6} = x = 1 - \frac{x}{9}$

(vi) $\frac{x}{3x-6} = 2 + \frac{2x}{x-2}, x \neq 2$

(vii) $\frac{2x-2}{2x+5} = \frac{5}{3} - \frac{5}{4x+10}, x \neq \frac{5}{2}$

(viii) $\frac{2x}{x-1} + \frac{1}{3} = \frac{5}{6} + \frac{2}{x-1}, x \neq 1$

(ix) $\frac{2}{x^2-1} - \frac{1}{x+1} = \frac{1}{x+1}, x \neq \pm 1$

MATHEMATICS FOR 9TH CLASS (UNIT # 7)

Pilot Superone Mathematics

311

Class 9th

$$(x) \quad \frac{2}{3x+6} - \frac{1}{6} = \frac{1}{2x+4} \quad x \neq -2$$

$$I(i) \quad \frac{2}{3}x - \frac{1}{2} = x + \frac{1}{6}$$

L.C.M. of 2, 3, 6 is 6 multiplying by 6

$$Solution: \quad (6) \frac{2}{3}x - (6) \frac{1}{2} = (6)x + (6) \frac{1}{6}$$

$$4x - 3x = 6x + 1$$

$$x = 6x + 1$$

$$x - 6x = 1$$

$$-5x = 1$$

$$x = -\frac{1}{5}$$

Solve

$$x = \left\{ -\frac{1}{5} \right\}$$

$$I(ii) \quad \frac{x-3}{3} - \frac{x-2}{2} = 1$$

$$Solution: \quad \left(\frac{x-3}{3} \right) \times 6 - \left(\frac{x-2}{2} \right) \times 6 = (1) \times 6$$

L.C.M. of 2, 3, 6 multiplying by 6

$$(x-3) \times 2 - (x-2) \times 3 = 6$$

$$2x - 6 - 3x + 6 = 6$$

$$2x - 3x = 6$$

$$-x = 6$$

$$x = 6$$

Solve

$$x = \{6\}$$

$$I(iii) \quad \frac{1}{2} \left(x - \frac{1}{6} \right) + \frac{2}{3} = \frac{5}{6} + \frac{1}{3} \left(\frac{1}{2} - 3x \right)$$

Solution:

$$\frac{x}{2} - \frac{1}{12} + \frac{2}{3} = \frac{5}{6} + \frac{1}{6} - x$$

L.C.M. of 2, 12, 3, 6 is 12 multiplying by 12

MATHEMATICS FOR 9TH CLASS (UNIT # 7)

Pilot Superone Mathematics 312 Class 9th

$$12\left(\frac{x}{7}\right) - 12\left(\frac{1}{12}\right) + 12\left(\frac{2}{3}\right) = 12\left(\frac{5}{6}\right) + 12\left(\frac{1}{6}\right) - 12(x)$$

$$6x - 1 + 8 - 10 + 2 = 12x$$

$$6x + 12x = 10 + 2 + 1 - 8$$

$$18x = 5$$

$$x = \frac{5}{18}$$

Solve

$$x = \left\{ \frac{5}{18} \right\}$$

$$I(iv) \quad x + \frac{1}{3} = 2\left(x - \frac{2}{3}\right) - 6x$$

$$\text{Solution:} \quad x + \frac{1}{3} = 2x - \frac{4}{3} - 6x$$

Multiplying by 3

$$3x + 3\left(\frac{1}{3}\right) = 3(2x) - 3\left(\frac{4}{3}\right) - 3(6x)$$

$$3x + 1 = 6x - 4 - 18x$$

$$3x - 6x + 18x = 4 - 1$$

$$15x = 5$$

$$x = \frac{5}{15}$$

$$x = -\frac{1}{3}$$

Solve set

$$x = \left\{ -\frac{1}{3} \right\}$$

$$I(v) \quad \frac{5(x-3)}{6} - x = 1 - \frac{x}{9}$$

Solution: L.C.M. of 6, 9, is 18 multiplying by 18

MATHEMATICS FOR 9TH CLASS (UNIT # 7)

Pilot Superone Mathematics 313 Class 9th

$$\frac{18(5)(x-3)}{6} - 18x = 18 - 18\left(\frac{x}{9}\right)$$

$$15(x-3) - 18x = 18 - 2x$$

$$15x - 45 - 18x = 18 - 2x$$

$$15x - 18x + 2x = 18 + 45$$

$$-x = 63$$

$$x = -63$$

$$\text{Solve set } x = \{-63\}$$

$$I(vi) \quad \frac{x}{3x-6} = 2 - \frac{2x}{x-2}, x \neq 2$$

$$\text{Solution: } \frac{x}{3x-6} + \frac{2x}{x-2} = 2$$

$$\frac{x + 3(2x)}{3(x-2)} = 2$$

$$\frac{7x}{3(x-2)} = 2$$

Multiplying by $2(x-2)$

$$7x = 2(3)(x-2)$$

$$7x = 6(x-2)$$

$$7x = 6x - 12$$

$$7x - 6x = -12$$

$$x = -12$$

$$\text{Solve Set } x = \{-12\}$$

$$I(vii) \quad \frac{2x}{2x+5} = \frac{2}{3} - \frac{5}{4x+10}, x \neq -\frac{5}{2}$$

$$\text{Solution: } \frac{2x}{2x+5} + \frac{5}{4x+10} = \frac{2}{3}$$

$$\frac{2x}{2x+5} + \frac{5}{2(2x+5)} = \frac{2}{3}$$

MATHEMATICS FOR 9TH CLASS (UNIT # 7)

Pilot Superone Mathematics 314 Class 9th

$$\frac{2(2x) + 5}{2(2x + 5)} = \frac{2}{3}$$

$$\frac{4x + 10}{2(2x + 5)} = \frac{2}{3}$$

Multiplying by $2 \times 3(2x + 5)$

$$2 \times 3(2x + 5) \left(\frac{4x + 10}{2(2x + 5)} \right) = \frac{2}{3} \times 2 \times 3 \times (2x + 5)$$

$$3(4x + 5) = 4(2x + 5)$$

$$12x + 15 = 8x + 20$$

$$12x - 8x = 20 - 15$$

$$4x = 5$$

$$x = \frac{5}{4}$$

Solve Set $x = \left\{ \frac{5}{4} \right\}$

I(viii) $\frac{2x}{x-1} + \frac{1}{3} = \frac{5}{6} + \frac{2}{x-1}, x \neq 1$

Solution: $\frac{2x}{x-1} - \frac{2}{x-1} = \frac{5}{6} - \frac{1}{3}$

$$\frac{2x-2}{x-1} = \frac{5-2}{6}$$

$$\frac{2(x-1)}{x-1} = \frac{3}{6}$$

$$2 = \frac{3}{6}$$

Solution is not possible

MATHEMATICS FOR 9TH CLASS (UNIT # 7)

Pilot Superone Mathematics 315 Class 9th

$$I(ix) \quad \frac{2}{x^2 - 1} - \frac{1}{x + 1} = \frac{1}{x + 1}, x \neq \pm 1$$

$$\text{Solution:} \quad \frac{2}{(x+1)(x-1)} - \frac{1}{x+1} = \frac{1}{x+1}$$

Multiplying by $(x+1)(x-1)$

$$\frac{2(x+1)(x-1)}{(x+1)(x-1)} - (x+1)(x-1) \times \frac{1}{x+1} = (x+1)(x-1) \times \frac{1}{x+1}$$

$$2(x-1) = x-1$$

$$2 - x + 1 = x - 1$$

$$x - x = 1 - 2 + 1$$

$$2x = -4$$

$$x = \frac{-4}{2}$$

$$x = -2$$

$$x \neq \{-2\}$$

Solve Set

$$I(x) \quad \frac{2}{3x+6} = \frac{1}{6} - \frac{1}{2x+4}, x \neq -2$$

$$\text{Solution:} \quad \frac{2}{3(x+2)} = \frac{1}{6} - \frac{1}{2(x+2)}$$

$$\frac{2}{3(x+2)} + \frac{1}{2(x+2)} = \frac{1}{6}$$

$$\frac{2 \times 2 + 1 \times 3}{6(x+2)} = \frac{1}{6}$$

$$\frac{4+3}{6(x+2)} = \frac{1}{6}$$

Multiplying by $6(x+2)$

$$7 = x + 2$$

$$x = 7 - 2$$

$$x = 5$$

MATHEMATICS FOR 9TH CLASS (UNIT # 7)

Pilot Superone Mathematics 316 Class 9th

Q.2 Solve each equation and check for extraneous solution, if any:

- (i) $\sqrt{3x+4} = 2$ (ii) $\sqrt[3]{2x-4} - 2 = 0$
 (iii) $\sqrt{x-3} - 7 = 0$ (iv) $2\sqrt{t+4} = 5$
 (v) $\sqrt[3]{2x+3} = \sqrt[3]{x-2}$ (vi) $\sqrt[3]{2-1} = \sqrt[3]{2t-28}$
 (vii) $\sqrt{2t+6} - \sqrt{2t-5} = 0$ (viii) $\sqrt{\frac{x+1}{2x+5}} = 2x - \frac{5}{2}$

2(i) $\sqrt{3x+4} = 2$

Solution: $(\sqrt{3x+4})^2 = (2)^2$ (Squaring)

$$3x + 4 = 4$$

$$3x = 4 - 4$$

$$3x = 0$$

$$x = 0$$

2(ii) $\sqrt[3]{2x-4} - 2 = 0$

Solution: $\sqrt[3]{2x-4} = 2$

$$(2x-4)^{1/3} = 2$$

$$[(2x-4)^{1/3}]^3 = (2)^3$$
 (Cubing)

$$(2x-4)^{1/3 \cdot 3} = 8$$

$$2x - 4 = 8$$

$$2x = 8 + 4$$

$$2x = 12$$

$$x = \frac{12}{2}$$

$$x = 6$$

MATHEMATICS FOR 9TH CLASS (UNIT # 7)

Pilot Superone Mathematics

317

Class 9th

2(iii) $\sqrt{x-3} - 7 = 0$

Solution:

$$\sqrt{x-3} = 7$$

$$(x-3)^{\frac{1}{2}} = 7$$

$$[(x-3)^{\frac{1}{2}}]^2 = (7)^2 \quad \text{(Squaring)}$$

$$(x-3)^{\frac{1}{2} \times 2} = 49$$

$$x-3 = 49$$

$$x = 49 + 3$$

$$x = 52$$

2(iv) $2\sqrt{t+4} = 5$

Solution:

$$2\sqrt{t+4} = 5$$

$$2(t+4)^{\frac{1}{2}} = 5$$

$$4(t+4)^{\frac{1}{2} \times 2} = (5)^2 \quad \text{(Squaring)}$$

$$4(t+4) = 25$$

$$4t + 16 = 25$$

$$4t = 25 - 16$$

$$4t = 9$$

$$t = \frac{9}{4}$$

2(v) $\sqrt[3]{2x+3} = \sqrt[3]{x-2}$

Solution:

$$\sqrt[3]{2x+3} = \sqrt[3]{x-2}$$

or

$$(2x+3)^{1/3} = (x-2)^{1/3}$$

$$[(2x+3)^{1/3}]^3 = [(x-2)^{1/3}]^3 \quad \text{(Cubing)}$$

$$(2x+3)^{1/3 \times 3} = (x-2)^{1/3 \times 3}$$

$$(2x+3) = x-2$$

$$2x - x = -2 - 3$$

$$x = -5$$

Solve Set

$$x = \{-5\}$$

MATHEMATICS FOR 9TH CLASS (UNIT # 7)

Pilot Superone Mathematics 318 Class 9th

2(vi) $\sqrt[3]{2} - 1 = \sqrt[3]{21 - 28}$

Solution: $\sqrt[3]{2} - 1 = \sqrt[3]{21 - 28}$
 or $(2 - 1)^{1/3} = (21 - 28)^{1/3}$
 $[(2 - 1)^{1/3}]^3 = [(21 - 28)^{1/3}]^3$ (Cubing)
 $(2 - 1)^{1/3 \cdot 3} = (21 - 28)^{1/3 \cdot 3}$
 $2 - 1 = 21 - 28$
 $1 - 21 = -28 - 2$
 $31 = -30$
 $1 = \frac{-30}{3}$
 $1 = 10$

2(vii) $\sqrt{21 - 6} = \sqrt{21 - 5} = 0$

Solution: $\sqrt{21 - 6} = \sqrt{21 - 5}$
 or $(21 - 6)^{1/2} = (21 - 5)^{1/2}$
 $[(21 - 6)^{1/2}]^2 = [(21 - 5)^{1/2}]^2$
 $21 - 6 = 21 - 5$
 $21 - 21 = -5 + 6$
 $0 = 1$

Sol. set is not possible

Sol. set = \emptyset

2(viii) $\sqrt{\frac{x+1}{2x+5}} = 2$

Solution: $\sqrt{\frac{x+1}{2x+5}} = 2$
 or $\left(\frac{x+1}{2x+5}\right)^2 = 2$

MATHEMATICS FOR 9TH CLASS (UNIT # 7)

Pilot Superone Mathematics 319 Class 9th

$$\left[\left(\frac{x+1}{2x+5} \right)^2 \right] = (2)^2 \quad (\text{Squaring})$$

$$\left(\frac{x+1}{2x+5} \right) = 4$$

Multiplying $2x+5$

$$\left(\frac{x+1}{2x+5} \times 2x+5 \right) = 4 \times (2x+5)$$

$$x+1 = 4(2x+5)$$

$$x+1 = 8x+20$$

$$x-8x = 20-1$$

$$-7x = 19$$

$$x = -\frac{19}{7}$$

Equation Involving Absolute Value

Another type of linear equation is the one that contains absolute value. To solve equations involving absolute value we first give the following definition.

Definition

The absolute of a real number 'a' denoted by $|a|$, is defined as

$$|a| = a, \quad \text{if } a \geq 0$$

$$= -a, \quad \text{if } a < 0$$

$$\text{e.g., } |6| = 6, |0| = 0 \text{ and } |-6| = 6$$

Some properties of absolute value

If $a, b \in \mathbb{R}$, then

$$(i) \quad |a| \geq 0$$

$$(ii) \quad |-a| = |a|$$

$$(iii) \quad |ab| = |a| \cdot |b|$$

$$(iv) \quad \left| \frac{a}{b} \right| = \frac{|a|}{|b|}, b \neq 0$$

Exercise 7.2

Q.1 Identify the following statements as True or False.

- (i) $|x| = 0$ has only one solution.
- (ii) All absolute value equations have two solution.
- (iii) The equation $|x| = 2$ is equivalent to $x = 2$ or $x = -2$.
- (iv) The equation $|x - 4| = -4$ has no solution.
- (v) The equation $|2x - 3| = 5$ is equivalent to $2x - 3 = 5$ or $2x + 3 = 5$.

Answers:

(i)	T	(ii)	F	(iii)	T	(iv)	T	(v)	F
-----	---	------	---	-------	---	------	---	-----	---

Q.2 Solve for x:

(i) $|3x - 5| = 4$

(ii) $\frac{1}{2}|3x + 2| - 4 =$

II

(iii) $|2x + 5| = 11$

(iv) $|3 + 2x| = |6x - 7|$

(v) $|x + 2| - 3 = 5 - |x + 2|$

(vi) $\frac{1}{2}|x + 3| + 21 = 9$

(vii) $\left| \frac{3 - 5x}{4} \right| - \frac{1}{3} = \frac{2}{3}$

(viii) $\left| \frac{x + 5}{2 - x} \right| = 2$

2(i) $|3x - 5| = 4$

Solution: $|3x - 5| = 4$

$$3x = 4 + 5$$

$$3x = 9$$

$$x = \frac{9}{3}$$

$$x = 3$$

$$x = 3$$

$$-(3x - 5) = 4$$

$$3x - 5 = -4$$

$$3x = -4 + 5$$

$$3x = 1$$

$$x = \frac{1}{3}$$

$$= \left\{ 3, \frac{1}{3} \right\}$$

MATHEMATICS FOR 9TH CLASS (UNIT # 7)

Pilot Superone Mathematics 321 Class 9th

$$2(ii) \quad \frac{1}{2} |3x + 2| - 4 = 11$$

$$\text{Solution:} \quad \frac{1}{2} |3x + 2| = 11 + 4$$

$$\frac{1}{2} |3x + 2| = 15$$

$$|3x + 2| = 30 \quad (\text{Multiplying by 2})$$

Now

$$3x + 2 = 30$$

$$3x = 30 - 2$$

$$3x = 28$$

$$x = \frac{28}{3}$$

$$-(3x + 2) = 30$$

$$3x + 2 = -30$$

$$3x = -30 - 2$$

$$3x = -32$$

$$x = \frac{-32}{3}$$

$$\text{Sol. Set} = \left\{ \frac{28}{3}, \frac{-32}{3} \right\}$$

$$2(iii) \quad |2x + 5| = 11$$

$$\text{Solution:} \quad 2x + 5 = 11$$

$$2x = 11 - 5$$

$$2x = 6$$

$$x = \frac{6}{2}$$

$$x = 3$$

$$-(2x + 5) = 11$$

$$2x + 5 = -11$$

$$2x = -11 - 5$$

$$2x = -16$$

$$x = \frac{-16}{2}$$

$$x = -8$$

$$\text{Sol. Set} = \{3, -8\}$$

$$2(iv) \quad |3 + 2x| = |6x - 7|$$

$$\text{Solution:} \quad 3 + 2x = 6x - 7$$

$$3 + 7 = 6x - 2x$$

$$10 = 4x$$

$$3 + 2x = -(6x - 7)$$

$$3 + 2x = -6x + 7$$

$$2x + 6x = 7 - 3$$

MATHEMATICS FOR 9TH CLASS (UNIT # 7)

Pilot Superone Mathematics 322 Class 9th

$$x = \frac{10}{4}$$

$$x = \frac{5}{2}$$

$$8x = 4$$

$$x = \frac{4}{8}$$

$$x = \frac{1}{2}$$

Sol. Set $\left\{ \frac{5}{2}, \frac{1}{2} \right\}$

2(v) $|x + 2| - 3 = 5$ $|x + 2|$

Solution: $|x + 2| + 3 = 5 + 3$
 $2|x + 2| = 8$

Dividing by 2

$$|x + 2| = 4$$

$$x + 2 = 4$$

$$x = 4 - 2$$

$$x = 2$$

$$-(x + 2) = 4$$

$$x + 2 = -4$$

$$x = -4 - 2$$

$$x = -6$$

Sol. Set = $\{2, -6\}$

2(vi) $\frac{1}{2}|x + 3| + 21 = 9$

Solution: $\frac{1}{2}|x + 3| = 9 - 21$

$$\frac{1}{2}|x + 3| = -12$$

$$|x + 3| = -24$$

(Multiplying by 2)

Sol. Set = ϕ

(Absolute value is never negative)

2(vii) $\left| \frac{3 - 5x}{4} \right| - \frac{1}{3} = \frac{2}{3}$

Solution: $\left| \frac{3 - 5x}{4} \right| - \frac{1}{3} = \frac{2}{3}$

$$\left| \frac{3 - 5x}{4} \right| = \frac{2}{3} + \frac{1}{3}$$

MATHEMATICS FOR 9TH CLASS (UNIT # 7)

Pilot Superone Mathematics 323 Class 9th

$$\left| \frac{3-5x}{4} \right| = 1$$

Now

$$\frac{3-5x}{4}$$

and

$$\begin{array}{r} 3-5x \\ -4 \\ \hline 3-5x-4 \\ -5x-4 \\ 5x \\ x = -\frac{1}{5} \end{array}$$

$$\left| \frac{3-5x}{4} \right| = 1$$

$$\begin{array}{r} 3-5x \\ 4 \\ \hline 3-5x \\ 4 \\ 3-5x-4 \\ -5x-4 \\ 5x \\ 5x \\ x = \frac{7}{5} \end{array}$$

$$\text{Sol Set} = \left\{ -\frac{1}{5}, \frac{7}{5} \right\}$$

$$2(\text{viii}) \quad \left| \frac{x+5}{2-x} \right| = 6$$

Solution:

$$\begin{array}{r} \frac{x+5}{2-x} = 6 \\ x+5 = 6(2-x) \\ x+5 = 12-6x \\ x+6x = 12-5 \\ 7x = 7 \\ x = 1 \end{array}$$

$$\left| \frac{x+5}{2-x} \right| = 6$$

$$\begin{array}{r} \frac{x+5}{2-x} = 6 \\ x+5 = 6(2-x) \\ x+5 = 12-6x \\ x+6x = 12-5 \\ 7x = 7 \\ x = \frac{17}{5} \end{array}$$

$$\text{Sol Set} = \left\{ 1, \frac{17}{5} \right\}$$

Defining Inequalities

Let a, b be real numbers. Then a is greater than b if the difference $a - b$ is positive and we denote this order relation by the inequality $a > b$. An equivalent statement is that b is less

==
re.

MATHEMATICS FOR 9TH CLASS (UNIT # 7)

Pilot Superone Mathematics 324 Class 9th

than a symbolized by $b < a$. Similarly, if a, b is negative, then a is less than b and expressed in symbols as $a < b$.

Properties of Inequalities

1. Law of Trichotomy

For any $a, b \in \mathbb{R}$, one and only one of the following statements is true

$$a < b \text{ or } a = b, \text{ or } a > b$$

An important special case of this property is the case for $b = 0$, namely

$$a < 0 \text{ or } a = 0 \text{ or } a > 0 \text{ for any } a \in \mathbb{R}.$$

2. Transitive Property

Let $a, b, c \in \mathbb{R}$

- (i) If $a > b$ and $b > c$, then $a > c$
- (ii) If $a < b$ and $b < c$, then $a < c$

3. Additive Closure Property

For $a, b, c \in \mathbb{R}$,

- (i) If $a > b$, then $a + c > b + c$
 If $a < b$, then $a + c < b + c$
- (ii) If $a > 0$ and $b > 0$, then $a + b > 0$
 If $a < 0$ and $b < 0$, then $a + b < 0$

4. Multiplicative Property

Let $a, b, c, d \in \mathbb{R}$

- (i) If $a > 0$ and $b > 0$, then $ab > 0$.
 whereas $a < 0$ and $b < 0 \Rightarrow ab > 0$
- (ii) If $a > b$ and $c > 0$, then $ac > bc$
 or if $a < b$ and $c > 0$, then $ac < bc$
- (iii) If $a > b$ and $c < 0$, then $ac < bc$
 or if $a < b$ and $c < 0$, then $ac > bc$

The above property (iii) states that the sign of inequality is reversed if each side is multiplied by a negative real number

- (iv) If $a > b$ and $c > d$, then $ac > bd$

MATHEMATICS FOR 9TH CLASS (UNIT # 7)

Pilot Superone Mathematics 325 Class 9th

Exercise 7.3

Q.1 Solve the following inequalities.

(i) $3x + 1 < 5x - 4$ (ii) $4x - 10.3 \leq 21x - 1.8$

(iii) $4 - \frac{1}{2}x \geq -7 + \frac{1}{4}x$ (iv) $x - 2(5 - 2x) \leq 6x -$

$3\frac{1}{2}$

(v) $\frac{3x+2}{9} - \frac{2x+1}{3} > -1$

(vi) $3(2x + 1) - 2(2x + 5) < 5(3x - 2)$

(vii) $3(x - 1) - (x - 2) > -2(x + 4)$

(viii) $2\frac{2}{3}x + \frac{2}{3}(5x - 4) > \frac{1}{3}(8x + 7)$

1(i) $3x + 1 > 5x - 4$

Solution:

$$3x + 1 > 5x - 4$$

$$3x + 1 - 5x > 5x - 4 - 5x$$

$$-2x + 1 < -4$$

$$-2x + 1 - 1 < -4 - 1$$

$$-2x < -5 \quad \text{Dividing by } -2$$

$$x > \frac{5}{2} \quad \text{As } -1 \text{ is change of sign}$$

$$\text{Sol. Set} = \{x | x > \frac{5}{2}\}$$

1(ii) $4x - 10.3 \leq 21x - 1.8$

Solution:

(Subtracting $21x$)

$$4x - 10.3 - 21x < 21x - 1.8 - 21x$$

$$-17x - 10.3 < -1.8$$

$$-17x - 10.3 + 10.3 \leq -1.8 + 10.3 \quad \text{(Adding 10.3)}$$

$$-17x < 8.5$$

MATHEMATICS FOR 9TH CLASS (UNIT # 7)

Pilot Superone Mathematics 326 Class 9th

N.T.S. Dividing by 17 (change of sign)

$$x > \frac{8.5}{-17}$$

$$x \geq -5$$

$$\text{Sol. Set } \{x|x \geq -5\}$$

$$1(\text{iii}) \quad 4 - \frac{1}{2}x \geq -7 + \frac{1}{4}x$$

$$\text{Solution:} \quad 4 - \frac{1}{2}x \geq -7 + \frac{1}{4}x$$

Multiplying by 4

$$16 - 2x \geq -28 + x$$

Subtracting by x

$$16 - 2x - x \geq -28 + x - x$$

$$16 - 3x \geq -28$$

Subtracting by 16

$$16 - 3x - 16 \geq -28 - 16$$

$$-3x \geq -44$$

Subtracting by -3

$$x \leq \frac{-44}{-3} \quad \text{N.T.S (change of}$$

sign)

$$x < \frac{44}{3}$$

$$\text{Sol. Set} = \{x|x < \frac{44}{3}\}$$

$$1(\text{iv}) \quad x - 2(5 - 2x) \geq 6x - 3\frac{1}{2}$$

$$\text{Solution:} \quad x - 2(5 - 2x) \geq 6x - 3\frac{1}{2}$$

MATHEMATICS FOR 9TH CLASS (UNIT # 7)

Pilot Superone Mathematics 327 Class 9th

$$x - 10 + 4x \geq 6x \quad \frac{7}{2}$$

Multiplying by 2

$$2x - 20 + 8x > 12x \quad 7$$

$$10x - 20 \geq 12x \quad 7$$

Subtracting 12x

$$10x - 20 - 12x > 12x - 7 - 12x$$

$$-2x - 20 \geq -7$$

Adding 20

$$-2x - 20 + 20 \geq -7 + 20$$

$$-2x \geq 13$$

Dividing by -2

$$x \leq -\frac{13}{2} \quad (\text{change of sign})$$

$$x \leq -6.5$$

$$\text{Sol Set} = \{x | x \leq -6.5\}$$

$$I(v) \quad \frac{3x+2}{9} - \frac{2x+1}{3} > -1$$

Solution:

Multiplying by 9

$$\frac{9(3x+2)}{9} - \frac{9(2x+1)}{3} > -1(9)$$

$$3x+2 - 3(2x+1) > -9$$

$$3x+2 - 6x - 3 > -9$$

$$-3x - 1 > -9$$

$$-3x - 1 + 1 > -9 + 1 \quad (\text{Adding 1})$$

$$-3x > -8$$

Dividing by -3 (change of sign)

$$x < \frac{-8}{-3}$$

MATHEMATICS FOR 9TH CLASS (UNIT # 7)

Pilot Superone Mathematics 328 Class 9th

$$x < \frac{8}{3}$$

1(vi) $3(2x + 1) - 2(2x + 5) < 5(3x - 2)$

Solution: $6x + 3 - 4x - 10 < 15x - 10$
 $2x - 7 < 15x - 10$

Subtracting $15x$

$$2x - 7 - 15x < 15x - 10 - 15x$$

$$13x - 7 < -10$$

$$13x - 7 + 7 < -10 + 7 \quad (\text{Adding } 7)$$

$$13x < -3$$

Dividing by -13 (change of sign)

$$x > \frac{-3}{-13}$$

$$x > \frac{3}{13}$$

$$\text{Sol. Set} = \{x | x > \frac{3}{13}\}$$

1(vii) $3(x - 1) - (x - 2) > -2(x + 4)$

Solution: $3x - 3 - x + 2 > -2x - 8$
 $2x - 1 > -2x - 8$
 $2x + 2x - 1 > -2x + 2x - 8 \quad (\text{Adding } 2x)$

2x)

$$4x - 1 + 1 > -8 + 1 \quad (\text{Adding } 1)$$

$$4x > -7$$

$$x > \frac{-7}{4} \quad (\text{Dividing by } 4)$$

1(viii) $2\frac{2}{3}x + \frac{2}{3}(5x - 4) > -\frac{1}{3}(8x + 7)$

Solution: $\frac{4x}{3} + \frac{2}{3}(5x - 4) > -\frac{1}{3}(8x + 7)$

$$8x + 2(5x - 4) > -(8x + 7) \quad (\text{Multiplying by } 3)$$

$$8x + 10x - 8 > -8x - 7$$

$$18x - 8 > -8x - 7$$

Adding by $8x$

MATHEMATICS FOR 9TH CLASS (UNIT # 7)

Pilot Superone Mathematics 329 Class 9th

$$\begin{aligned} 18x - 8 + 8x &> 8x - 7 + 8x \\ 26x - 8 &> -7 && \text{(Adding by 8)} \\ 26x - 8 + 8 &> -7 + 8 \\ 26x &> 1 \\ x &> \frac{1}{26} && \text{(Dividing by 26)} \end{aligned}$$

$$\text{Sol. Set} = \{x | x > \frac{1}{26}\}$$

Q.2 Solve the following inequalities.

(i) $-4 < 3x + 5 < 8$ (ii) $-5 \leq \frac{4 - 3x}{2} < 1$

(iii) $-6 < \frac{x - 2}{4} < 6$ (iv) $3 \geq \frac{7 - x}{2} \geq 1$

(v) $3x - 10 \leq 5 < x + 3$ (vi) $-3 \leq \frac{x - 4}{5} < 4$

(vii) $1 - 2x < 5$ $x \leq 25 - 6x$

(viii) $3x - 2 < 2x + 1 < 4x + 17$

2(i) $-4 < 3x + 5 < 8$

Solution:
$$\begin{aligned} -4 < 3x + 5 < 8 \\ -4 - 5 < 3x + 5 - 5 < 8 - 5 && \text{(Subtracting 5)} \\ 9 < 3x < 3 \\ 3 < x < 1 && \text{(Dividing by 3)} \end{aligned}$$

2(ii) $-5 \leq \frac{4 - 3x}{2} < 1$

Solution:
$$\begin{aligned} 10 &\leq 4 - 3x < 2 && \text{(Multiplying by 2)} \\ 10 - 4 &\leq 4 - 3x - 4 < 2 - 4 && \text{(Subtracting 4)} \\ 14 &\leq -3x < -2 \end{aligned}$$

(Dividing by -3 (change of sign))

$$\frac{14}{-3} \geq \frac{-3x}{-3} > \frac{-2}{-3}$$

$$\frac{14}{3} \geq x > \frac{2}{3}$$

$$\frac{2}{3} > x \leq \frac{14}{3}$$

MATHEMATICS FOR 9TH CLASS (UNIT # 7)

Pilot Superine Mathematics 330 Class 9th

$$2(\text{iu}) \quad -6 < \frac{x-2}{4} < 6$$

Solution: $24 \cdot \frac{x-2}{4} < 24$ (Multiplying by 4)
 $24 \cdot 2 < x-2+2 < 24+2$ (Adding 2)
 $22 < x < 26$

$$2(\text{iv}) \quad 3 \geq \frac{7}{2}x \geq 1$$

Solution: $6 \geq 7x \geq 2$ (Multiplying by 2)
 $6-7 \geq 7x-7 \geq 2-7$ (Subtracting 7)
 $-1 \geq x \geq -5$

Multiplying by -1 (change of sign)

$$1 \leq x \leq 5$$

$$2(\text{v}) \quad 3x-10 \leq 5 < x+3$$

Solution: Here

$$3x-10 < 5$$

$$3x-10+10 < 5+10$$

$$3x < 15$$

$$\frac{3x}{3} < \frac{15}{3}$$

$$x < 5 \quad (i)$$

$$5 < x+3$$

$$\text{or } x+3 > 5$$

$$x+3-3 > 5-3$$

$$x > 2$$

(ii)

$$\text{or } 2 < x$$

From (i) and (ii)

$$2 < x \leq 5$$

$$2(\text{vi}) \quad -3 \leq \frac{x-4}{-5} < 4$$

Solution: Multiplying by -5 (Change of sign)

$$15 \geq x-4 > -20$$

$$15+4 \geq x-4+4 > -20+4 \quad (\text{Adding } +4)$$

$$19 \geq x > -16$$

$$\text{or } -16 < x \leq 19$$

MATHEMATICS FOR 9TH CLASS (UNIT # 7)

Pilot Superone Mathematics 331 Class 9th

2(vii) $1 - 2x < 5$ & $x \leq 25 - 6x$

Solution:

$ \begin{aligned} 1 - 2x &< 5 && -x \\ 1 - 2x + x &< 5 && -x + x \\ 1 - x &< 5 && \\ -x &< 5 - 1 && \\ -x &< 4 && \\ \text{or } x &> 4 && \quad (i) \end{aligned} $	$ \begin{aligned} 5 - x &< 25 - 6x \\ 5 - x + 6x &< 25 - 6x + 6x \\ 5 + 5x &< 25 \\ 5 + 5x - 5 &\leq 25 - 5 \\ 5x &\leq 20 \\ \frac{5x}{5} &\leq \frac{20}{5} \\ x &\leq 4 && \quad (ii) \end{aligned} $
--	--

From (i) and (ii)

$$4 < x < 4$$

2(viii) $3x - 2 < 2x + 1 < 4x + 17$

Solution: Here

$ \begin{aligned} 3x - 2 &< 2x + 1 \\ 3x - 2 - 1 &< 2x + 1 - 1 \\ 3x - 3 &< 2x \\ 3x - 2x - 3 &< 2x - 2x \\ x - 3 &< 0 \\ x - 3 + 3 &< 0 + 3 \\ x &< 3 && \quad (i) \end{aligned} $	$ \begin{aligned} 2x + 1 &< 4x + 17 \\ 2x + 1 - 1 &< 4x + 17 - 1 \\ 2x &< 4x + 16 \\ 2x - 4x &< 4x + 16 - 4x \\ -2x &< 16 \\ \text{Dividing by 2 (Change of sign)} \\ x &> \frac{16}{-2} \\ x &> -8 && \quad (ii) \end{aligned} $
---	--

From (i) and (ii)

$$-8 < x < 3$$

MATHEMATICS FOR 9TH CLASS (UNIT # 7)

Pilot Superone Mathematics 332 Class 9th

Review Exercise 7

Q.1 Multiple choice question, choose the correct answers.

- (i) Which of the following is the solution of the inequality $3 - 4x < 11$?
- (a) 8 (b) 2
- (c) $\frac{14}{4}$ (d) None of these
- (ii) A statement involving any of the symbols $<$ or \geq is called
- (a) equation (b) identity
- (c) inequality (d) linear equation
- (iii) x is a solution of the inequality $-2 < x < \frac{3}{2}$.
- (a) 5 (b) 3
- (c) 0 (d) $\frac{3}{2}$
- (iv) If x is no larger than 10, then
- (a) $x > 8$ (b) $x = 10$
- (c) $x = 10$ (d) $x = 10$
- (v) If the capacity c of an elevator is at most 1600 pounds, then
- (a) $c < 1600$ (b) $c = 1600$
- (c) $c \geq 1600$ (d) $c > 1600$
- (vi) $x = 0$ is a solution of the inequality
- (a) $x = 0$ (b) $3x - 5 < 0$
- (c) $x + 2 < 0$ (d) $x - 2 < 0$

Answers:

- (i) b (ii) c (iii) c (iv) b
- (v) c (vi) d

MATHEMATICS FOR 9TH CLASS (UNIT # 7)

Pilot Superone Mathematics

333

Class 9th

Q.2 Identify the following statements as True or False.

- (i) The equation $3x - 5 = 7 - x$ is a linear equation.
- (ii) The equation $x = 0, 3x = 0, 7x$ is an identity.
- (iii) The equation $2x + 3 = 8$ is equivalent to $2x = 11$.
- (iv) To eliminate fractions, we multiply each side of an equation by the L.C.M. of denominators.
- (v) $4(x + 3) = x + 3$ is a conditional equation.
- (vi) The equation $2(3x + 5) = 6x + 12$ is an inconsistent equation.
- (vii) To solve $\frac{2}{3}x = 12$, we should multiply each side by $\frac{2}{3}$.
- (viii) Equations having exactly the same solution are called equivalent equations.
- (ix) A solution that does not satisfy the original equation is called extraneous solution.

Answers:

(i)	True	(ii)	True	(iii)	False	(iv)	True
(v)	True	(vi)	True	(vii)	False	(viii)	True
(ix)	True						

Q.3 Answer the following short question.

- (i) Define a linear inequality in one variable.
- (ii) State the trichotomy and transitive properties of inequalities.
- (iii) The formula relating degrees Fahrenheit to degrees Celsius is $F = \frac{9}{5}C + 32$. For what value of C is $F < 0$?
- (iv) Seven times the sum of an integer and 12 is at least 50 and at most 60. Write and solve the inequality that expresses this relationship.

MATHEMATICS FOR 9TH CLASS (UNIT # 7)

Pilot Superone Mathematics 334 Class 9th

Answer:

(i) A linear inequality in one variable x is $ax + b < 0$, $a \neq 0$

(ii) **Trichotomy Property.**

$\forall a, b \in \mathbb{R}$, one and only one of the following statement is true

$a < b$ or $a = b$ or $a > b$

If $b = 0$ then $a < 0$ or $a = 0$ or $a > 0$ for any $a \in \mathbb{R}$.

Transitive Property:

$\forall a, b, c \in \mathbb{R}$

(i) If $a > b$ and $b > c$ then $a > c$

(ii) If $a < b$, and $b < c$ then $a < c$

(iii) $-\frac{9}{5}C + 32 < 0$

$$C < 32 \times \frac{5}{9}$$

$$C < \frac{160}{9}$$

(iv) Let the integer = x

According to the conditions

$$50 \leq 7(x + 12) \leq 60$$

Solution: $50 < 7x + 84 \leq 60$

$$50 - 84 < 7x + 84 - 84 \quad 84 < 60 - 84 \quad \text{(Subtracting 84)}$$

$$-34 < 7x \leq -24$$

$$-\frac{34}{7} < \frac{7x}{7} \leq \frac{-24}{7}$$

Q.4 Solve each of the following and check for extraneous solution if any

(i) $\sqrt{21+4} = \sqrt{1-1}$

(ii) $\sqrt{3x-1} - 2\sqrt{8-2x} = 0$

MATHEMATICS FOR 9TH CLASS (UNIT # 7)

Pilot Superone Mathematics 335 Class 9th

4(i) $\sqrt{2t+4} = \sqrt{t-1}$

Solution:
$$\begin{aligned} 2t+4 &= t-1 & (\text{Squaring}) \\ 2t-t &= -1-4 \\ t &= -5 \end{aligned}$$

Check Put $t = -5$

$$\begin{array}{ll} \sqrt{2t+4} & \sqrt{t-1} \\ \sqrt{2(-5)+4} & \sqrt{-5-1} \\ \sqrt{10+4} & \sqrt{-6} \\ \sqrt{6} & \end{array}$$

Solution in integers in null set (ϕ)

4(ii) $\sqrt{3x-1} - 2\sqrt{8-2x} = 0$

Solution:
$$\begin{aligned} \sqrt{3x-1} - 2\sqrt{8-2x} &= 0 \\ \sqrt{3x-1} &= 2\sqrt{8-2x} \end{aligned}$$

Squaring

$$\begin{aligned} 3x-1 &= 4(8-2x) \\ 3x-1 &= 32-8x \\ 3x+8x &= 32+1 \\ 11x &= 33 \\ x &= \frac{33}{11} \\ x &= 3 \end{aligned}$$

Check

$$\begin{aligned} &= \sqrt{3x-1} - 2\sqrt{8-2x} \\ \text{Putting } x &= 3 \\ &= \sqrt{3(3)-1} - 2\sqrt{8-2(3)} \\ &= \sqrt{9-1} - 2\sqrt{8-6} \\ &= \sqrt{8} - 2\sqrt{2} \end{aligned}$$

MATHEMATICS FOR 9TH CLASS (UNIT # 7)

Pilot Superone Mathematics 336 Class 9th

$$2\sqrt{2} - 2\sqrt{2} = 0 = \text{R.H.S.}$$

Q.5 Solve for x

$$(i) \quad |3x + 14| - 2 = 5x \quad (ii) \quad \frac{1}{3}|x - 3| = \frac{1}{2}|x +$$



5(i) $|3x + 14| - 2 = 5x$

Solution: $|3x + 14| - 2 = 5x$
 $|3x + 14| = 5x + 2$

$$3x + 14 = 5x + 2$$

$$3x - 5x = 2 - 14$$

$$-2x = -12$$

$$x = \frac{-12}{-2}$$

$$x = 6$$

$$-(3x + 14) = 5x + 2$$

$$-3x - 14 = 5x + 2$$

$$-3x - 5x = 2 + 14$$

$$-8x = 16$$

$$x = \frac{16}{-8}$$

$$x = -2$$

Check: L.H.S.

$$|3x + 14| - 2$$

$$|3(6) + 14| - 2$$

$$= |18 + 14| - 2$$

$$= |32| - 2$$

$$32 - 2$$

$$30$$

R.H.S.

$$x - 6$$

$$5(x)$$

$$= 5 \times 6$$

$$= 30$$

True

$$x = -2$$

Now check

$$|3x + 14| - 2 = 5x + 2$$

L.H.S.

$$|3(-2) + 14| - 2$$

$$|-6 + 14|$$

$$|8|$$

R.H.S.

$$5x + 2$$

$$= 5(-2) + 2$$

$$= -10 + 2$$

$$= -8$$

MATHEMATICS FOR 9TH CLASS (UNIT # 7)

Pilot Superone Mathematics 337 Class 9th

$$|8| = -8 \quad \text{False}$$

$$\text{Sol. Set} = \{6\}$$

$$5(ii) \quad \frac{1}{3}|x-3| = \frac{1}{2}|x+2|$$

$$\frac{1}{3}(x-3) = \frac{1}{2}(x+2)$$

Divide on 6

$$2(x-3) = 3(x+2)$$

$$2x-6 = 3x+6$$

$$2x-3x = 6+6$$

$$-x = 12$$

$$x = -12$$

$$\frac{1}{3}(x-3) = -\frac{1}{2}(x+2)$$

Divide on 6

$$2(x-3) = -3(x+2)$$

$$2x-6 = -3x-6$$

$$2x-5x = -6+6$$

$$7x = 0$$

$$x = 0$$

$$\text{Set } \{-12, 0\}$$

Q.6 Solve the following inequality.

$$(i) \quad -\frac{1}{3}x + 5 \leq 1 \quad (ii) \quad -3 < \frac{1-2x}{5} < 1$$

$$6(i) \quad -\frac{1}{3}x + 5 \leq 1$$

$$\text{Solution:} \quad -\frac{1}{3}x + 5 - 5 < 1 - 5$$

$$-\frac{1}{3}x + 5 - 5 \leq 1 - 5$$

$$-\frac{1}{3}x < -4$$

Multiplying by 3 (Change of sign)

$$(-3) \left(-\frac{1}{3}x \right) \geq (-3)(-4)$$

$$x \geq 12$$

MATHEMATICS FOR 9TH CLASS (UNIT # 7)

Pilot Superone Mathematics 338 Class 9th

6(ii) $-3 < 1 - \frac{2x}{5} < 1$

Solution: $-15 < 1 - 2x < 5$ Multiplying by 5

$15 - 1 < 1 - 2x < 5 - 1$ Subtracting 1

$16 < -2x < 4$

Dividing by -2 (change of sign)

$$\frac{-16}{-2} > \frac{-2x}{-2} > \frac{4}{-2}$$

$$8 > x > -2$$

MATHEMATICS FOR 9TH CLASS (UNIT # 8)

Pilot Super One Mathematics

439

Class 9th

**Unit
8**

LINEAR GRAPHS & THEIR APPLICATION

An Ordered Pair of Real Numbers

An ordered pair of real numbers x and y is a pair (x, y) in which elements are written in specific order i.e.,

- (i) (x, y) is an ordered pair in which first element is x and second is y such that $(x, y) \neq (y, x)$ where,
- (ii) $(2, 3)$ and $(3, 2)$ are two different ordered pairs.
- (iii) $(x, y) = (m, n)$ only if $x = m$ and $y = n$.

Cartesian Plane

The Cartesian plane establishes one-to-one correspondence between the set of ordered pairs $R \times R = \{(x, y) : x, y \in R\}$ and the points of the Cartesian plane.

In plane two mutually perpendicular straight lines are drawn. The lines are called the coordinate axes. The point O , where the two lines meet is called origin. This plane is called the coordinate plane or the Cartesian plane.


MATHEMATICS FOR 9TH CLASS (UNIT # 8)

Pilot Superone Mathematics 340 Class 9th

EXERCISE 8.1

Q.1 Determine the quadrant of the coordinate plane in which the following points lie: $P(-4, 3)$, $Q(-5, -2)$, $R(2, 2)$ and $S(2, -6)$.

Ordered Pair	Quadrant
$S(2, -6)$	IV - Q
$R(2, 2)$	I - Q
$Q(-5, -2)$	III - Q
$P(-4, 3)$	II - Q



Q.2 Draw the graph of each of the following.

(i) $x = 2$ (ii) $x = -3$ (iii) $y = -1$

(iv) $y = 3$ (v) $y = 0$ (vi) $x = 0$

(vii) $y = 3x$ (viii) $-y = 2x$ (ix) $\frac{1}{2} = x$

(x) $3y - 5x$ (xi) $2x - y = 0$ (xii) $2x - y = 2$

(xiii) $x - 3y + 1 = 0$ (xiv) $3x - 2y + 1 = 0$

(v) $x = 2$

MATHEMATICS FOR 9TH CLASS (UNIT # 8)


Pilot Superone Mathematics 341

Class 9th

1(i) $x = 2$

Solution: Table ordered pairs that lie on the graph of $x = 2$

$x =$	2	2	2	2
$y =$	2	4	2	-4



Graph $x = 2$

2(ii) $x = -3$

Solution: Table ordered pairs that lie on the graph of $x = -3$

$x =$...	3	3	-3	3	3
$y =$...	2	5	-2	4	...


MATHEMATICS FOR 9TH CLASS (UNIT # 8)

Prior Superone Mathematics 312 Class 9th

2(n) $x = -3$

Solution: Table ordered pairs that lie on the graph of $x = -3$

x	-3	-3	-3	-3	-3
y	1	2	3	4	5




Graph $x = -3$

2(n) $y = 1$

Solution: Table ordered pairs that lie on the graph of $y = 1$

x	1	2	3	4	5	6
y	1	1	1	1	1	1



Graph $y = 1$

MATHEMATICS FOR 9TH CLASS (UNIT # 8)

Pilot Superong Mathematics 343 Class 9th

2(iv) $y = 3$

Solution: Table ordered pairs that lie on the graph of $y = 3$

x	-3	-1	1	3	5	6
y	3	3	3	3	3	3




Graph $y = 3$

2(v) $y = 0$

Solution: Table ordered pairs that lie on the graph of $y = 0$

x	-3	-1	1	3	5	6
y	0	0	0	0	0	0



Graph $y = 0$

MATHEMATICS FOR 9TH CLASS (UNIT # 8)


Pilot Superone Mathematics 344 Class 9th

This is x-axis

2(vi) $x = 0$

Solution: Table ordered pairs that lie on the graph of $x = 0$

X =	0	0	0	0	0	0
y =		3	5	2	-6	



Graph $x = 0$


This is y-axis

2(vii) $y = 3x$

Solution: Table ordered pairs that lie on the graph of $y = 3x$

x =		2	-1	2
y =	3	6	-3	-6

Plotting the points on the graph paper



Graph of $y = 3x$

MATHEMATICS FOR 9TH CLASS (UNIT # 8)


Pilot Superone Mathematics 345

Class 9th

2(viii) $-y = 2x$

Solution: Table ordered pairs that lie on the graph of $y = 2x$

$x =$	1	-2	1	4
$y =$	2	4	2	-8




Graph $-y = 2x$

2(ix) $\frac{1}{2} = x$

or $x = \frac{1}{2}$

Solution: Table ordered pairs that lie on the graph of $x = \frac{1}{2}$.

$x =$	$\frac{1}{2}$	$\frac{1}{2}$	$\frac{1}{2}$	$\frac{1}{2}$
$y =$	2	2	-3	



Graph $x = \frac{1}{2}$


MATHEMATICS FOR 9TH CLASS (UNIT # 8)

Pilot Superone Mathematics 346 Class 9th

2(x) - 3y = 5x

Solution: $y = -\frac{5}{3}x$

Table ordered pairs that lie on the graph of $3y = -5x$




Graph $3y = -5x$

2(x) - 1x - y = 0

Solution: $y = x$

Table ordered pairs that lie on the graph of $2x - y = 0$



Graph $2x - y = 0$

MATHEMATICS FOR 9TH CLASS (UNIT # 8)


Pilot Superline Mathematics 347 Class 9th

2(vi) $2x - y = 2$

Solution. $y = 2 - 2x$

Table ordered pairs that lie on the graph of $2x - y = 2$

x	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30	31	32	33	34	35	36	37	38	39	40	41	42	43	44	45	46	47	48	49	50	51	52	53	54	55	56	57	58	59	60	61	62	63	64	65	66	67	68	69	70	71	72	73	74	75	76	77	78	79	80	81	82	83	84	85	86	87	88	89	90	91	92	93	94	95	96	97	98	99	100
y	0	2	4	6	8	10	12	14	16	18	20	22	24	26	28	30	32	34	36	38	40	42	44	46	48	50	52	54	56	58	60	62	64	66	68	70	72	74	76	78	80	82	84	86	88	90	92	94	96	98	100																																																	




2(vii) $x - 3y + 1 = 0$

Solution. $x = 3y - 1$

Table ordered pairs that lie on the graph of $x - 3y + 1 = 0$

x	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30	31	32	33	34	35	36	37	38	39	40	41	42	43	44	45	46	47	48	49	50	51	52	53	54	55	56	57	58	59	60	61	62	63	64	65	66	67	68	69	70	71	72	73	74	75	76	77	78	79	80	81	82	83	84	85	86	87	88	89	90	91	92	93	94	95	96	97	98	99	100	
y	0	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30	31	32	33	34	35	36	37	38	39	40	41	42	43	44	45	46	47	48	49	50	51	52	53	54	55	56	57	58	59	60	61	62	63	64	65	66	67	68	69	70	71	72	73	74	75	76	77	78	79	80	81	82	83	84	85	86	87	88	89	90	91	92	93	94	95	96	97	98	99	100



MATHEMATICS FOR 9TH CLASS (UNIT # 8)

Pilot Superone Mathematics 348 Class 9th


2(xiv) $3x - 2y + 1 = 0$

Solution: $-2y = -1 - 3x$

or $y = \frac{1+3x}{2}$

Table ordered pairs that lie on the graph of
 $3x - 2y + 1 = 0$.

$x =$	1	-1	3
$y =$	2	-1	5



Graph $3x - 2y + 1 = 0$

Q.3 Are the following lines (i) parallel to x-axis (ii) parallel to y-axis

(i) $2x - 1 = 3$ (ii) $x + 2 = 1$ (iii) $2y + 3 = 2$

(iv) $x + y = 0$ (v) $2x - 2y = 0$

3(i) $x - 1 = 3$

Solution. $2x - 1 = 3$

$2x = 3 + 1$

$2x = 4$

MATHEMATICS FOR 9TH CLASS (UNIT # 8)

Pilot Superone Mathematics 349

Class 9th

$$x = \frac{-1}{2} = -\frac{1}{2}$$

Parallel to y-axis

3(ii) $x + 2 = -1$

Solution: $x + 2 = -1$

$$x = -1 - 2$$

$$x = -3$$

Parallel to y-axis

3(iii) $2y + 3 = 2$

Solution: $2y + 3 = 2$

$$2y = 2 - 3$$

$$2y = -1$$

$$y = -\frac{1}{2}$$


Parallel to x-axis

3(iv) $x + y = 0$

Solution: $x + y = 0$

$$y = -x$$

x =	3	-4
y =	-3	4



x, y = 0

MATHEMATICS FOR 9TH CLASS (UNIT # 8)

Pilot Superior Mathematics 350 (Class 9)

It is neither parallel to x-axis, nor to y-axis


3(x) $2x - 2x = 0$

Solution: $2x - 2y = 0$

$2y = 2x$

$y = x$

x =	4	3
y =	4	3



It is neither parallel to x-axis, nor to y-axis

Q.4 Find the value of m and c of the following lines by expressing them in the form $y = mx + c$

(i) $2x + 3y - 1 = 0$

(ii) $x - 2y = -2$

(iii) $3x + y - 1 = 0$

(iv) $2x - y = 7$

(v) $3 - 2x + y = 0$

(vi) $2x = y + 3$

Sol. (i) $2x + 3y - 1 = 0$

Solution. $2x + 3y - 1 = 0$

or $3y = -2x + 1$

MATHEMATICS FOR 9TH CLASS (UNIT # 8)

Pilot Superone Mathematics 351 Class 9th

or $y = -\frac{2}{3}x + \frac{1}{3}$ (i) (Dividing by 3)

$y = mx + c$ (ii) Comparing (i) and (ii)

$m = -\frac{2}{3}$

$c = \frac{1}{3}$

4(ii) $x - 2y = 2$

Solution: $x - 2y = 2$

$2y = x - 2$

$-\frac{2}{2}y = \frac{-x}{2} - \frac{2}{2}$ (Dividing by -2)

$y = \frac{1}{2}x + 1$ (Comparing it with)

$y = mx + c$

$m = \frac{1}{2}, c = 1$

4(iii) $3x + y - 1 = 0$

Solution: $3x + y - 1 = 0$

or $y = -3x + 1$ (i)

$y = mx + c$ (ii) Comparing (i) and (ii)

$m = -3, c = 1$

4(iv) $2x - y = 7$

Solution: $2x - y = 7$

or $y = -2x + 7$

or $y = 2x - 7$ (i)

$y = mx + c$ (ii) Comparing (i) and (ii)

$m = 2, c = -7$

MATHEMATICS FOR 9TH CLASS (UNIT # 8)

Pilot Superone Mathematics 352 Class 9th

4(v) $3 - 2x + y = 0$

Solution: $3 - 2x + y = 0$

or $y = 2x - 3$ (i)

$y = mx + c$ (ii) Comparing (i) and (ii)

$m = 2, c = -3$

4(vi) $2x = y + 3$

Solution: $2x = y + 3$

or $y + 3 = 2x$

or $y = 2x - 3$ (i)

$y = mx + c$ (ii) Comparing (i) and (ii)

$m = 2, c = -3$

Q.5 Verify whether the following points lie on the line $2x - y + 1 = 0$ or not.

(i) (2, 3) (ii) (0, 0) (iii) (-1, 1)

(iv) (2, 5) (v) (5, 3)

5(i) (2, 3)

Solution: $2x - y + 1 = 0$

$-y = -2x + 1$

$y = 2x + 1$ (i) (2, 3)

Putting $x = 2, y = 3$ in the equation

$3 = 2(2) + 1$

$3 = 4 + 1$

$3 = 5$ (Impossible)

The point (2, 3) does not lie on the line.

5(ii) (0, 0)

Solution: $2x - y + 1 = 0$

$y = 2x + 1$

Putting $x = 0, y = 0$ in the equation

$0 = 2(0) + 1$

$0 = 0 + 1$ (Impossible)

MATHEMATICS FOR 9TH CLASS (UNIT # 8)

Pilot Superone Mathematics 353 Class 9th

The point (0, 0) does not lie on the line.

5(iii) (-1, 1)

Solution: $2x - y + 1 = 0$

$$y = 2x + 1$$

Putting $x = -1, y = 1$ in the equation

$$1 = 2(-1) + 1$$

$$1 = -2 + 1$$

$$1 = -1 \quad (\text{Impossible})$$

The point (-1, 1) does not lie on the line.

5(iv) (2, 5)

Solution: $2x - y + 1 = 0$

$$y = 2x + 1$$

Putting $x = 2, y = 5$ in the equation

$$5 = 2(2) + 1$$

$$5 = 4 + 1$$

$$5 = 5 \quad (\text{True})$$

The point (2, 5) lie on the line.

5(v) (5, 3)

Solution: $2x - y + 1 = 0$

$$y = 2x + 1$$

Putting $x = 5, y = 3$ in the equation

$$3 = 2(5) + 1$$

$$3 = 10 + 1$$

$$3 = 11 \quad (\text{Impossible})$$

The point (5, 3) does not lie on the graph of the line

MATHEMATICS FOR 9TH CLASS (UNIT # 8)

Pilot Superone Mathematics 354 Class 9th

EXERCISE 8.2

Q.1 Draw the conversion graph between litres and gallons using the relation 9 litres = 2 gallons (approximately), and taking litres along horizontal axis and gallons along vertical axis. From the graph, read

- (i) The number of gallons in 18 litres.
- (ii) The number of litres in 8 gallons.

Solution: (Litre) 9 = 2 (gallon)


$$y = \frac{2}{9}x$$

Value table for (x, y)

x =	3	4.5	21	27
y =	67	1	4.67	6

From graph

18 (litre) = 4 (gallon)
 8 (gallon) = 26 (litre)



Scale on y-axis 2 sq. = 1

MATHEMATICS FOR 9TH CLASS (UNIT # 8)

Pilot Superone Mathematics 355 Class 9th

Scale on x-axis 1 sq = 3

Q.2 On 15.03.2008 the exchange rate of Pakistani currency and Saudi Riyal was as under:

1 S. Riyal = 16.70 Rupees

If Pakistani currency y is an expression of S. Riyal x , expressed under the rule $y = 16.70x$, then draw the conversion graph between these two currencies by taking S. Riyal along x-axis.

Solution:-

$y = 16.70x$ where y is Pak. Rupee and x is S. Riyal

Table value for x and y


$x =$	1	2	3	4
$y =$	16.70	33.40	50.10	66.8

From graph

$$y = 16.70 \times 2 = 33.40$$

$$y = 16.70 \times 3 = 50.10$$

$$y = 16.70 \times 4 = 66.8$$



MATHEMATICS FOR 9TH CLASS (UNIT # 8)

Pilot Superone Mathematics 356 Class 9th

Graph. y 16.70

Q.3 Sketch the graph of each of the following lines.

- (a) $x - 3y + 2 = 0$ (b) $3x - 2y - 1 = 0$
 (c) $2y - x + 2 = 0$ (d) $y - 2x = 0$
 (e) $3y - 1 = 0$ (f) $y + 3x = 0$
 (g) $2x + 6 = 0$

3(a) $x - 3y + 2 = 0$

Solution:


$$-3y = -x - 2$$

$$y = \frac{-x-2}{-3}$$

$$y = \frac{x+2}{3}$$

Table value for x and y

x =	1	2
y =	1	0



Graph: $x - 3y + 2 = 0$

MATHEMATICS FOR 9TH CLASS (UNIT # 8)

Pilot Superone Mathematics 357 Class 9th

3(b) $3x - 2y - 1 = 0$

Solution.


$$-2y = -3x + 1$$

$$y = \frac{-3x + 1}{-2}$$

$$y = \frac{3x - 1}{2}$$

Table of ordered pairs for (x, y)

$x =$	1	-1
$y =$	1	-2



Graph: $3x - 2y - 1 = 0$

3(c) $2y - x + 2 = 0$

Solution.

$$2y = x - 2$$

$$y = \frac{x - 2}{2}$$

Table of ordered pairs for (x, y)


$x =$	2	-2	0
$y =$	0	-2	-1

MATHEMATICS FOR 9TH CLASS (UNIT # 8)

Pilot Superone Mathematics

358

Class 9th



Graph: $2y - x + 2 = 0$


3(d) $y - 2x = 0$

Solution:

$y = 2x$

Table of ordered pairs for (x, y)

$x =$	1	-1
$y =$	2	-2



Graph: $y - 2x = 0$

MATHEMATICS FOR 9TH CLASS (UNIT # 8)


Pilot Superone Mathematics 359 **Class 9th**

3(c) $y - 1 = 0$

Solution. $3y - 1 = 0 \quad y = \frac{1}{3}$

Table of ordered pairs for (x, y)

x	-	1	2	3	-2	...
y =	$\frac{1}{3}$	$\frac{1}{3}$	$\frac{1}{3}$	$\frac{1}{3}$	$\frac{1}{3}$




Graph: $3y - 1 = 0$

3(f) $y + 3x = 0$

Solution $y = -3x$

Table of ordered pairs for (x, y)

x =	-1	1
y =	3	-3



Graph: $y + 3x = 0$

MATHEMATICS FOR 9TH CLASS (UNIT # 8)

Pilot Superone Mathematics 360 Class 9th


3(g) $2x + 6 = 0$

Solution: $2x = -6$

or $x = -3$

Table of ordered pairs for (x, y)

x =	-3	3	-3	-3	-3
y =		2	3	-3	...



Graph: $2x + 6 = 0$

Q.4 Draw the graph for following relations.

(i) One mile = 1.6 km

(ii) One Acre = 0.4 Hectare

(iii) $F = \frac{9}{5}C + 32$

(iv) One Rupee = $\frac{1}{86}$ \$

4(i) One mile = 1.6 km

Solution: one mile = 1.6 km

Let us show it by an equation


$$y = 1.6x$$

We prepare a table in form of ordered pairs in x, y

x =	0	2	5	4
y =	0	3.2	8.0	6.4

MATHEMATICS FOR 9TH CLASS (UNIT # 8)

Pilot Superone Mathematics 361 Class 9th



Graph: Scale 2 sq. = 1

4(ii) One Acre = 0.4 Hectare

Solution: One Acre = 0.4 Hectare

Let us show it by an equation

$$y = 0.4x$$


Table of ordered pairs in (x, y)

x =	0	1	2	4	5
y =	0	0.4	.8	1.6	2.0

Putting the ordered pair on the graph and joining them,
 we get a straight line

MATHEMATICS FOR 9TH CLASS (UNIT # 8)

Pilot Superone Mathematics 362 Class 9th



Graph: Scale 3 sq. = 1, Scale 2 sq. = 1

4(ii) $F = \frac{9}{5}C + 32$

Solution:

Table value for F and C in form of (x, y)

x	C	0	10	20	50	100
y	F	32	50	68	122	212

$$F = \frac{9}{5}C + 32$$

$$F = \frac{9}{5}(0) + 32 = 32$$

$$F = \frac{9}{5}(10) + 32 = 50$$

$$F = \frac{9}{5}(20) + 32 = 68$$


$$F = \frac{9}{5}(100) + 32 = 212$$

MATHEMATICS FOR 9TH CLASS (UNIT # 8)

Pilot Superone Mathematics

303

Class 9th




4(iv) One Rupee = $\frac{1}{86}$ \$

Solution: One rupee = $\frac{1}{86}$ \$

We show the relation with the help of an equation

$$y = \frac{1}{86}x$$

x =	0	86	172	258	344
y =	0	1	2	3	4



EXERCISE 8.3

Solve the following pair of equations in x and y graphically.

1. $x + y = 0, 2x - y + 3 = 0$
2. $x - y + 1 = 0, x - 2y = -1$
3. $2x + y = 0, x + 2y = 2$
4. $x + y - 1 = 0, x - y + 1 = 0$
5. $2x + y - 1 = 0, x = -y$

1. $x + y = 0, 2x - y + 3 = 0$

Solution:

$$\begin{aligned} x + y &= 0 \\ y &= -x \end{aligned}$$

$$\begin{aligned} 2x - y + 3 &= 0 \\ -y &= -2x - 3 \\ y &= 2x + 3 \end{aligned}$$


Table for x and y values

$x =$	4	-4
$y =$	-4	4

Table for x and y value

$x =$	2	-4
$y =$	7	-5

Plotting points on the graph paper



Point of intersection $(-1, 1)$

Sol. Set = $\{(-1, 1)\}$

MATHEMATICS FOR 9TH CLASS (UNIT # 8)

Pilot Superone Mathematics 365 Class 9th

2. $x - y + 1 = 0$, $x - 2y = -1$

Solution:

$$\begin{aligned} x - y + 1 &= 0 \\ -y &= -x - 1 \\ y &= x + 1 \end{aligned}$$

Table for x and y values


x =	4	-5
y =	5	-4

$$\begin{aligned} x - 2y &= -1 \\ -2y &= -1 - x \\ y &= \frac{-1 - x}{2} \\ y &= \frac{1 + x}{2} \end{aligned}$$

Table for x and y value

x =	5	-5
y =	3	-2

Plotting the pair of points of both the equations.



Point of intersection $(-1, 0)$

Sol. Set = $\{(-1, 0)\}$

MATHEMATICS FOR 9TH CLASS (UNIT # 8)

Pilot Superone Mathematics 366 Class 9th

3. $2x + y = 0, x + 2y = 2$

Solution:

$$\begin{aligned} 2x + y &= 0 \\ y &= -2x \end{aligned}$$

Table value for x, y


x =	2	-3
y =	-4	6

$$\begin{aligned} x + 2y &= 2 \\ 2y &= 2 - x \\ y &= \frac{2-x}{2} \end{aligned}$$

Table value for x, y

x =	4	-2
y =	-1	3

Plotting points on the graph paper



Point of intersection $\left(-\frac{2}{3}, \frac{4}{3}\right)$

Sol. Set = $\left\{\left(-\frac{2}{3}, \frac{4}{3}\right)\right\}$

MATHEMATICS FOR 9TH CLASS (UNIT # 8)

Pilot Superone Mathematics 367 Class 9th

4. $x + y - 1 = 0, x - y + 1 = 0$

Solution:

$$\begin{aligned} x + y - 1 &= 0 \\ y &= 1 - x \end{aligned}$$

Table value of x, y


x =	-4	6
y =	5	-5

$$\begin{aligned} x - y + 1 &= 0 \\ -y &= -x - 1 \\ y &= x + 1 \end{aligned}$$

Table value of x, y

x =	4	-5
y =	5	-4

Plotting points on the graph paper



Point of intersection (0, 1)

Sol. Set = {(0, 1)}

5. $2x + y - 1 = 0, x = -y$

Solution:

$$\begin{aligned} 2x + y - 1 &= 0 \\ y &= 1 - 2x \end{aligned}$$

Table values for x, y

x =	-2	3
y =	5	-5

$$\begin{aligned} x &= -y \\ y &= -x \end{aligned}$$


Table value for x, y

x =	4	-5
y =	-4	5

MATHEMATICS FOR 9TH CLASS (UNIT # 8)

Pilot Superone Mathematics 368 Class 9th

Plotting points on the graph paper



Point of intersection $(1, -1)$

Sol. Set = $\{(1, -1)\}$

Review Exercise 8

Q.1 Choose the correct answers.

- (i) If $(x - 1, y + 1) = (0, 0)$, then (x, y) is
 (a) $(1, -1)$ (b) $(-1, 1)$
 (c) $(1, 1)$ (d) $(-1, -1)$
- (ii) If $(x, 0) = (0, y)$, then (x, y) is
 (a) $(0, 1)$ (b) $(1, 0)$
 (c) $(0, 0)$ (d) $(1, 1)$
- (iii) Point $(2, -3)$ lies in quadrant
 (a) I (b) II
 (c) III (d) IV
- (iv) Point $(-3, -3)$ lies in quadrant
 (a) I (b) II

MATHEMATICS FOR 9TH CLASS (UNIT # 8)

Pilot Superone Mathematics 369 Class 9th

- (c) III (d) IV
- (v) If $y = 2x + 1$ $x = 2$ then y is
 (a) 2 (b) 3
 (c) 4 (d) 5
- (vi) Which ordered pair satisfy the equation $y = 2x$
 (a) (1, 2) (b) (2, 1)
 (c) (2, 2) (d) (0, 1)

Answers:

(i)	a	(ii)	C	(iii)	d	(iv)	a
(v)	d	(vi)	A				

Q.2 Identify the following statements as True or False.

- (i) The point $O(0, 0)$ is a vertical line.
 (i) The point $P(2, 0)$ lies on x-axis
 (ii) The graph of $x = -2$ is a vertical line.
 (iv) $3 - y = 0$ is a horizontal line
 (v) The point $Q(-1, 2)$ is in quadrant III
 (vi) The point $R(-1, -2)$ is in quadrant IV
 (vii) $y = x$ is a line on which origin lies.
 (viii) The point $P(1, 1)$ lies on the line $x + y = 0$.
 (ix) The point $S(1, -3)$ lies in quadrant III
 (x) The point $R(0, 1)$ lies on the x-axis

Answers:


(i)	F	(ii)	T	(iii)	T	(iv)	T
(v)	F	(vi)	F	(vii)	T	(viii)	F
(ix)	F	(x)	F				

MATHEMATICS FOR 9TH CLASS (UNIT # 8)

Pilot Superone Mathematics 370 Class 9th

Q.3 Draw the following points on the graph paper.

$(-3, -3), (-6, 4), (4, -5), (5, 3)$



Graph

Q.4 Draw the graph of the following equations.

(i) $x = -6$ (ii) $y = 7$ (iii) $x = \frac{5}{2}$

(iv) $y = \frac{9}{2}$ (v) $y = 4x$ (vi) $y = -2x + 1$

4(i) $x = -6$

Solution: Table value for x, y


$x =$	-6	-6	-6	-6	6	6	
$y =$		2	4	5	1	-2	..

MATHEMATICS FOR 9TH CLASS (UNIT # 8)

Pilot Superone Mathematics

371

Class 9th




Graph of $x = -6$ to parallel to y-axis and at the left side of it.

4(ii) $y = 7$

Solution: Table value for x, y

$x =$		2	3	4	-1	-3	..
$y =$	7	7	7	7	7	7	7



MATHEMATICS FOR 9TH CLASS (UNIT # 8)

Pilot Superone Mathematics 372 Class 9th


Graph of $y = 7$ is parallel to x-axis and it is box it at a distance of 7 units

4(iii) $x = \frac{5}{2}$

Solution: $x = 2.5$

Table value for x, y

x =	2.5	2.5	2.5	2.5	2.5	2.5
y =	3	4	2	-1	



Graph of $x = \frac{5}{2}$ is parallel to y-axis and is on the right of it at a distance of $\frac{5}{2}$ units.

4(iv) $y = -\frac{9}{2}$

Solution: $y = -4.5$

Table value for x, y


x	-	3	4	2	
y =	-4.5	-4.5	-4.5	-4.5	-4.5

MATHEMATICS FOR 9TH CLASS (UNIT # 8)

Pilot Superone Mathematics

373

Class 9th




Graph of $y = -\frac{9}{2}$ is parallel to x-axis and it is below at
 a distance of $-\frac{9}{2}$ units.

4(v) $y = 4x$

Solution: Table value for x, y

$x =$	0	1	-1
$y =$	0	4	-4



$y = 4x$


MATHEMATICS FOR 9TH CLASS (UNIT # 8)

Pilot Superone Mathematics 374 Class 9th

4(vi) $y = -2x + 1$

Solution: Table value for x, y

x =	2	4	-3
y =	5	-7	7




Q.5 Draw the graph of the following equation.

(i) $y = 0.62x$ (ii) $y = 2.5x$

Sol(i) $y = 0.62x$

Solution: Table value of x, y

x =	2	6	-6	0
y =	1.24	3.72	-3.72	0



Graph $y = 0.62x$


MATHEMATICS FOR 9TH CLASS (UNIT # 8)

Pilot Superstar Mathematics 375 Class 9th

5(ii) $y = 2.5x$

Solution: Table value for ,

x =	0	2	-2
y =	0	5	-5



Graph $y = 2.5x$

Q.6 Solve the following equations

(i) $x - y = 1$, $x + y = \frac{1}{2}$

(ii) $x = 3y$, $2x - 3y = -6$

(iii) $\frac{1}{3}(x + y) = \frac{1}{2}$, $\frac{1}{3}(x - y) = -1$

6(i) $x + y = \frac{1}{2}$, $x - y = 1$

Solution:

$$x + y = \frac{1}{2}$$

$$y = \frac{1}{2} - x$$

$$= \frac{1 - 2x}{2}$$

$$x - y = 1$$

$$-y = 1 - x$$

$$y = x - 1$$

Table value for x, y


Table value for x, y

x =	3	4	5
y =	2.5	4.5	4.5

x =	3	4	-3
y =	4	5	4

MATHEMATICS FOR 9TH CLASS (UNIT # 8)

Pilot Superone Mathematics 376 **Class 9th**
 Plotting the points



Point of intersection $(\frac{3}{4}, -\frac{1}{4})$

Sol Set = $\{(\frac{3}{4}, -\frac{1}{4})\}$

6(ii) $2x - 3y = -6$ $x = 3y$

Solution:

$$\begin{aligned} 2x - 3y &= -6 \\ -3y &= -6 - 2x \\ &= \frac{-6 - 2x}{-3} \\ &= \frac{2x + 6}{3} \end{aligned}$$

$$\begin{aligned} x &= 3y \\ 3y &= x \\ y &= \frac{x}{3} \end{aligned}$$

Table values for x, y

x =	0	2	-3
y =	2	10/3	0

Table value for x, y

x =	0	6	-6
y =	0	2	-2


Plotting the points

MATHEMATICS FOR 9TH CLASS (UNIT # 8)

Pilot Superone Mathematics

377

Class 9th



Point of intersection $(-6, 2)$

Common Sol. Set = $\{(-6, 2)\}$

6(ii) $\frac{1}{2}(x - y) = -1$, $\frac{1}{3}(x + y) = 2$

Solution:

$$\frac{1}{2}(x - y) = -1$$

$$x - y = -2$$

$$y = -2 - x$$

$$y = x + 2$$

$$\frac{1}{3}(x + y) = 2$$

$$x + y = 6$$

$$y = 6 - x$$

Table values for x, y

x =	4	-6	0
y =	6	4	2

Table value for x, y


x =	4	2	0
y =	2	8	6

MATHEMATICS FOR 9TH CLASS (UNIT # 8)

Pilot Superone Mathematics

378

Class 9th



Point of intersection (2, 4)

Common Sol. Set = $\{(2, 4)\}$

www.downloadclassnotes.com

**Unit
9**

**INTRODUCTION TO CO-ORDINATE
GEOMETRY**


- (i) The distance formula.
- (ii) Collinear Points.
- (iii) Mid Point Formula.

Distance Formula

(Finding distance between two points).

Let $P(x_1, y_1)$ and $Q(x_2, y_2)$ be any two points in the coordinate plane where distance between P and Q is ' d ' (a real number).

i.e. $|PQ| = d$



MATHEMATICS FOR 9TH CLASS (UNIT # 0)

Pilot Superone Mathematics 380 Class 9th

$\overline{PM} \perp X'OX$, $\overline{QN} \perp X'OX$ and $\overline{PR} \perp \overline{QN}$ therefore,
 \overline{PM} , \overline{QN} are parallel to $Y'OY$ and PR is parallel to $X'OX$.

Thus ΔPRQ is a right angled triangle and $m \angle PRQ = 90^\circ$

Now $m \overline{PR} = |x_2 - x_1|$ and $m \overline{RQ} = |y_2 - y_1|$

Using Pythagoras Theorem

$$\begin{aligned} (m \overline{PQ})^2 &= (m \overline{PR})^2 + (m \overline{RQ})^2 \\ \text{or } d^2 &= |x_2 - x_1|^2 + |y_2 - y_1|^2 \\ \text{or } d &= \sqrt{|x_2 - x_1|^2 + |y_2 - y_1|^2} \\ &= \sqrt{|x_2 - x_1|^2 + |y_2 - y_1|^2} \end{aligned}$$

Corollary:

Distance of any point (x_1, y_1) from the origin $(0, 0) = \sqrt{x_1^2 + y_1^2}$

MATHEMATICS FOR 9TH CLASS (UNIT # 9)

Pilot Superone Mathematics 381 Class 9th

EXERCISE 9.1

1. Find the distance between the following pairs of points.

(a) $A(9, 2), B(7, 2)$ (b) $A(2, -6), B(3, -6)$

(c) $A(-8, 1), B(6, 1)$ (d) $A(-4, \sqrt{2}), B(-4, -3)$

(e) $A(3, 1), B(3, 4)$ (f) $A(0, 0), B(0, -9)$

Solution:-

1(a) $A(9, 2), B(7, 2)$

$$d = \sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2} \text{ (Formula)}$$

$$|AB| = \sqrt{(7 - 9)^2 + (2 - 2)^2}$$

$$= \sqrt{(-2)^2 + 0}$$

$$= \sqrt{4} = 2$$

1(b) $A(2, -6), B(3, -6)$

$$d = \sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2} \text{ (Formula)}$$

$$|AB| = \sqrt{(3 - 2)^2 + (-6 - (-6))^2}$$

$$= \sqrt{(1)^2 + (-6 + 6)^2}$$

$$= \sqrt{1 + 0} = 1$$

1(c) $A(-8, 1), B(6, 1)$

$$d = \sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2} \text{ (Formula)}$$

$$|AB| = \sqrt{(6 - (-8))^2 + (1 - 1)^2}$$

$$= \sqrt{(6 + 8)^2 + 0}$$

$$= \sqrt{(14)^2} = 14$$

1(d) $A(-4, \sqrt{2}), B(-4, -3)$

$$d = \sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2} \text{ (Formula)}$$

$$|AB| = \sqrt{((-4) - (-4))^2 + ((-3) - \sqrt{2})^2}$$

MATHEMATICS FOR 9TH CLASS (UNIT # 0)

Pilot Superstar Mathematics 342 Class 9th

$$= \sqrt{(-4+0)^2 + (-3-\sqrt{2})^2}$$

$$= \sqrt{0 + (-1)^2(3+\sqrt{2})^2}$$

$$3+\sqrt{2}$$

1(e) $A(3, -1), B(3, -4)$

$$d = \sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2}$$

(Formula)

$$|AB| = \sqrt{(3-3)^2 + ((-4)-(-1))^2}$$

$$= \sqrt{0 + (-4+1)^2}$$

$$= \sqrt{(-3)^2} = 3$$

1(f) $A(0, 0), B(0, -5)$

$$d = \sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2}$$

(Formula)

$$|AB| = \sqrt{(0-0)^2 + (-5-0)^2}$$

$$= \sqrt{0 + (-5)^2}$$

$$= \sqrt{(-5)^2} = 5$$

2. Let P be the point on x -axis with x -coordinate ' a ' and Q be the point on y -axis with y -coordinate ' b ' as given below. Find the distance between P and Q .

(i) $a = 9, b = 7$ (ii) $a = 2, b$

(iii) $a = -8, b = 6$ (iv) $a = -2, b$


(v) $a = \sqrt{2}, b = 1$ (vi) $a = 9, b$

Solution:-

The points which are on x -axis have y -coordinate ' 0 ' and the points which are on y -axis have x coordinate ' 0 '

MATHEMATICS FOR 9TH CLASS (UNIT # 9)

Pilot Summer Assignment



2(i) $a = 9, b = 7$

Points are $P(9, 0), Q(0, 7)$

$$d = \sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2}$$

(Formula)

$$\begin{aligned} PQ &= \sqrt{(0 - 9)^2 + (7 - 0)^2} \\ &= \sqrt{(-9)^2 + (7)^2} \\ |PQ| &= \sqrt{81 + 49} \\ &= \sqrt{130} \end{aligned}$$

2(ii) $a = 2, b = 3$

Points are $P(2, 0), Q(0, 3)$

$$d = \sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2}$$

(Formula)

$$\begin{aligned} |PQ| &= \sqrt{(0 - 2)^2 + (3 - 0)^2} \\ &= \sqrt{(-2)^2 + (3)^2} \\ &= \sqrt{4 + 9} \\ &= \sqrt{13} \end{aligned}$$

2(iii) $a = -8, b = 6$

Points are $P(-8, 0), Q(0, 6)$

$$d = \sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2}$$

(Formula)

$$|PQ| = \sqrt{(0 - (-8))^2 + (6 - 0)^2}$$

MATHEMATICS FOR 9TH CLASS (UNIT # 0)

Pilot Superior Mathematics 384 Class 9th

$$\begin{aligned} &= \sqrt{(8)^2 + (6)^2} \\ &= \sqrt{64 + 36} \\ &= \sqrt{100} = 10 \end{aligned}$$

2(iv) $a = -2, b = -3$

Points are $P(-2, 0)$ $Q(0, 3)$

$$d = \sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2}$$

(Formula)

$$\begin{aligned} |PQ| &= \sqrt{[0 - (-2)]^2 + [(-3) - 0]^2} \\ &= \sqrt{(2)^2 + (-3)^2} \\ &= \sqrt{4 + 9} \\ &= \sqrt{13} \end{aligned}$$

2(v) $a = \sqrt{2}, b = 1$

Points are $P(\sqrt{2}, 0), Q(0, 1)$

$$d = \sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2}$$

(Formula)

$$\begin{aligned} |PQ| &= \sqrt{(0 - \sqrt{2})^2 + (1 - 0)^2} \\ &= \sqrt{(-\sqrt{2})^2 + (1)^2} \\ &= \sqrt{2 + 1} \\ &= \sqrt{3} \end{aligned}$$

2(vi) $a = -9, b = 4$

Points are $P(-9, 0), Q(0, -4)$

$$d = \sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2}$$

(Formula)

$$\begin{aligned} |PQ| &= \sqrt{[0 - (-9)]^2 + [(-4) - 0]^2} \\ &= \sqrt{(9)^2 + (-4)^2} \\ &= \sqrt{81 + 16} \\ &= \sqrt{97} \end{aligned}$$

Collinear Points

Two or more than two points which lie on the same straight line are called collinear points w.r.t that line, otherwise they are non-collinear points.


MATHEMATICS FOR 9TH CLASS (UNIT # 9)

Pilot Superone Mathematics

185


Class 9th

• C



Points A, B lie on straight line 'l', these points are collinear with respect to line 'l'. Points A, B, C are not collinear w.r.t the line l.

We can prove the collinearity of more than two points with the help of distance formula.



We can prove by using distance formula that points P, Q, R are not collinear with respect to line 'm'.

$$|PQ| + |QR| = |PQ|$$

While points P, Q, R, S are not collinear w.r.t m

as

$$|PQ| + |QS| \neq |PS|$$

and

$$|PR| + |RS| \neq |PS|$$

Use of distance formula to different types of a triangle.

- | | |
|----------------------------|-------------------------|
| (i) Equilateral triangle | (ii) Isosceles triangle |
| (iii) Right angle triangle | (iv) Scalene triangle |

Definitions:-

(i) **Equilateral Triangle:**

Equilateral triangle has its sides equal in length

(ii) **Isosceles Triangle:**

Two of its sides are of equal in length while its third side has a different length.

(iii) **Right Angle Triangle:**

It has one of its angle is of 90°

(iv) **Scalene Triangle:**

Measures of all the three sides are different

Using distance formula to show that given four non-collinear points form a square, rectangle or a parallelogram.

MATHEMATICS FOR 9TH CLASS (UNIT # 9)

Pilot Superone Mathematics 386 Class 9th

- (i) **Square:**
It is a closed plane figure formed by four non-collinear points such that length of all sides are equal and measures of each angle is 90°
- (ii) **Rectangle.**
It is a closed plane figure formed by four non-collinear points if
(i) its opposite sides are equal in length.
(ii) angle of each vertex is of 90°
- (iii) **Parallelogram**
It is a closed plane figure formed by four non-collinear points if
(i) lengths of opposite sides are equal
(ii) its opposite sides are parallel
(iii) measure of none of the angle is 90°

MATHEMATICS FOR 9TH CLASS (UNIT # 9)


Pilo. Superone Mathematics 387 Class 9th

EXERCISE 9.2

Q.1. Show whether the points with vertices $(5, -2)$, $(5, 4)$ and $(-4, 1)$ are vertices of an equilateral or an isosceles triangle?

Solution:-

Let $P(-4, 1)$, $Q(5, 4)$, $R(5, -2)$



We find $|RP|$, $|QR|$ and $|PQ|$

$$d = \sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2}$$

(Formula)

$$\begin{aligned} |PQ| &= \sqrt{[5 - (-4)]^2 + [4 - 1]^2} \\ &= \sqrt{(5 + 4)^2 + (3)^2} = \sqrt{81 + 9} = \end{aligned}$$

$\sqrt{90}$ (i)

$Q(5, 4)$, $R(5, -2)$

$$d = \sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2}$$

(Formula)

$$\begin{aligned} |QR| &= \sqrt{(5 - 5)^2 + (-2 - 4)^2} \\ &= \sqrt{0 + (-6)^2} = \sqrt{36} = 6 \end{aligned}$$

(ii)

$R(5, -2)$, $P(-4, 1)$

MATHEMATICS FOR 9TH CLASS (UNIT # 9)

Pilot Superong Mathematics, 388 Class 9th

$$d = \sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2}$$

(Formula)

$$\begin{aligned} |RP| &= \sqrt{(-4 - 5)^2 + [1 - (-2)]^2} \\ &= \sqrt{(-9)^2 + (1 + 2)^2} \\ &= \sqrt{81 + (3)^2} \quad \sqrt{81 + 9} = \sqrt{90} \end{aligned}$$


(iii)

$$PQ = RP \quad (\text{From i and iii})$$

Thus PQR is an isosceles triangle

Q.2. Show whether or not the points with vertices $(-1, 1)$, $(5, 4)$, $(2, -2)$ and $(-4, 1)$ form a square.

Solution:-



Let $A(-4, 1)$, $B(2, -2)$, $C(5, 4)$, $D(-1, 1)$

We find

$$\begin{aligned} |AB| &= \sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2} \\ |AB| &= \sqrt{[2 - (-4)]^2 + [(-2) - (1)]^2} \\ &= \sqrt{(2 + 4)^2 + (-2 - 1)^2} \\ &= \sqrt{(6)^2 + (-3)^2} = \sqrt{36 + 9} = \sqrt{45} \end{aligned}$$

(i)

$$|BC| = \sqrt{(5 - 2)^2 + [4 - (-2)]^2}$$

MATHEMATICS FOR 9TH CLASS (UNIT # 9)

Pilot Superone Mathematics 389 Class 9th

$$= \sqrt{(3)^2 + (6)^2} = \sqrt{9 + 36} = \sqrt{45}$$

(ii)

$$\begin{aligned} |CD| &= \sqrt{[(-1) - (5)]^2 + [1 - 4]^2} \\ &= \sqrt{(-1 - 5)^2 + (-3)^2} \\ &= \sqrt{(-6)^2 + (-3)^2} = \sqrt{36 + 9} = \sqrt{45} \end{aligned}$$

$\sqrt{45}$

(iii)

$$\begin{aligned} |DA| &= \sqrt{[(-4) - (-1)]^2 + [2 - 1]^2} \\ &= \sqrt{(-4 + 1)^2 + 0} \\ &= \sqrt{(-3)^2} = \sqrt{9} = 3 \end{aligned}$$

(iv)

We find that

$$|AB| = |BC| = |CD| \quad (\text{From i, ii, iii})$$

But


$$|AB| = |BC| = |CD| \neq |DA|$$

Therefore $(-4, 1)$, $(2, -2)$, $(3, 0)$, $(-1, 1)$ do not form a square figure.

Q.3. Show whether or not the points with coordinates $(1, 3)$, $(4, 2)$ and $(-2, 6)$ are vertices of a right triangle.

MATHEMATICS FOR 9TH CLASS (UNIT # 9)

Pilot Superone Mathematics 390 Class 9th



Solution:-

Let $P(-2, 6)$, $Q(4, 2)$, $R(1, 3)$ be the points

We find: $|PQ|$, $|QR|$, $|RP|$

$$|PQ| = \sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2}$$

$$= \sqrt{(4 - (-2))^2 + (2 - 6)^2}$$

$$= \sqrt{(4 + 2)^2 + (-4)^2}$$

$$= \sqrt{(6)^2 + (-4)^2} = \sqrt{36 + 16} =$$

$\sqrt{52}$

and

$$|PQ|^2 = (\sqrt{52})^2 = 52 \quad (i)$$

$$|QR| = \sqrt{(1 - 4)^2 + (3 - 2)^2}$$

$$= \sqrt{(-3)^2 + (1)^2}$$

$$= \sqrt{9 + 1} = \sqrt{10}$$

$$|QR|^2 = (\sqrt{10})^2 = 10 \quad (ii)$$

$$|RP| = \sqrt{(-2 - 1)^2 + (6 - 3)^2}$$

$$= \sqrt{(-2 - 1)^2 + (3)^2}$$

$$= \sqrt{(-3)^2 + (3)^2} = \sqrt{9 + 9} =$$

$\sqrt{18}$

MATHEMATICS FOR 9TH CLASS (UNIT # 9)

Pilot Superone Mathematics 391 Class 9th

and $RP^2 + (\sqrt{18})^2 = 18$ (iii)
 $PQ^2 = 52$ From (iii), (ii), (i)
 $QR^2 = 10$
 $RP^2 = 18$

We see that

$$52 + 10 \neq 18$$

$$10 + 18 \neq 52$$

$$18 + 52 \neq 10$$

Therefore, points (1, 3), (-2, 6) do not form a right angle triangle

Q.4. Use the distance formula to prove whether or not the points (1, 1), (-2, -8), (4, 10) lie on a straight line?

Solution:-

Let $P(4, 10)$, $Q(-2, -8)$, $R(1, 1)$ be the points

We find

RP , QR and PQ

$$d = \sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2}$$

(Formula)

$$\begin{aligned} PQ &= \sqrt{(-2 - 4)^2 + (-8 - 10)^2} \\ &= \sqrt{(-6)^2 + (-18)^2} \\ &= \sqrt{36 + 324} = \sqrt{360} = 6\sqrt{10} \end{aligned}$$

(i)

$$d = \sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2}$$

(Formula)

$$\begin{aligned} QR &= \sqrt{(1 - (-2))^2 + (1 - (-8))^2} \\ &= \sqrt{(1 + 2)^2 + (1 + 8)^2} \\ &= \sqrt{(3)^2 + (9)^2} \\ &= \sqrt{9 + 81} = \sqrt{90} = 3\sqrt{10} \end{aligned}$$

(ii)

$$d = \sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2}$$

(Formula)

MATHEMATICS FOR 9TH CLASS (UNIT # 9)

Pilot Superone Mathematics 392 Class 9th

$$\begin{aligned} RP &= \sqrt{(4-1)^2 + (10-1)^2} \\ &= \sqrt{3^2 + 9^2} \\ &= \sqrt{9 + 81} = \sqrt{90} = 3\sqrt{10} \end{aligned}$$

(iii)

$$QR = 3\sqrt{10}$$

$$RP = 3\sqrt{10}$$

$$\begin{aligned} QR + RP &= 3\sqrt{10} + 3\sqrt{10} \\ &= \sqrt{10}(3 + 3) \\ &= 6\sqrt{10} \end{aligned}$$


$$PQ = 6\sqrt{10}$$

From (i)

$$QR + RP = PQ$$

Points are collinear.

See the graph also



Q.5. Find k , given that the points $(2, k)$ is equidistant from $(3, 7)$ and $(9, 1)$

Solution:-

MATHEMATICS FOR 9TH CLASS (UNIT # 9)

Pilot Superone Mathematics 393 Class 9th

$$= \sqrt{(2-k)^2}$$

$$|A(1, 7)| =$$

$$= C(9, 1)$$

We find Let $A(2, k), B(3, 7), C(9, 1)$
 $|AB|, |AC|$

$$d = \sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2}$$

(Formula)

$$\begin{aligned} |AB| &= \sqrt{(3-2)^2 + (7-k)^2} \\ &= \sqrt{(1)^2 + (7-k)^2} \\ &= \sqrt{1 + (7-k)^2} \quad (i) \end{aligned}$$

and $|AC| = \sqrt{(9-2)^2 + (1-k)^2}$
 $= \sqrt{(7)^2 + (1-k)^2}$
 $= \sqrt{49 + (1-k)^2} \quad (ii)$

According to the given condition

$$\begin{aligned} |AB| &= |AC| \\ \sqrt{1 + (7-k)^2} &= \sqrt{49 + (1-k)^2} \quad (\text{from i, ii}) \\ \text{Or } 1 + (7-k)^2 &= 49 + (1-k)^2 \\ 1 + 49 + k^2 - 14k &= 49 + 1 + k^2 - 2k \\ -12k &= 0 \\ k &= 0 \end{aligned}$$

Q.6. Use distance formula to verify that the points $A(0, 7), B(3, -5), C(-2, 15)$ are collinear.

Solution:- $A(0, 7), B(3, -5), C(-2, 15)$ are given

We find $|CA|, |BC|, |AB|$
 $d = \sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2}$
 (Formula)

$$\begin{aligned} |AB| &= \sqrt{(3-0)^2 + [(-5)-7]^2} \\ &= \sqrt{(3)^2 + (-12)^2} \\ &= \sqrt{9 + 144} \end{aligned}$$

MATHEMATICS FOR 9TH CLASS (UNIT # 9)

Pilot Superone Mathematics 394 Class 9th

$$= \sqrt{9 + 144} = \sqrt{153}$$

$$= \sqrt{3 \times 3 \times 17} = 3\sqrt{17}$$

(i)

$$d = \sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2} \quad \text{(Formula)}$$

$$BC_1 = \sqrt{(2 - 3)^2 + (15 - (-5))^2}$$

$$= \sqrt{(-1)^2 + (20)^2}$$

$$= \sqrt{1 + 400} = \sqrt{401}$$

$$= \sqrt{1 \times 1 \times 401} = \sqrt{401}$$

(ii)

$$d = \sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2} \quad \text{(Formula)}$$

$$CA_1 = \sqrt{(4 - (-2))^2 + (-15)^2}$$

$$= \sqrt{(6)^2 + (-15)^2}$$

$$= \sqrt{36 + 225} = \sqrt{261}$$

$$= \sqrt{9 \times 29} = 3\sqrt{29}$$

(iii)

$$AB_1 + CA_1 = 3\sqrt{17} + 2\sqrt{17} \quad \text{(from i + ii)}$$

iii)

$$= \sqrt{17} (3 + 2)$$

$$= 5\sqrt{17}$$

$$|BC| = 5\sqrt{17} \quad \text{(from ii)}$$

$$|AB| + |CA| = |BC|$$

Therefore A(0, 7), B(3, -5) and C(-2, 15) are collinear.

Q.7. Verify whether or not the points O(0, 0), A($\sqrt{3}$, 1), B($\sqrt{3}$, -1) are the vertices of an equilateral triangle.

Solution:-

We find the lengths of the sides of the triangle

$$O(0, 0), A(\sqrt{3}, 1), B(\sqrt{3}, -1)$$

MATHEMATICS FOR 9TH CLASS (UNIT # 9)

Pilot Superone Mathematics 395 Class 9th

$$d = \sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2}$$

(Formula)

We find

$$\begin{aligned} |AB| &= \sqrt{(\sqrt{3} - \sqrt{3})^2 + ((-1) - (-1))^2} \\ &= \sqrt{0 + (-2)^2} \\ &= \sqrt{4} = 2 \end{aligned} \quad (i)$$

$$d = \sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2}$$

(Formula)

$$\begin{aligned} |OB| &= \sqrt{(\sqrt{3} - 0)^2 + (-1 - 0)^2} \\ &= \sqrt{(\sqrt{3})^2 + (-1)^2} \\ &= \sqrt{3 + 1} = \sqrt{4} = 2 \end{aligned} \quad (ii)$$

$$\begin{aligned} |OA| &= \sqrt{(\sqrt{3} - 0)^2 + (1 - 0)^2} \\ &= \sqrt{(\sqrt{3})^2 + (1)^2} \\ &= \sqrt{3 + 1} = \sqrt{4} = 2 \end{aligned} \quad (iii)$$

$$|AB| = |OB| = |OA| = 2 \quad (\text{From i, ii, iii})$$

iii)

The lengths of the sides of the triangle are equal therefore, the triangle is equilateral

Q.8 Show that the points $A(-6, -5)$, $B(5, -5)$, $C(5, -8)$ and $D(-6, -8)$ are the vertices of rectangle. Find the lengths of the diagonals. Are they equal?

Solution:-

Points are $D(-6, -8)$, $C(5, -8)$, $B(5, -5)$, $A(-6, -5)$

We find $|DA|$, $|CD|$, $|BC|$, $|AB|$

$$d = \sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2}$$

(Formula)

$$|AB| = \sqrt{[(5) - (-6)]^2 + [(-5) - (-5)]^2}$$



MATHEMATICS FOR 9TH CLASS (UNIT # 9)

Pilot Superone Mathematics 396 Class 9th

$$= \sqrt{(5+6)^2 + (-5+5)^2}$$

$$= \sqrt{(11)^2 + 0} = \sqrt{11^2} = 11$$

(i)

$$d = \sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2}$$

(Formula)

$$|BC| = \sqrt{(3-5)^2 + [(8) - (-5)]^2}$$

$$= \sqrt{0 + (-8+5)^2}$$

$$= \sqrt{(-3)^2} = \sqrt{9} = 3$$

(ii)

$$d = \sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2}$$

(Formula)

$$|CD| = \sqrt{[(-6) - (5)]^2 + [(-8) - (-8)]^2}$$

$$= \sqrt{(-6-5)^2 + (-8+8)^2}$$

$$= \sqrt{(-11)^2} = \sqrt{11^2} = 11$$

(iii)

$$d = \sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2}$$

(Formula)

$$|DA| = \sqrt{[(-6) - (-6)]^2 + [(-5) - (-8)]^2}$$


$$= \sqrt{(-6+6)^2 + (-5+8)^2}$$

$$= \sqrt{0 + (3)^2} = \sqrt{3^2} = 3$$

(iv)

MATHEMATICS FOR 9TH CLASS (UNIT # 9)

Pilot Superone Mathematics 397 Class 9th



$$|AB| = |CD| \quad (\text{From i, iii})$$

$$|BC| = |DA| \quad (\text{From ii, iv})$$

Lengths of opposite sides are equal we find the lengths of the diagonals.

$$\begin{aligned} |AC| &= \sqrt{[5 - (-6)]^2 + [(-8) - (-5)]^2} \\ &= \sqrt{(5+6)^2 + (-8+5)^2} \\ &= \sqrt{(11)^2 + (-3)^2} = \sqrt{121+9} \end{aligned}$$

$$= \sqrt{130}$$

$$|AC|^2 = 130 \quad (\text{iv})$$

$$|AB|^2 = (11)^2 = 121 \quad (\text{v})$$

$$|BC|^2 = (3)^2 = 9 \quad (\text{vi})$$

$$121 + 9 = 130 \quad (\text{From iv, v and vi})$$

MATHEMATICS FOR 9TH CLASS (UNIT # 9)

Pilot Superone Mathematics 398 Class 9th

$$AB^2 + BC^2 = AC^2 = 130$$


$\triangle ABC$ is a right angled triangle $m\angle B = 90^\circ$

Similarly all the angle of ABCD are of 90° Therefore point A, B, C, D are the vertices of a rectangle

Q.9. Show that the points $M(-1, 4)$, $N(-5, 3)$, $P(1, 3)$ and $Q(5, -2)$ are the vertices of a parallelogram.

Solution:-

We plot the given points on the graph paper



We find the lengths of the sides with distance formula.

$$d = \sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2}$$

(Formula)

$$\begin{aligned} |NM| &= \sqrt{(-1 - (-5))^2 + (4 - 3)^2} \\ &= \sqrt{(-1 + 5)^2 + (1)^2} \\ &= \sqrt{(4)^2 + 1} = \sqrt{16 + 1} = \sqrt{17} \end{aligned}$$

(i)

MATHEMATICS FOR 9TH CLASS (UNIT # 9)

Pilot Superone Mathematics 399 Class 9th

$$d = \sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2}$$

(Formula)

$$\begin{aligned} PQ &= \sqrt{(5 - 1)^2 + [(-2) - (-3)]^2} \\ &= \sqrt{(4)^2 + (-2 + 3)^2} \\ &= \sqrt{16 + 1} = \sqrt{16 + 1} = \sqrt{17} \end{aligned}$$

(ii)

$$d = \sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2}$$

(Formula)

$$\begin{aligned} |MQ| &= \sqrt{[5 - (-1)]^2 + [(-2) - (-4)]^2} \\ &= \sqrt{(5 + 1)^2 + (-2 + 4)^2} \\ &= \sqrt{(6)^2 + (-2)^2} = \sqrt{36 + 4} = \end{aligned}$$

$\sqrt{40}$ (iii)

$$\begin{aligned} |NP| &= \sqrt{[1 - (-5)]^2 + [(-3) - (3)]^2} \\ &= \sqrt{(1 + 5)^2 + (-3 - 3)^2} \\ &= \sqrt{(6)^2 + (-6)^2} = \sqrt{36 + 36} = \end{aligned}$$

$\sqrt{72}$ (iv)

$$|NM| = |PQ|, |MQ| = |NP|$$

(From i, ii, iii and iv)

Lengths of opposite sides are equal

We find the length of the diagonal NQ ,

$$d = \sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2}$$

(Formula)

$$\begin{aligned} |NQ| &= \sqrt{[(-5) - (-3)]^2 + [(-2) - (3)]^2} \\ &= \sqrt{(-5 + 3)^2 + (-2 - 3)^2} \\ &= \sqrt{(-2)^2 + (-5)^2} = \sqrt{4 + 25} \\ &= \sqrt{29} \end{aligned}$$

$\sqrt{29}$

Now in ΔNPQ

$$NQ^2 = 29, NP^2 = 72, PQ^2 = 17$$

$$\text{and } NP^2 + PQ^2 = 72 + 17 = 89$$

$$\text{thus } NQ^2 \neq NP^2 + PQ^2$$

MATHEMATICS FOR 9TH CLASS (UNIT # 9)

Pilot Superone Mathematics 404 Class 9th

$$M \left(0, \frac{-5}{2} \right)$$

Q.2. The end point P of a line segment PQ is (-3, 6) and its end point is (5, 8). Find the coordinate of the end point Q.

Solution.

Let Q (x, y) be the other end point

Mid-point is R (5, 8)

One end point is P, (-3, 6)

$$M(x, y) = M \left(\frac{x_1 + x_2}{2}, \frac{y_1 + y_2}{2} \right) \text{ (Mid-point Formula)}$$

$$5 = \frac{-3 + x}{2}$$

$$10 = -3 + x$$

$$x = 13$$

$$\text{and } 8 = \frac{6 + y}{2}$$

$$16 = 6 + y$$

$$y = 10$$

thus Q (x, y) = Q (13, 10).

Q.3. Prove that mid-point of the hypotenuse of a right triangle is equidistant from its three vertices P (-2, 5), Q (1, 3) and R (-1, 0)

Solution. Given points are

R (-1, 0), Q (1, 3), P (-2, 5)

We find the lengths of the sides of the triangle.

$$d = \sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2} \text{ (Formula)}$$

$$RQ = \sqrt{[1 - (-1)]^2 + [3 - 0]^2}$$

$$= \sqrt{(1 + 1)^2 + (3)^2}$$

$$= \sqrt{(2)^2 + (3)^2}$$

$$= \sqrt{4 + 9} = \sqrt{13}$$

(i)

$$d = \sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2} \text{ (Formula)}$$

$$RP = \sqrt{[(-2) - (-1)]^2 + [5 - 0]^2}$$

$$= \sqrt{(-2 + 1)^2 + (5)^2}$$

MATHEMATICS FOR 9TH CLASS (UNIT # 9)

Pilot Superone Mathematics 405 Class 9th

$$\begin{aligned}
 &= \sqrt{(-3)^2 + (2)^2} \\
 &= \sqrt{9 + 4} = \sqrt{13} \quad \text{(ii)} \\
 RP &= \sqrt{[(-2) - (-1)]^2 + [5 - 0]^2} \\
 RP &= \sqrt{(-2 + 1)^2 + (5)^2} \\
 &= \sqrt{(-1)^2 + (5)^2} \\
 &= \sqrt{1 + 25} = \sqrt{26} \quad \text{(iii)} \\
 |QP|^2 + |RQ|^2 &= 13 + 13 \quad \text{(From i, ii)} \\
 &= 26 \quad \text{(iv)} \\
 |RP|^2 &= 26 \quad \text{(v)} \\
 |RP|^2 &= |QP|^2 + |RQ|^2 \quad \text{(From iv, v)}
 \end{aligned}$$

Thus hypotenuse is $|RP|$ where $R(-1, 0)$ and $P(-2, 5)$ are its end point

Let $M(x, y)$ be mid point of \overline{RP}

$$(x, y) = \left(\frac{x_1 + x_2}{2}, \frac{y_1 + y_2}{2} \right) \quad \text{(Mid-point Formula)}$$

$$\begin{aligned}
 M(xy) &= M\left(\frac{-2 + (-1)}{2}, \frac{5 + 0}{2}\right) \\
 &= M\left(\frac{-3}{2}, \frac{5}{2}\right)
 \end{aligned}$$

Now, we find distance of $M\left(\frac{-3}{2}, \frac{5}{2}\right)$ from the three vertices.

We find MP, MQ, MR

$$M\left(\frac{-3}{2}, \frac{5}{2}\right), R(-1, 0), Q(1, 3), P(-2, 5)$$

$$d = \sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2} \quad \text{(Formula)}$$

$$\begin{aligned}
 |MP| &= \sqrt{\left[(-2) - \left(\frac{-3}{2}\right)\right]^2 + \left(5 - \frac{5}{2}\right)^2} \\
 &= \sqrt{\left(-2 + \frac{3}{2}\right)^2 + \left(\frac{0 - 5}{2}\right)^2} \\
 &= \sqrt{\left(\frac{-4 + 3}{2}\right)^2 + \left(\frac{0 - 5}{2}\right)^2}
 \end{aligned}$$

MATHEMATICS FOR 9TH CLASS (UNIT # 9)

Pilot Superfine Mathematics 406 Class 9th

$$= \sqrt{\left(-\frac{1}{2}\right)^2 + \left(\frac{5}{2}\right)^2}$$

$$= \sqrt{\frac{1}{4} + \frac{25}{4}} = \sqrt{\frac{26}{4}} = \frac{1}{2}\sqrt{26} \quad (A)$$

$$d = \sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2} \quad (\text{Formula})$$

$$MQ = \sqrt{\left(1 + \frac{3}{2}\right)^2 + \left(3 - \frac{5}{2}\right)^2}$$

$$= \sqrt{\left(\frac{2+3}{2}\right)^2 + \left(\frac{6-5}{2}\right)^2}$$

$$= \sqrt{\left(\frac{5}{2}\right)^2 + \left(\frac{1}{2}\right)^2}$$

$$= \sqrt{\frac{25}{4} + \frac{1}{4}} = \sqrt{\frac{26}{4}} = \frac{1}{2}\sqrt{26} \quad (B)$$

$$d = \sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2} \quad (\text{Formula})$$

$$MR = \sqrt{\left(1 + \frac{1}{2}\right)^2 + \left(0 - \frac{5}{2}\right)^2}$$

$$= \sqrt{\left(\frac{2+1}{2}\right)^2 + \left(-\frac{5}{2}\right)^2}$$

$$= \sqrt{\frac{9}{4} + \frac{25}{4}}$$

$$= \sqrt{\frac{34}{4}} = \frac{1}{2}\sqrt{34} \quad (C)$$

$$MP = MQ = MR = \frac{1}{2}\sqrt{26} \quad (\text{From A, B, C})$$

M is equidistant from P, Q, R.

Q.4. If $O(0, 0)$, $A(3, 0)$ and $B(3, 5)$ are three points in the plane, find M_1 and M_2 as

mid-points of the line segments AB and OB respectively. Find M_1, M_2 .

Solution: $A(3, 0)$, $O(0, 0)$ and $B(3, 5)$ are the


vertices of $\triangle OAB$. M_1 is mid-point of OB .

MATHEMATICS FOR 9TH CLASS (UNIT # 9)

Pilot Superone Mathematics

407

Class 9th



$$M(x, y) = M\left(\frac{x_1 + x_2}{2}, \frac{y_1 + y_2}{2}\right) \quad (\text{Formula})$$

$$M_1\left(\frac{0 + 3}{2}, \frac{0 + 5}{2}\right)$$

$$M_1\left(\frac{3}{2}, \frac{5}{2}\right)$$

M_2 is mid-point of \overline{AB}

Therefore $M_2 = M_2\left(\frac{3 + 3}{2}, \frac{0 + 5}{2}\right)$

$$= M_2\left(\frac{6}{2}, \frac{5}{2}\right)$$

$$= M_2\left(3, \frac{5}{2}\right)$$

Now $\left(\frac{3}{2}, \frac{5}{2}\right)$ and $\left(3, \frac{5}{2}\right)$ are mid points. We find

(M_1M_2)

$$d = \sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2} \quad (\text{Formula})$$

MATHEMATICS FOR 9TH CLASS (UNIT # 9)

Pilot Superone Mathematics 408 Class 9th


$$\begin{aligned} |M_1 M_2| &= \sqrt{\left(3 - \frac{3}{2}\right)^2 + \left(\frac{5}{2} - \frac{5}{2}\right)^2} \\ &= \sqrt{\left(\frac{6-3}{2}\right)^2 + 0} \\ &= \sqrt{\left(\frac{3}{2}\right)^2} = \frac{3}{2} \end{aligned}$$

Q.5. Show that the diagonals of the parallelogram having vertices A (1, 2), B (4, 2), C (-1, -3) and D (-4, -3) bisect each other.

Solution: ABCD is a parallelogram with vertices A(1, 2), B(4, 2), C(-1, -3) and

D(-4, -3). Suppose diagonals \overline{DB} and \overline{AC} intersect at M.

A (1, 2) and C (-1, -3), we find mid-point of AC



Mid point is $\left(\frac{x_1 + x_2}{2}, \frac{y_1 + y_2}{2}\right)$ Formula

Mid-point of AC $= \left(\frac{-1+1}{2}, \frac{-3+2}{2}\right)$

MATHEMATICS FOR 9TH CLASS (UNIT # 9)

Pilot Superone Mathematics 409 Class 9th

$$\left(0, \frac{1}{2}\right) \quad (i)$$

Now B (4, 2) and D (-4, -3) are end points of the second diagonal BD.

$$\begin{aligned} \text{Mid-point of } \overline{BD} &= \left(\frac{-4+4}{2}, \frac{-3+2}{2}\right) \\ &= \left(0, \frac{-1}{2}\right) \quad (ii) \end{aligned}$$


Mid-point of diagonal \overline{AC} and \overline{BD} is the same

therefore, diagonals \overline{AC} and \overline{BD} bisect each other

Q.6. The vertices of a triangle are P (4, 6) Q (-2, -4) and R (-8, 2). Show that the length of the line segment joining the mid-points of the line segment \overline{PR} , \overline{QR} is $\frac{1}{2} \overline{PQ}$.

Solution: Vertices of $\triangle PQR$ are P(4, 6), Q(-2, -4) and R(-8, 2)

Let M_1 be mid-point of \overline{RQ}



MATHEMATICS FOR 9TH CLASS (UNIT # 9)

Pilot Superzone Mathematics 410 Class 9th

Then M_1 is

$$M_1 \left(\frac{-8+2}{2}, \frac{2-4}{2} \right) \quad \left(\text{Formula } \frac{x_1+x_2}{2}, \frac{y_1+y_2}{2} \right)$$

$$M_1 \left(\frac{-10}{2}, \frac{2}{2} \right)$$

$$M_1 (-5, 1)$$

M_2 is mid-point of RP then M_2 is

$$M_2 \left(\frac{-8+4}{2}, \frac{2+6}{2} \right)$$

$$= M_2 \left(\frac{-4}{2}, \frac{8}{2} \right)$$

$$= M_2 (-2, 4)$$

Now we use distance formula

$$d = \sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2} \quad \text{to find } |PQ|$$

$P(4, 6), Q(-2, -4)$

$$PQ = \sqrt{(-2-4)^2 + (-4-6)^2}$$

$$= \sqrt{(-6)^2 + (-10)^2}$$

$$= \sqrt{36 + 100} = \sqrt{136}$$

$$= \sqrt{2 \times 2 \times 34} = 2\sqrt{34} \quad (A)$$

and $M_2(-2, 4), M_1(-5, -1)$

Therefore,

$$M_1M_2 = \sqrt{(-2+5)^2 + (4+1)^2}$$

$$= \sqrt{(3)^2 + (5)^2}$$

$$= \sqrt{9 + 25} = \sqrt{34} \quad (B)$$

$$\therefore M_1M_2 = \frac{1}{2} PQ \quad (\text{From A and B})$$

MATHEMATICS FOR 9TH CLASS (UNIT # 9)

Pilot Superone Mathematics 411 (Class 9th)

Review Exercise 9

Q.1 Choose the correct answer

- (i) Distance between points (0, 0) and (1, 1) is
 (a) 0 (b) 1
 (c) 2 (d) $\sqrt{2}$
- (ii) Distance between the points (1, 0) and (0, 1) is
 (a) 0 (b) 1
 (c) $\sqrt{2}$ (d) 2
- (iii) Mid-point of the points (2, 2) and (0, 0) is
 (a) (1, 1) (b) (1, 0)
 (c) (0, 1) (d) (1, 4)
- (iv) Mid-point of the points (2, 2) and (-2, -2) is:
 (a) (2, 2) (b) (-2, -2)
 (c) (0, 0) (d) (1, 1)
- (v) A triangle having all sides equal is called
 (a) Isosceles (b) Scalene
 (c) Equilateral (d) None of these
- (vi) A triangle having all sides equal is called
 (a) Isosceles (b) Scalene
 (c) Equilateral (d) None of these

Answers:

- (i) d (ii) c (iii) a
 (iv) c (v) c (vi) b

2. Answer the following, which is true and which is false.

- (i) A line has two end points.
- (ii) A line segment has one end point.
- (iii) A triangle is formed by three collinear points.
- (iv) Each side of a triangle has two collinear vertices.
- (v) The end points of each side of rectangle are collinear.
- (vi) All the points that lie on the x-axis are collinear.
- (vii) Origin is the only point collinear with the points of both the axes separately.

Answers:

- (i) F (ii) F (iii) F (iv) T
 (v) T (vi) T (vii) T

MATHEMATICS FOR 9TH CLASS (UNIT # 9)

Pilot Superone Mathematics 412 Class 9th

3. Find the distance between the following pairs of points.

(i) (6, 3), (3, 3) (ii) (7, 5), (1, -1)

(iii) (0, 0), (4, 3)

3(i) Let P(6, 3) Q(3, -3)

$$d = \sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2}$$

$$|PQ| = \sqrt{(3 - 6)^2 + (-3 - 3)^2}$$

$$= \sqrt{(-3)^2 + (-6)^2}$$

$$= \sqrt{9 + 36} = \sqrt{45}$$

$$= \sqrt{3 \times 3 \times 5} = 3\sqrt{5} \text{ Units}$$

3(ii) (7, 5), (1, -1)

Let A(7, 5), B(1, -1)

$$d = \sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2}$$

$$|AB| = \sqrt{(1 - 7)^2 + (-1 - 5)^2}$$

$$= \sqrt{(-6)^2 + (-6)^2}$$

$$= \sqrt{36 + 36} = \sqrt{72}$$

$$= \sqrt{6 \times 6 \times 2} = 6\sqrt{2} \text{ Units}$$

3(iii) (0, 0), (-4, -3)

Let O(0, 0), Q(-4, -3)

$$d = \sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2}$$

$$|OQ| = \sqrt{(-4 - 0)^2 + (-3 - 0)^2}$$

$$= \sqrt{(-4)^2 + (-3)^2}$$

$$= \sqrt{16 + 9} = \sqrt{25} = 5 \text{ Units}$$

4. Find the mid-point between following pairs of points.

(i) (6, 6), (4, -2)

(ii) (5, 7), (7, -5)

(iii) (8, 0), (0, 12)

Solutions:-

4(i) Let M be mid-point of P(6, 6), (4, -2)

MATHEMATICS FOR 9TH CLASS (UNIT # 9)

Pilot Superone Mathematics 413 Class 9th

$$\text{Mid point formula } \frac{x_1 + x_2}{2}, \frac{y_1 + y_2}{2}$$

$$\text{Thus } = M\left(\frac{6+4}{2}, \frac{6-2}{2}\right)$$

$$= M\left(\frac{10}{2}, \frac{4}{2}\right)$$

= M (5, 2) is the mid-point

4(ii) Let M be mid-point of B(-7, 5), A(-5, -7)

$$\text{Mid point formula } \frac{x_1 + x_2}{2}, \frac{y_1 + y_2}{2}$$

$$\text{Thus } M\left(\frac{-7-5}{2}, \frac{5-7}{2}\right)$$

$$= M\left(\frac{-12}{2}, \frac{-2}{2}\right)$$

$$= M(-6, -1)$$

4(iii) Points P(8, 0) and Q(1, -12)

Let M be the mid-point of P(8, 0) and Q(1, -12)

$$\text{Mid point formula } \frac{x_1 + x_2}{2}, \frac{y_1 + y_2}{2}$$

$$\text{Thus mid-point M is } \left(\frac{8+1}{2}, \frac{0-12}{2}\right)$$

$$= \left(\frac{9}{2}, \frac{-12}{2}\right)$$

$$= (4.5, -6)$$

5. Define the following

- | | |
|--------------------------|---------------------------|
| (i) Co-ordinate Geometry | (ii) Collinear |
| (iii) Non-collinear | (iv) Equilateral Triangle |
| (v) Scalene Triangle | (vi) Isosceles Triangle |
| (vii) Right Triangle | (viii) Square |

MATHEMATICS FOR 9TH CLASS (UNIT # 0)

Pilot Superior Mathematics 4/4 Class 9th

Definitions

(i) Co-ordinate Geometry

Co-ordinate geometry is the study of geometrical shapes in the Cartesian plane (co-ordinate plane)

(ii) Collinear Points

Two or more than two points which lie the same straight line are called collinear points w.r.t that line

(iii) Non-collinear

Three or more than three points that do not lie on a straight line are called non-collinear points

(iv) Equilateral Triangle

A triangle having lengths of its three sides the same is called an equilateral triangle

(v) Scalene Triangle

If measures of three sides of a triangle are different then the triangle is called a scalene triangle

(vi) Isosceles Triangle

If lengths of two sides of a triangle is equal and length of the third side is different then that triangle is called an isosceles triangle.

(vii) Right Triangle

A triangle having one of its angle as a right angle (90°) is called a right angled triangle

(viii) Square

A figure formed by four non-collinear points in the plane is called a square if:

- (i) All its sides have equal lengths
- (ii) Measure of each angle is 90°

MATHEMATICS FOR 9TH CLASS (UNIT # 10)

Pilot Superone Mathematics 115

Class 9th

Unit
10


CONGRUENT TRIANGLES

Congruent Triangles:

Two triangles are said to be congruent if at least one (1) correspondence can be established between them in which the corresponding sides and angles are congruent.

For example:

If in the correspondence $\triangle ABC \leftrightarrow \triangle DEF$



(i) $\angle A \cong \angle D$

(ii) $\angle B \cong \angle E$

(iii) $\angle C \cong \angle F$

(iv) $\overline{BC} \cong \overline{EF}$

(v) $\overline{CA} \cong \overline{FD}$

(vi) $\overline{AB} \cong \overline{DE}$

Then $\triangle ABC \cong \triangle DEF$

In such a case we can also say that the correspondence $\triangle ABC \leftrightarrow \triangle DEF$ is a congruence.

Angle Side Postulate (S.A.S)

If in a given correspondence of two triangles, the two sides and the included angle of the one triangle are congruent to the corresponding two sides and the included angle of the other, the triangles are congruent.

In $\triangle LMN \leftrightarrow \triangle PQR$

If $\overline{LM} \cong \overline{PQ}$, $m\angle M = m\angle Q$, $\overline{MN} \cong \overline{QR}$

Then $\triangle LMN \cong \triangle PQR$

MATHEMATICS FOR 9TH CLASS (UNIT # 10)


Pilot Superone Mathematics

416

Class 9th

Theorem

In any correspondence of two triangles, if one side and any two angles of one triangle are congruent to the corresponding side and angles of the other, then the triangles are congruent (A.S.A \equiv A.S.A)



Given

In $\triangle ABC \leftrightarrow \triangle DEF$

$\angle B \cong \angle E$, $\overline{BC} \cong \overline{EF}$, $\angle C \cong \angle F$

To Prove

$\triangle ABC \leftrightarrow \triangle DEF$

Construction

Suppose $\overline{AB} \not\cong \overline{DE}$, take a point M on \overline{DE} such that $\overline{AB} \cong \overline{ME}$. Join M to F

Proof:

Statements		Reasons
In	$\triangle ABC \leftrightarrow \triangle MEF$	
	$\overline{AB} \cong \overline{ME}$(i)	Construction
	$\overline{BC} \cong \overline{EF}$. . . (ii)	Given
	$\angle B \cong \angle E$. . .(iii)	Given
	$\triangle ABC \cong \triangle MEF$	S.A.S postulate
So,	$\angle C \cong \angle MFE$	(Corresponding angles of congruent triangles)
But	$\angle C \cong \angle DFE$	Given
		Both congruent to $\angle C$

MATHEMATICS FOR 9TH CLASS (UNIT # 10)

Pilot Superone Mathematics

417

Class 9th

$\angle DFE \cong \angle MFE$
 This is possible only if D
 and M are the same
 points and $\overline{AB} \cong \overline{DE}$

So, $\overline{AB} \cong \overline{DE}$ (iv)
 Thus from (i), (iii) and
 (iv), we have

$$\triangle ABC \cong \triangle DEF$$


$\overline{AB} \cong \overline{ME}$ (construction)

and $\overline{ME} \cong \overline{DE}$ (proved)

S.A.S. postulate

Exercise 10.1

Q.1. In the given figure $\overline{AB} \cong \overline{CD}$ $\angle 1 \cong \angle 2$ prove that
 $\triangle ABD \cong \triangle CBE$.



Given:
 In the given figure

$$\overline{AB} \cong \overline{CD}$$

$$\angle 1 \cong \angle 2$$

To Prove: $\triangle ABC \cong \triangle CBI$

Proof:


	Statements	Reasons
In	$\triangle ABC$ and CBI	
	$\overline{AB} \cong \overline{CD}$	Given
	$\angle A \cong \angle C$	Given
And	$\angle B \cong \angle B$	Common
	$\triangle ABC \cong \triangle CBE$	S.A.S postulate

MATHEMATICS FOR 9TH CLASS (UNIT # 10)

Plot Superone Mathematics 418 Class 9th

Q.2. From a point on the bisector of an angle perpendiculars are drawn to the arms of the angle. Prove that these perpendiculars are equal in measure.

Solution:



Given:

$\angle PQR$ is given QS is bisector of $\angle PQR$ i.e. $\angle 1 = \angle 2$

(i) M is a point on QS

(ii) $MS \perp QP$, $MN \perp QR$
 i.e. $\angle 3 = \angle 4$ (each 90°)

To Prove:


Proof:

Statements	Reasons
$\angle QMS = \angle QMN$	Given
$\angle 1 = \angle 2$	Given
$\angle 3 = \angle 4$	
$QM = QM$	Common
$\triangle QMS \cong \triangle QMN$	S.A.S postulate
$MS = MN$	Corresponding sides of congruent triangles

MATHEMATICS FOR 9TH CLASS (UNIT # 10)

Pilot Superine Mathematics 419 Class 9th

Q.3 In a triangle ABC the bisectors of $\angle B$ and $\angle C$ meet in a point I . Prove that I is equidistant from the three sides of $\triangle ABC$.



Given:

In triangle ABC

- (i) BI is bisector of $\angle B$ i.e. $\angle 1 = \angle 2$
- (ii) CI is bisector of $\angle C$ i.e. $\angle 3 = \angle 4$
- (iii) I is point of intersection of BI and CI
- (iv) $IP \perp AB$, $IQ \perp BC$, $IR \perp AC$

To Prove: $IP = IQ = IR$

Proof:


Statements	Reasons
$\angle IPB \leftrightarrow \angle IQB$	
$\angle 1 = \angle 2$	Given
$\angle P = \angle Q$	Each 90°
$BI = BI$	Common
$\triangle IPB \cong \triangle IQB$	S.A.S postulate
(v) $IP = IQ$	Corresponding sides of congruent triangles
From $\triangle IRC \cong \triangle IQB$	
(vi) $IQ = IR$	From (v) and (vi)
$IP = IQ = IR$	

MATHEMATICS FOR 9TH CLASS (UNIT # 10)

Prior Superone Mathematics 420 Class 9th

Theorem

If two angles of a triangle are congruent then the sides opposite in them are also congruent



Given: In $\triangle ABC$, $\angle B \cong \angle C$

To Prove: $\overline{AB} \cong \overline{AC}$

Construction:

Draw \overrightarrow{AP} bisector of angle A i.e. $\angle 1 \cong \angle 2$, \overrightarrow{AP} cuts \overline{BC} at D


Proof:

Statements	Reasons
In $\triangle ABD \cong \triangle ACD$	
$\angle A \cong \angle B \cong \angle C$	Given
$\angle 1 \cong \angle 2$	Construction
$AD \cong \overline{AD}$	Common
$\triangle ABD \cong \triangle ACD$	S.A.S postulate
$\overline{AB} \cong \overline{AC}$	Corresponding sides of congruent triangles

MATHEMATICS FOR 9TH CLASS (UNIT # 10)

Exercise 10.2

Q.1. Prove that any two medians of an equilateral triangle are equal in measure.



Given: In $\triangle ABC$

(i) $\overline{AB} \cong \overline{BC} \cong \overline{CA}$ points P, Q, R are respectively mid points of \overline{BC} , \overline{CA} and \overline{AB} i.e: $m\overline{AR} = m\overline{BR} = \frac{1}{2} m\overline{AB}$

(ii) $m\overline{BP} = m\overline{CP} = \frac{1}{2} m\overline{BC}$
 $m\overline{CQ} = m\overline{AQ} = \frac{1}{2} m\overline{CA}$

Thus $\overline{AR} \cong \overline{BR} \cong \overline{BP} \cong \overline{CP} \cong \overline{CQ} \cong \overline{AQ}$ (From I, II)

Medians are \overline{CR} , \overline{BQ} , \overline{AP}

To Prove: $\overline{AP} \cong \overline{BQ} \cong \overline{CR}$

Proof:

Statements	Reasons
In $\triangle ABC$	Given
$\overline{AB} \cong \overline{BC} \cong \overline{CA}$	Opposite angles of equal sides
$\angle ABC \cong \angle BCA \cong \angle CAB$	
Now $\triangle BCR \leftrightarrow \triangle CBQ$	
$\overline{BC} \cong \overline{CB}$	Common


MATHEMATICS FOR 9TH CLASS (UNIT # 10)

Pilot Superone Mathematics

Class 9

$\angle ABC = \angle ACB$	Proved
1. $BR \cong CQ$	Given
$\triangle BCR \cong \triangle CBQ$	S.A.S \cong S.A.S
Thus $CR \cong BQ$	Corresponding sides of congruent triangles
Similarly $BQ \cong AP$	
$AP \cong BQ \cong CR$	

Q.2. Prove that a point, which is equidistant from the end points of a line segment is on the right bisector of the line segment.



Given:

AB is a line-segment. There is a point C that $CA = CB$

To Prove:

Point C lies on the right bisector of AB

Construction:

- Take P as mid-point of AB i.e. $AP \cong BP$
- Join point C to A, P, B

Proof:

Statements	Reasons
1. $\triangle ABC$	
2. $CA = CB$	Given
3. $\angle A \cong \angle B$	Corresponding angles of congruent \triangle

MATHEMATICS FOR 9TH CLASS (UNIT # 10)

Pilot Superone Mathematics

423

Class 9th

Now $\triangle CAP \leftrightarrow \triangle CBP$

$$\overline{CA} \cong \overline{CB}$$

Given

$$\angle A \cong \angle B$$

Proved

$$\overline{AP} \cong \overline{BP}$$

Construction

Thus $\triangle CAP \cong \triangle CBP$

S.A.S \cong S.A.S

$$(i) \quad \angle 1 \cong \angle 2$$

$$(ii) \quad m\angle 1 + m\angle 2 = 180^\circ$$

Adj. angles on one side of a line

$$\text{Thus } m\angle 1 = m\angle 2 = 90^\circ$$


Hence \overline{CP} is right

bisector of \overline{AB} and point

C lies on \overline{CP}

Theorem

In a correspondence two triangles, if three sides of one triangle are congruent to the corresponding three sides of the other, then the two triangles are congruent (S.S.S. \cong S.S.S.)



Given: In $\triangle ABC \leftrightarrow \triangle DEF$

$$\overline{AB} \cong \overline{DE}$$

$$\overline{BC} \cong \overline{EF}$$

$$\overline{CA} \cong \overline{FD}$$

MATHEMATICS FOR 9TH CLASS (UNIT # 10)

Pilot Superone Mathematics 424 Class 9th

To Prove: $\triangle ABC \cong \triangle DEF$

Construction:

Suppose the side BC is not shorter than any of the other two sides of the triangle ABC . Construct $\triangle GEF$ with G and D on opposite sides of EF such that

$GE \cong AB$ and $\angle GEF \cong \angle B$

Join G and D . Let the angles be named by the numbers as shown in the figure

Proof:

Statements	Reasons
In $\triangle ABC \leftrightarrow \triangle GEF$	
$BC \cong EF$	Given
$\angle B \cong \angle GEF$	Construction
and $AB \cong GE$	Construction
Thus $\triangle ABC \cong \triangle GEF$	S.A.S. \cong S.A.S
Now in $\triangle EGD$	
$DE \cong GE$	Each $\cong AB$
$\angle 2 \cong \angle 1$	Opposite angles of congruent sides
Thus $\angle 3 \cong \angle 4$... (i)	From (i) - (ii)
$m\angle 2 + m\angle 3 = m\angle 1 + m\angle 4$	Addition postulate of angles
or $m\angle G = m\angle D$	
i.e. $\angle D \cong \angle G$	
Now in $\triangle DGF \leftrightarrow \triangle GEF$	From (iv) and (v)
$DE \cong GE$	Each $\cong AB$
$\angle D \cong \angle G$	Proved
$DF \cong GF$	Each $\cong CA$
$\triangle DGF \cong \triangle GEF$	S.A.S. \cong S.A.S
But $\triangle ABC \cong \triangle GEF$	Transitive property of congruency
$\triangle ABC \cong \triangle DGF$	

MATHEMATICS FOR 9TH CLASS (UNIT # 10)

Pilot Superone Mathematics 425

Class 9th

Exercise 10.3

Q.1. In the figure

$$\overline{AB} \cong \overline{DC}$$


$$\overline{AD} \cong \overline{BC}$$

Prove that

$$\angle A \cong \angle C$$

And

$$\angle ABC \cong \angle ADC$$



Given:

In the given figure

$$\overline{AB} \cong \overline{DC}$$

$$\overline{AD} \cong \overline{BC}$$

To Prove:

$$\angle A \cong \angle C$$

$$\angle ABC \cong \angle ADC$$


Proof:

Statements	Reasons
In $\triangle ABD \leftrightarrow \triangle CDB$	
$\overline{AB} \cong \overline{DC}$	Given
$\overline{AD} \cong \overline{BC}$	Given
$\overline{BD} \cong \overline{BD}$	Common
$\triangle ABD \cong \triangle CDB$	S.S.S \cong S.S.S
Hence, $\angle A \cong \angle C$	Corresponding angles of congruent Δ 's
(i) $\angle 1 \cong \angle 3$	Corresponding angles of congruent Δ 's
$\angle 2 \cong \angle 4$	From (i)
$m\angle 1 + m\angle 2 = m\angle 3 + m\angle 4$	
or $m\angle ABC = m\angle ADC$	
$\angle ABC \cong \angle ADC$	

MATHEMATICS FOR 9TH CLASS (UNIT # 10)

Pilot Superone Mathematics 436 Class 9

Q.2. In the figure:



$$\overline{LN} \cong \overline{MN}$$

$$\overline{NP} \cong \overline{LP}$$

$$\angle N \cong \angle P \quad (i)$$

$$\angle NML \cong \angle PLM$$

Prove that
and

Given:

In the figure

$$\overline{LN} \cong \overline{MN}$$

$$\overline{NP} \cong \overline{LP}$$

To Prove:

$$\angle N \cong \angle P \quad (i)$$

$$\angle NML \cong \angle PLM$$


Proof:

Statements	Reasons
In $\triangle LNM$ and $\triangle LPM$	
$\overline{LN} \cong \overline{MP}$	Given
$\overline{NP} \cong \overline{LP}$	Given
$\overline{LM} \cong \overline{LM}$	Common
$\therefore \triangle LNM \cong \triangle LPM$	S.A.S \cong S.A.S
Thus $\angle N \cong \angle P$	
$\angle NML \cong \angle PLM$	

MATHEMATICS FOR 9TH CLASS (UNIT # 10)

For Supreme Mathematics 427 Class 9

Q.3. Prove that the median bisecting the base of an isosceles triangle bisects the vertex angle and it is perpendicular to the base.



Given: In $\triangle ABC$

(i) $AB = AC$

(ii) Point P is mid-point of \overline{BC} i.e. $\overline{BP} \cong \overline{CP}$

(iii) P is joined to A i.e. \overline{AP} is median
 To Prove,

$$\angle 1 \cong \angle 2$$

$$\overline{AP} \perp \overline{BC}$$

Proof:


Statements	Reasons
In $\triangle ABP \cong \triangle ACP$	
$AB \cong AC$	Given
$\overline{BP} \cong \overline{CP}$	Given
$\overline{AP} \cong \overline{AP}$	Common
$\triangle ABP \cong \triangle ACP$	SAS \cong SAS
Thus $\angle 1 \cong \angle 2$	Corresponding angles of congruent triangles
(i) $\angle 3 \cong \angle 4$	
But (ii) $m\angle 3 + m\angle 4 = 180^\circ$	Corresponding angles of congruent triangles
Thus $m\angle 3 = m\angle 4 = 90^\circ$	
therefore $\overline{AP} \perp \overline{BC}$	From (i) and (ii)

MATHEMATICS FOR 9TH CLASS (UNIT # 10)

Pilot Superone Mathematics 428

Theorem

If in the corresponding of the two right-angled triangles, the hypotenuse and one side of one triangle are congruent to the hypotenuse and the corresponding side of the other, then the triangles are congruent (H.S. ≅ H.S.)



Given: In $\triangle ABC \leftrightarrow \triangle DEF$
 $\angle ABC \cong \angle DEF$ (Right angle)
 $\overline{AC} \cong \overline{DF}$ Hypotenuse
 $\overline{BC} \cong \overline{EF}$

To Prove: $\triangle ABC \cong \triangle DEF$

Construction.

Produce \overline{AB} towards B and cut off $\overline{BC} \cong \overline{DE}$

Statements	Reasons
In $\triangle GBC$	
$m\angle CBG = 90^\circ$	Supplementary of 90°
In $\triangle GBC \leftrightarrow \triangle DEF$	
$\overline{GB} \cong \overline{DE}$	Construction
$\angle GBE \cong \angle E$	Each 90°
$\overline{BC} \cong \overline{EF}$	Given
$\triangle GBC \cong \triangle DEF$	S.A.S. \cong S.A.S.
$\overline{GC} \cong \overline{DF}$	Corresponding sides of congruent triangles
But $\overline{DF} \cong \overline{AC}$	Given
	Transitive property

MATHEMATICS FOR 9TH CLASS (UNIT # 10)

Pilot Superone Mathematics

429

Class 9th

Thus $\overline{GC} \cong \overline{AC}$

Now in $\triangle ACG$

$\angle A \cong \angle G$

Now in $\triangle ABC \leftrightarrow \triangle GBC$

$\overline{AC} \cong \overline{GC}$

$\angle A \cong \angle G$

$\angle ABC \cong \angle GBC$

Thus $\triangle ABC \cong \triangle GBC$

But $\triangle GBC \cong \triangle DEF$

Hence $\triangle ABC \cong \triangle DEF$

Corresponding angles of
congruent sides

Proved

Proved

Each 90°


S.A.A. \cong S.A.A.

Proved

Transitive property of
congruency

Exercise 10.4

Q.1. In $\triangle PAB$ of figure, $\overline{PQ} \perp \overline{AB}$ and $\overline{PA} \cong \overline{PB}$,
 prove that $\overline{AQ} \cong \overline{BQ}$ and $\angle APQ \cong \angle BPQ$



Given: In $\triangle PAB$

(i) $\overline{PA} \cong \overline{PB}$

(ii) $\overline{PQ} \perp \overline{AB}$ i.e. $m\angle 1 = m\angle 2 = 90^\circ$

To Prove: $\overline{AQ} \cong \overline{BQ}$
 $\angle APQ \cong \angle BPQ$

MATHEMATICS FOR 9TH CLASS (UNIT # 10)

Pilot Superone Mathematics

430

Class 9th

Proof:

Statements	Reasons
In $\triangle APQ \leftrightarrow \triangle BPQ$	
$\angle 1 \cong \angle 2$	Given each 90°
$\overline{PA} \cong \overline{PB}$	Given
$\overline{PQ} \cong \overline{PQ}$	Common
Thus $\triangle APQ \cong \triangle BPQ$	H.S. \cong H.S.
$\overline{AQ} \cong \overline{BQ}$	Corresponding sides of congruent triangles
$\angle APQ \cong \angle BPQ$	Corresponding angles of congruent triangles

Q.2 In the figure $m\angle C = m\angle D = 90^\circ$ and $\overline{BC} \cong \overline{AD}$

Prove that $\overline{AC} \cong \overline{BD}$ and $\angle BAC \cong \angle ABD$.



Given: In the given figure
 $m\angle C = m\angle D = 90^\circ$ (i)

$\overline{BC} \cong \overline{AD}$

To Prove $\overline{AC} \cong \overline{BD}$ (ii)
 $\angle BAC \cong \angle ABD$

Proof:

Statements	Reasons
In $\triangle ABD \leftrightarrow \triangle BAC$	
$\overline{AD} \cong \overline{BC}$	Given
$\angle D \cong \angle C$	Each 90°
$\overline{AB} \cong \overline{BA}$	Common
Thus $\triangle ABD \cong \triangle BAC$	H.S. \cong H.S.

MATHEMATICS FOR 9TH CLASS (UNIT # 10)

Pilot Superone Mathematics

131

Class 9th


$$\overline{AC} \cong \overline{BD}$$

Corresponding sides of congruent triangles.

$$\angle BAC \cong \angle ABD$$

Corresponding angles of congruent triangles

Q.3. In the figure, $m\angle B = m\angle D = 90^\circ$ and $\overline{AD} \cong \overline{BC}$.
 Prove that ABCD is a rectangle.



Given. In the figure

(i) $m\angle B = m\angle D = 90^\circ$

(ii) $\overline{AD} \cong \overline{BC}$

To Prove,

ABCD is a rectangle i.e. $\overline{AB} \parallel \overline{DC}$

Construction:

Join A to C

Proof:

Statements	Reasons
In $\triangle ABC \cong \triangle CDA$	
$\angle B \cong \angle D$	Given (each 90°)
$\overline{BC} \cong \overline{AD}$	Given
$\overline{AC} \cong \overline{AC}$	Common
$\triangle ABC \cong \triangle CDA$	HS \cong HS
Hence $\overline{AB} \cong \overline{DC}$	Corresponding sides of congruent triangles.

MATHEMATICS FOR 9TH CLASS (UNIT # 10)

Pilot Superone Mathematics 432 (Class 9th)

REVIEW EXERCISE 10

1. Which of the following are true and which are false?

- (i) A ray has two end points
- (ii) In a triangle, there can be only one right angle
- (iii) Three points are said to be collinear if they lie on same line.
- (iv) Two parallel lines intersect at a point.
- (v) Two lines can intersect only at one point
- (vi) A triangle of congruent sides has non-congruent angles.

Answers:

(i)	False	(ii)	True	(iii)	True
(iv)	False	(v)	True	(vi)	False

Q.2 If $\triangle ABC \cong \triangle LMN$




Then
 $m\angle M$
 $m\angle N$
 $m\angle A$

- ()
- (ii)
- (iii)

Answers:

- (i) $m\angle M = 30^\circ$ (ii) $m\angle N = 60^\circ$ (iii) $m\angle A = 90^\circ$

Q.3. If $\triangle ABC \cong \triangle LMN$, then find the unknown x .



MATHEMATICS FOR 9TH CLASS (UNIT # 10)

Prior Superone Mathematics 433


Class 9th

Answer:

$$m\angle N = x^\circ = 60^\circ$$

Q.4. Find the value of unknowns for the given congruent triangles.


Solution:-



$$\begin{aligned} \text{(i)} \quad 15x + 5 &= 55 \\ 15x &= 55 - 5 \\ 15x &= 50 \\ x &= \frac{50}{15} \end{aligned}$$

$$\begin{aligned} \text{(ii)} \quad 5m &= 2m + 6 \\ 5m - 2m &= 6 + 3 \\ 3m &= 9 \\ m &= 3 \end{aligned}$$

Q.5. If $\triangle PQR \cong \triangle ABC$, then find the unknowns.



Solution:-

$$\begin{aligned} \text{(i)} \quad x &= 3m \\ \text{(ii)} \quad y &= 1 \\ z &= 5 + 1 \end{aligned}$$

MATHEMATICS FOR 9TH CLASS (UNIT # 10)

Pilot Superone Mathematics 114 Class 9th

(iii) $\frac{3}{4} = \frac{6}{8}$
- 4cm

Remember

- In any correspondence of two triangles, if one side and any two angles of one triangle are congruent to the corresponding side and angles of the other, the two triangles are congruent (A.S.A \cong A.S.A)
- If two angles of a triangle are congruent, then the sides opposite to them are also congruent
- In a correspondence of two triangles, if three sides of one triangle are congruent to the corresponding three sides of the other, then the two triangles are congruent (S.S.S \cong S.S.S)
- If in the correspondence of the two right-angled triangles, the hypotenuse and one side of one triangle are congruent to the hypotenuse and the corresponding side of the other, then the triangles are congruent (H.S \cong H.S)
- Two triangles are said to be congruent, if there exists a correspondence between them such that all the corresponding sides and angles are congruent.

www.download10

MATHEMATICS FOR 9TH CLASS (UNIT # 11)

Pilot Superone Mathematics 135

Class 9th


Unit
11

PARALLELOGRAMS AND TRIANGLES

Theorem

In a parallelogram

- (i) Opposite sides are congruent
- (ii) Opposite angles are congruent
- (iii) The diagonals bisect each other



Given: ABCD is a \parallel^m that $\overline{AD} \parallel \overline{BC}$ and $\overline{AB} \parallel \overline{DC}$. Diagonals \overline{AC} and \overline{BD} cut each other at O.

- To prove
- (i) $\overline{AD} \cong \overline{BC}$, $\overline{AB} \cong \overline{DC}$
 - (ii) $\angle ABC \cong \angle ADC$
 $\angle BAD \cong \angle BCD$
 - (iii) $\overline{OA} \cong \overline{OC}$
 $\overline{OB} \cong \overline{OD}$

Proof

Statements	Reasons
$\triangle ABD \leftrightarrow \triangle CDB$	
$\angle 2 \cong \angle 1$	Alternate angles
$BD \cong BD$	Common
$\angle 4 \cong \angle 3$	Alternate angles
$\triangle ABD \cong \triangle CDB$	A.S.A \cong A.S.A
	Corresponding sides

MATHEMATICS FOR 9TH CLASS (UNIT # 11)

Pilot Superlong Mathematics	436	Class 9 th
$\overline{AB} \cong \overline{DC}$	of congruent triangles	
and $\overline{AD} \cong \overline{BC}$	proved	
Now $\angle 2 \cong \angle 1$ (i)	proved	
$\angle 4 \cong \angle 3$ (ii)	from (i) + (ii)	
$m\angle 2 + m\angle 4 = m\angle 1 + m\angle 3$		
or $m\angle ADC = m\angle ABC$		
or $\angle ADC \cong \angle ABC$		
and $\angle BAD \cong \angle BCD$		
or $\triangle ABC \cong \triangle DCB$	proved	
$\overline{BC} \cong \overline{AD}$	proved	
$\angle 6 \cong \angle 5$		
$\angle 3 \cong \angle 4$		
$\triangle BCK \cong \triangle DKA$	proved	
Thus $\overline{CK} \cong \overline{AK}$	opp Vertical angles	
and $\overline{OB} \cong \overline{OA}$	Alternate angles	
	A.A.S \cong A.A.S	

MATHEMATICS FOR 9TH CLASS (UNIT # 11)

Pilot Superone Mathematics

437

Class 9th

EXERCISE 11.1

Q.1 One angle of a parallelogram is 130° . Find the measure of its remaining angles.



Given: ABCD is a parallelogram in which $m\angle DAB = 130^\circ$

To prove: To find $m\angle B$, $m\angle D$, $m\angle C$

Construction: Produce \overline{BA} towards A

Proof:

Statements	Reasons
ABCD is a parallelogram in which	
$m\angle DAB = 130^\circ$	Given
$m\angle C = 130^\circ$	Opposite angles of "
$m\angle DAP = 180^\circ - 130^\circ$	Adj. supplementary angles
$= 50^\circ$	
$m\angle B = 50^\circ$	Corresponding angle of $\angle DAP$
$m\angle D = 50^\circ$	Opposite angle of $\angle B$

Q.2. One exterior angle formed on producing one side of a parallelogram is 40° . Find the measures of its interior angles.



Given: ABCD is a parallelogram \overline{BA} is produced towards A and angle $\angle DAP = 40^\circ$

Required: $m\angle B$, $m\angle D$, $m\angle C$

MATHEMATICS FOR 9TH CLASS (UNIT # 11)

Prior Superone Mathematics Page Class 9th


$$\begin{aligned} m\angle B &= ? \\ m\angle C &= ? \\ m\angle D &= ? \end{aligned}$$

Proof:

Statements	Reasons
ABCD is a parallelogram and	
$m\angle DAP = 40^\circ$	Given
$m\angle DAP + m\angle DAB = 180^\circ$	Adj. supp. angles
$40^\circ + m\angle DAB = 180^\circ$	$m\angle DAP = 40^\circ$ given
$m\angle DAB = 180^\circ - 40^\circ$	
$m\angle DAB = 140^\circ$ (i)	
$m\angle C = 110^\circ$ (ii)	Opp. Angles of \square^{th}
$m\angle B = m\angle DAP$	Corresponding angles
$m\angle B = 40^\circ$	
$m\angle D = 40^\circ$ $m\angle D = m\angle B$	Opp. Angles of \square^{th}
Thus $m\angle DAB = 140^\circ$	Proved
$m\angle B = 40^\circ$	
$m\angle C = 110^\circ$	
$m\angle D = 40^\circ$	

Theorem

If two opposite sides of a quadrilateral are congruent and parallel, it is a parallelogram



Given: In a quadrilateral ABCD, we have

(i) $AB = DC$

MATHEMATICS FOR 9TH CLASS (UNIT # 11)

Pilot Superone Mathematics 439

Class 9th

(ii) $AB \parallel DC$

Required: ABCD is a parallelogram

i.e. $\overline{AD} \parallel \overline{BC}$

$\overline{AD} \parallel \overline{BC}$

Construction Join B to D

Proof

Statements	Reasons
In $\triangle ABD \leftrightarrow \triangle CDB$	
$\overline{AB} = \overline{DC}$	Given
$\angle 2 = \angle 1$	Alternate angles of lines
$BD = BD$	Common
$\triangle ABD \cong \triangle CDB$	S.A.S \cong S.A.S
∴ $\angle 4 = \angle 3$ (i)	Corresponding angles of congruent triangles
$\overline{AD} \parallel \overline{BC}$ (ii)	from (i)
and $\overline{AD} = \overline{BC}$ (iii)	Corresponding sides of congruent triangles
also $\overline{AB} \parallel \overline{DC}$	given
$ABDC$ is a \parallel^m	


MATHEMATICS FOR 9TH CLASS (UNIT # 11)

Pilot Superone Mathematics 44) Class 9th

Exercise 11.12

- Q.1. Prove that a quadrilateral is a parallelogram if its
 (a) opposite angles are congruent
 (b) diagonals bisect each other

Part (a)



Given: In quadrilateral ABCD

(i) $\angle 3 \cong \angle 1$

(ii) $\angle 2 \cong \angle 4$

To prove: ABCD is a \parallel gm

i.e. $\overline{AD} \parallel \overline{BC}$ and $\overline{AB} \parallel \overline{CD}$

Const: Produce \overline{AB} towards B and \overline{CD} towards D

Proof


Statements	Reasons
Now $\angle 3 \cong \angle 1$ (i)	Given
$\angle 5 \cong \angle 1$ (ii)	Vertical opposite angles
$\angle 3 \cong \angle 5$	From (i) and (ii)
$\therefore \overline{CQ} \parallel \overline{AD}$	
$\overline{AP} \parallel \overline{CD}$	
or $\overline{BC} \parallel \overline{AD}$	
and $\overline{AB} \parallel \overline{CD}$	
ABCD is a \parallel gm	

MATHEMATICS FOR 9TH CLASS (UNIT # 11)

Pilot Superone Mathematics 111

Class 9

Part (b)



Given: ABCD is a quadrilateral in which

\overline{AC} and \overline{BD} diagonals that bisect each at O, i.e.


$\overline{OA} \cong \overline{OC}$ and $\overline{OB} \cong \overline{OD}$

Required: Quadrilateral ABCD is parallelogram

Proof

Statements	Reasons
In $\triangle OBI \leftrightarrow \triangle ODA$	
$\overline{OC} \cong \overline{OA}$	Given
$\overline{OB} \cong \overline{OD}$	Given
$\angle 1 \cong \angle 2$	Vertical opp. angles
$\triangle OBI \cong \triangle ODA$	$S.A.S \cong S.A.S$
$\overline{AD} \cong \overline{BC}$	Corresponding sides of congruent \triangle 's
$\angle 4 \cong \angle 3$	Corresponding angles of congruent \triangle 's
$\therefore \overline{AD} \parallel \overline{BC}$	Alternate angles are congruent
and $\overline{AB} \parallel \overline{CD}$	
Then ABCD is a \parallel^m	

Q.2 Prove that a quadrilateral is a parallelogram if its opposite sides are congruent.



MATHEMATICS FOR 9TH CLASS (UNIT # 11)

Pilot Superone Mathematics 442 Class 9th

Given:

In quadrilateral ABCD

(i) $\overline{AB} = \overline{DC}$ (ii) $\overline{AD} = \overline{BC}$

To Prove


ABCD is a parallelogram i.e; $\overline{AD} \parallel \overline{BC}$, $\overline{AD} \parallel \overline{BC}$

Proof.

Statements	Reasons
In $\triangle ABD \leftrightarrow \triangle CDB$	
$\overline{AD} = \overline{BC}$	Given
$\overline{AB} = \overline{DC}$	Given
$\overline{BD} = \overline{BD}$	Common
$\triangle ABD = \triangle CDB$	S.S.S
Thus $\angle 1 = \angle 2$	Corresponding angles of congruent \triangle
and $\angle 4 = \angle 3$	Proved
Now $\angle 1 = \angle 2$	Alternate angles are congruent
(ii) $\overline{AD} \parallel \overline{BC}$	
and $\angle 4 = \angle 3$	
Thus (i) $\overline{AB} \parallel \overline{DC}$	
ABCD is a	

Theorem

The line segment joining the mid-points of two sides of a triangle, is parallel to the third side and is equal to one half of its length.



MATHEMATICS FOR 9TH CLASS (UNIT # 11)

Plot Superone Mathematics

443

Class 9th

Given:

In $\triangle ABC$, the line segment \overline{DE} , joins the mid-points of \overline{AC} and \overline{BC}

Required:

$$\overline{DE} \parallel \overline{AB}$$

$$m\overline{DE} = \frac{1}{2}(m\overline{AB})$$

Construction.

- (i) Produce \overline{DE} towards E and cut off $m\overline{EF} = m\overline{DE}$
 (ii) Join B to F

Proof:


Statements	Reasons
In $\triangle DEC \leftrightarrow \triangle FEB$	Construction
$\overline{DE} \cong \overline{EF}$	Vertical opp. Angles
$\angle 1 \cong \angle 2$	Given
$\overline{CE} \cong \overline{FB}$	S.A.S. \cong S.A.S.
$\triangle DEC \cong \triangle FEB$	Corresponding angles of congruent triangles
$\angle 3 \cong \angle 4$	Corresponding sides of congruent triangles
$\overline{CD} \cong \overline{BF}$	As $\overline{CD} \cong \overline{AD}$
$\overline{AD} \cong \overline{BF}$	As Alternate angles are congruent
$\overline{AC} \parallel \overline{BF}$	As $\overline{AD} \parallel \overline{BF}$ and $\overline{AD} = \overline{BF}$
Thus $\overline{AD} \parallel \overline{BF}$	
$\triangle ADE$ is \triangle	
$\overline{DE} \parallel \overline{AB}$	
Le: $\overline{DE} \parallel \overline{AB}$	
and $\overline{DE} \cong \overline{AB}$	
But $m\overline{DE} = \frac{1}{2}(m\overline{DE})$	
Thus $m\overline{DE} = \frac{1}{2}(m\overline{AB})$	

MATHEMATICS FOR 9TH CLASS (UNIT # 11)

Pilot Superone Mathematics 111 Class 9th

Exercise 11.3

Q.1. Prove that the line-segment joining the mid-points of the opposite sides of a quadrilateral bisect each other.



Given:

- (i) ABCD is a quadrilateral points P, Q, R, S are the mid-points of \overline{AB} , \overline{BC} , \overline{CD} and \overline{DA} respectively
- (ii) P is joined to R
- (iii) S is joined to Q

\overline{PR} and \overline{SQ} intersect each other at O

Required:

$$\overline{OS} \cong \overline{OQ}$$

$$\overline{OR} \cong \overline{OP}$$

Construction:

- (i) Join P, Q, R, S in order
- (ii) Join B to D

Proof:

Statements	Reasons
In $\triangle BDC$	
R is mid-point of \overline{DC}	Given
Q is mid-point of \overline{BC}	Given
Therefore (i) $\overline{RQ} \parallel \underline{\hspace{1cm}}$	

MATHEMATICS FOR 9TH CLASS (UNIT # 11)

Pilot Superone Mathematics

445

Class 9th

$$\text{arc } m\overline{RQ} = \frac{1}{2} (m\widehat{DB})$$

In $\rightarrow \triangle BAD$

S is mid-point of \overline{AD}

P is mid-point of \overline{AB}

$$(iii) \quad \overline{PS} \parallel \overline{DB}$$

$$(iv) \quad m\overline{PS} = \frac{1}{2} (m\widehat{DB})$$

$$\overline{PQ} \parallel \overline{PS}$$

$$\overline{PQ} \cong \overline{PS}$$

BQRS is a \parallel^m

\overline{PR} , \overline{SQ} are its diagonals

$$\text{Thus } \overline{OS} \cong \overline{OQ}$$

$$\overline{OR} \cong \overline{OP}$$

From (i), (ii) each is \parallel to


\overline{DB}

From (iii) and (iv) each

$$= \frac{1}{2} (m\widehat{DB})$$

Diagonals of a \parallel^m bisect each other

Q.2. Prove that the line-segments joining the mid-points of the opposite sides of a rectangle are the right-bisectors of each other.



Given:

(i) ABCD is a rectangle

MATHEMATICS FOR 9TH CLASS (UNIT # 11)

Pilot Superone Mathematics 4-76 Class 9th

(ii) P, Q, R, S are mid-points of \overline{AB} , \overline{BC} , \overline{CD} and \overline{DA} respectively.

(iii) \overline{RP} and \overline{SQ} cut other at O
 Required.

$$\overline{OP} \cong \overline{OR}$$

$$\overline{OS} \cong \overline{OQ}$$

and $\overline{RP} \cong \overline{SQ}$

Construction:

(i) Join P, Q, R, S in order

(ii) Join A to C

Proof:

Statements	Reasons
1. $\triangle ABC$	
P is mid-point of \overline{AB}	Given
Q is mid-point of \overline{BC}	Given
(i) $\therefore \overline{PQ} \parallel \overline{AC}$	
(ii) $m\overline{PQ} = \frac{1}{2} m\overline{AC}$	
Similarly, In $\triangle ADC$	
(iii) $\overline{SR} \parallel \overline{AC}$	
(iv) $m\overline{SR} = \frac{1}{2} m\overline{AC}$	
$\overline{PQ} \parallel \overline{SR}$	From (i), (iii) each $\parallel \overline{AC}$
$m\overline{PQ} \parallel m\overline{SR}$	Each = $\frac{1}{2} m\overline{DB}$
By joining B to D, we can prove	
$\overline{RQ} \parallel \overline{SP}$	
$m\overline{RQ} \parallel m\overline{SP}$	Each = $\frac{1}{2} m\overline{AC}$

MATHEMATICS FOR 9TH CLASS (UNIT # 11)

Pilot Superone Mathematics

447

Class 9

$$m\overline{AC} \parallel m\overline{DB}$$

Diagonals of rect angle

PQRS is a \square

All its sides are equal

Now \overline{PR} and \overline{SQ} are
diagonals of \square

Now $\overline{OP} \cong \overline{OR}$

$$\overline{OS} \cong \overline{OQ}$$

In $\triangle OQR \leftrightarrow \triangle OQP$

$$\overline{OR} \cong \overline{OP}$$

Proved

$$\overline{OQ} \cong \overline{OQ}$$

Common

$$\overline{RQ} \cong \overline{PQ}$$

Proved

$$\triangle OQR \cong \triangle OQP$$

Thus (M) $\angle ROQ \cong \angle POQ$

(N) $m\angle ROQ + m\angle POQ$
 $= 180^\circ$


Adj. supp. angles.

Thus $m\angle ROQ = m\angle POQ$
 $= 90^\circ$

From M. N

Thus $\overline{RP} \perp \overline{SQ}$

Q.3. Prove that the line-segment passing through the mid-point of one side and parallel to another side of a triangle also bisects the third side.



MATHEMATICS FOR 9TH CLASS (UNIT # 11)

Pilot Superone Mathematics 448 Class 9th

Given: (i) $\triangle ABC$ point Q is mid-point of AC

i.e. $\overline{AQ} \cong \overline{CQ}$

(ii) $\overline{QR} \parallel \overline{BC}$ which cuts \overline{AB} at R.

Required: (i) $\overline{RA} \cong \overline{BR}$

Construction: (i) Take P as mid-point of BC

(ii) Join R to Q.

Proof:

Statements	Reasons
Now $\overline{RQ} \parallel \overline{BC}$	Given
$\therefore m\angle RQ = \frac{1}{2}(m\angle BC)$	
$m\angle RQ = m\angle PC$	P is mid point of BC
In $\triangle AQR \leftrightarrow \triangle QCP$	given
$\overline{AQ} \cong \overline{QC}$	Corresponding angles
$\angle 1 \cong \angle 2$	Proved
$\therefore \overline{RQ} \cong \overline{PC}$	S.A.S \cong S.A.S
$\triangle AQR \cong \triangle QCP$	
$\overline{AR} \cong \overline{QP}$	construction
(M) $\overline{BP} \cong \overline{PC}$	proved
(N) But $\overline{RQ} \cong \overline{PC}$	From M, N
$\overline{RQ} \cong \overline{BP}$	Given,
$\overline{RQ} \parallel \overline{BP}$	
$\therefore \overline{BPQR} \parallel$	
(I) $\overline{RB} \cong \overline{QP}$	Proved
(II) $\overline{AR} \cong \overline{QP}$	
Thus $\overline{AR} \cong \overline{RB}$	from (i) and (ii)
or $\overline{RA} \cong \overline{BR}$	

MATHEMATICS FOR 9TH CLASS (UNIT # 11)


Pilot Superone Mathematics

449

Class 9th

Theorem

The medians of a triangle are concurrent and their point of concurrency is the point of trisection of each median.



Given:

In $\triangle ABC$ \overline{BE} and \overline{CF} are medians that intersect at point G.

- (i) All the three medians of triangle ABC are concurrent at point G.
- (ii) Point G is the point of trisection of each medians.

Required:

- (i) Join A to G and produce ahead. It cuts \overline{BC} at D.
- Take $\overline{GH} \cong \overline{AG}$
- (ii) Join H to B and C

Proof:

Statements	Reasons
In $\triangle ABH$	
(i) F is mid-point of \overline{AB}	Given

MATHEMATICS FOR 9TH CLASS (UNIT # 11)

Pilot Superior Mathematics

450

Class 9th

(ii) G is mid-point of \overline{AF}

$$\overline{FG} \parallel \overline{EH}$$

or $\overline{GC} \parallel \overline{EH}$

Similarly $\overline{GB} \parallel \overline{HC}$

Thus BHCG is a \overline{BC} and

\overline{GH} are its diagonals

Hence

(M) Thus $\overline{BD} \cong \overline{CD}$

(N) and $\overline{GD} \cong \overline{HD}$

Point D is mid-point of \overline{BC}

We can say \overline{AD} is third median

It passes through G

$$\overline{AD} \cong \overline{GF}$$

Are concurrent

Now $m\angle G = m\angle GH$

$$m\angle AG = 2m\angle GD$$

$$\frac{m\angle AG}{m\angle GD} = \frac{2}{1}$$

$$\frac{AG}{GD} = \frac{2}{1}$$

Point G is point of trisection

Similarly it can be proved that G is point of trisection

of \overline{CF} and \overline{BE} is called centroid of the triangle ABC

Construction

Construction

$$\overline{GD} \cong \overline{DH}$$

MATHEMATICS FOR 9TH CLASS (UNIT # 11)


Pilot Superone Mathematics 451

Class 9th

Exercise 11.14

Q.1. The distances of the point of concurrency of the medians of a triangle from its vertices are respectively 1.2 cm, 1.4 cm and 1.6 cm. Find the lengths of the medians.

Solution:-




Let \overline{AP} , \overline{BQ} , \overline{CR} be medians of $\triangle ABC$

$$\text{Length of } \overline{AP} = 1.6 + \frac{1.6}{2} = 1.6 + 0.8 = 2.4 \text{ cm}$$

$$\text{Length of } \overline{BQ} = 1.2 + \frac{1.2}{2} = 1.2 + 0.6 = 1.8 \text{ cm}$$

$$\text{Length of } \overline{CR} = 1.4 + \frac{1.4}{2} = 1.4 + 0.7 = 2.1 \text{ cm}$$

Q.2. Prove that the point of concurrency of the medians of a triangle and the triangle which is made by joining the mid-points of the sides is the same.



MATHEMATICS FOR 9TH CLASS (UNIT # 11)

Pilot Superone Mathematics 452 Class 9th

Given:

$\triangle ABC$ is a triangle \overline{AP} , \overline{BQ} , \overline{CR} are its medians that are concurrent at point G . $\overline{AR} \cong \overline{BR}$, $\overline{BP} \cong \overline{CP}$;

$\overline{CQ} \cong \overline{AQ}$ points P , Q and R are joined together. $\triangle PQR$ is a triangle formed by joining the mid-point of the sides of $\triangle ABC$

Required:


\overline{RF} , \overline{PD} , \overline{QE} are the medians of triangle PQR and these pass through G also

Proof:

Statements	Reasons
In $\triangle ABC$	
$\overline{QP} \parallel \overline{AB}$	As P , Q are mid-point of \overline{AC} , \overline{BC}
or $\overline{QF} \parallel \overline{AB}$	
or $\overline{QF} \parallel \overline{AR}$	
Thus F is mid-point of \overline{AP}	
Similarly	
E is mid-point of \overline{PR} and D is mid-point of \overline{QR}	
Therefore, \overline{RF} , \overline{PD} and \overline{QE} are medians of triangle PQR and these pass through G	
Hence the result	

MATHEMATICS FOR 9TH CLASS (UNIT # 11)

Plus Superior Mathematics 453 Class 9th
Theorem
 If three or more parallel lines make congruent segments on a transversal then also intercept segments on any other line that cuts them.



- Given
 (iii) $l \parallel m \parallel n$ and transversal t cuts these parallel lines at A, B, C respectively that $\overline{AB} \cong \overline{BC}$
 (iv) Another transversal p cuts these lines at D, E, F respectively

Gi

Required:

$\overline{DE} \cong \overline{EF}$

Require

Construction:
 Passing through F and E take line segment FH, EG parallel to t .

Solution.

Ans.
 \overline{AB}

Transversal t

L, M, N

as $\overline{AB} \cong \overline{BC}$

$\overline{LM} \cong \overline{MN}$

when as

Therefore $\overline{LM} \cong \overline{MN}$

Proof	Statements	Reasons
	AGEB is a \parallel^m	$\overline{AB} \parallel \overline{GE}$ and $\overline{AG} \parallel \overline{BE}$

$$\therefore m\overline{PQ} = 1cm$$

MATHEMATICS FOR 9TH CLASS (UNIT # 11)

Pilot Superone Mathematics

$$GF \cong AB$$

Similarly $HF \cong BF$

454

Opp. Ang.

Pilot Superone Mathematics

456

Class 9th

Thus

$$m\angle N = m\angle M + m\angle MN$$


$$= 1 + 1 = 2c$$

$$m\angle Q = m\angle M + m\angle MN + m\angle NP + m\angle PQ$$

$$= 1 + 1 + 1 + 1$$

$$= 4c$$

Q.2 Take a line segment of length 5.5 cm and divide it into five congruent parts.



Construction: (i) Take a line segment 5.5 cm

(ii) Draw an acute angle $\angle BAX$

(iii) On AX take $AP \cong PQ \cong QR \cong RS \cong ST$

(iv) Join T to B

(v) Draw line parallel to TB from the points P, Q, R and S.

Result: Line segment AB is divided into congruent line

segments $AC \cong CD \cong DE \cong EF \cong FB$

Proof: $AP \cong PQ \cong QR \cong RS \cong ST$

and $TB \parallel SF \parallel RE \parallel QD \parallel PC$

Therefore: $AC \cong CD \cong DE \cong EF \cong FB$

MATHEMATICS FOR 9TH CLASS (UNIT # 11)

Pilot Super One Mathematics

457

Class 9th

REVIEW EXERCISE 11


1. Fill in the blanks.

- (i) In a parallelogram opposite sides are
- (ii) In a parallelogram opposite angles are
- (iii) Diagonals of a parallelogram each other at a point
- (iv) Medians of a triangle are
- (v) Diagonal of a parallelogram divides the parallelogram into two triangles

Answers

- | | |
|---------------|-----------------|
| (i) congruent | (ii) congruent |
| (iii) bisect | (iv) concurrent |
| (v) congruent | |

Q.2. In the parallelogram ABCD



- (i) $m\overline{AB} \dots m\overline{DC}$ (i)
- (ii) $m\overline{BC} \dots m\overline{AD}$ (ii)
- (iii) $m\angle 1 \cong$ (iii)
- (iv) $m\angle 2 \cong$ (iv)


Answers

- (i) $m\overline{AB} = m\overline{DC}$
- (ii) $m\overline{BC} = m\overline{AD}$
- (iii) $m\angle 1 = m\angle 3$
- (iv) $m\angle 2 = m\angle 4$

MATHEMATICS FOR 9TH CLASS (UNIT # 11)

Pilot Superone Mathematics 458 Class 9th

Q.3 Find the unknowns in the given figure.



Solution: $m\angle C = m\angle A$

$$n^\circ = 75^\circ \quad (i)$$

$$y^\circ = n^\circ \quad (\text{Alternate angle})^\circ$$

$$y^\circ = 75^\circ \quad (ii)$$

$$x^\circ + y^\circ = 180^\circ \quad \text{Adj. Supp. angle}$$

$$x^\circ + 75^\circ = 180^\circ$$


$$x^\circ = 180^\circ - 75^\circ$$

$$x^\circ = 105^\circ \quad (iii)$$

$$m^\circ = x^\circ$$

$$m^\circ = 105^\circ \quad (iv)$$

Q.4 If the given figure ABCD is a parallelogram, then find x, m



Solution: $11x^\circ = 55^\circ$ opp. angles of "

$$x = \frac{55}{11} = 5 \quad (i)$$

$$\text{Now } (5m + 10) + 55 = 180$$

MATHEMATICS FOR 9TH CLASS (UNIT # 11)

Pilot Superone Mathematics 459 Class 9th


$$5m + 10 + 55 = 180$$

$$5m = 180 - 10 - 55$$

$$5m = 115$$

$$m = \frac{115}{5} = 23 \quad (ii)$$

Q.5. The given figure LMNP is a parallelogram.
 Find the value of m , n .



Solution: $4m + n = 10$ opposite of $\angle L$ (i)

$$8m - 4n = 8 \quad (ii)$$

$$16m + 4n = 40$$

$$8m - 4n = 8 \quad \text{multiplying by 4}$$

$$24m = 48 \quad (\text{Adding})$$

$$m = \frac{48}{24} = 2$$

Putting $m = 2$

$$4(2) + n = 10$$


$$8 + n = 10$$

$$n = 10 - 8 = 2$$

MATHEMATICS FOR 9TH CLASS (UNIT # 11)

Pilot Superionic Mathematics 400 Class 9th

Q.6. In the question 5, sum of the opposite angles of the parallelogram is 110° , find the remaining angles.



Solution $m\angle P + m\angle L = 180^\circ$

$$m\angle P + 55 = 180$$

$$m\angle P = 180 - 55$$

$$m\angle P = 125^\circ$$

$$m\angle M = m\angle P$$

$$m\angle M = 125^\circ$$

MATHEMATICS FOR 9TH CLASS (UNIT # 12)

Pilot Super Line Mathematics 461


Class 9

**Unit
12**

LINE BISECTORS AND ANGLE BISECTORS

Theorem

Any point on the right bisector of a line segment is equidistant from its end points.



- Given:** (i) \overline{AB} is a line segment
 (ii) \overline{QT} is a right bisector of \overline{AB} i.e. $AP \cong BP$
 $\overline{QT} \perp \overline{AB}$
 $m\angle 1 = m\angle 2 = 90^\circ$
 (iii) Point C lies on \overline{QT}
 (iv) C is joined to A and B

Required: $\overline{CA} \cong \overline{CB}$


Proof:	Statements	Reasons
	In $\triangle CAP \leftrightarrow \triangle CBP$	
	$\angle 1 \cong \angle 2$	Given
	$\overline{CP} \cong \overline{CP}$	Common
	$\overline{AP} \cong \overline{BP}$	Given
So	$\triangle CAP \cong \triangle CBP$	SAS \cong SAS
	$\overline{CA} \cong \overline{CB}$	Corresponding sides of congruent Δ s

MATHEMATICS FOR 9TH CLASS (UNIT # 12)

Pilot Superone Mathematics 462 Class 9th

Converse of Theorem

Any point equidistant from the end points of a line segment is on the right bisector of it.



Given: AB is a line segment
 Point P is such that $\overline{PA} \cong \overline{PB}$
Required: Point P is on the right bisector of AB
Construction: Take C mid point of \overline{AB}
 $\overline{AC} \cong \overline{BC}$ and point P to C

Proof

Statements	Reasons
In $\triangle ACP$	Construction
$\overline{CA} \cong \overline{BC}$	Given
$\overline{PA} \cong \overline{PB}$	Common
$\overline{PC} \cong \overline{PC}$	S.S.S \cong S.S.S
So $\triangle ACP \cong \triangle BCP$ (i)	Corresponding angle of congruent \triangle s
$m\angle 1 = m\angle 2$ (ii)	Adj. Supp. Angles
$m\angle 1 + m\angle 2 = 180^\circ$	$m\angle 2 = m\angle 1$
$m\angle 1 + m\angle 2 = 180^\circ$	
$2m\angle 1 = 180$	
$m\angle 1 = 90^\circ$	
$\overline{PC} \perp \overline{AB}$	


Thus $\overline{PC} \perp \overline{AB}$ and point C on it.

MATHEMATICS FOR 9TH CLASS (UNIT # 12)

Pilot Superone Mathematics 463 Class 9

EXERCISE 12.1

Q.1 Prove that the centre of a circle is on the right bisector of each of its chords.



- Given:** (i) O is the centre of a circle
 (ii) \overline{AB} is chord of the circle

Required: Centre O lies on the right bisector of \overline{AB}

Construction: (i) Take mid-point of \overline{AB} as P.
 (ii) Join P to O O to A and B.


Proof:

Statements	Reasons
In $\triangle OPA \leftrightarrow \triangle OPB$	
$\overline{PA} \cong \overline{PB}$	Construction
$\overline{OP} \cong \overline{OP}$	Common
$\overline{OA} \cong \overline{OB}$	Radii of a circle
Therefore $\triangle OPA \cong \triangle OPB$	S.S.S \cong S.S.S
$m\angle 1 \cong m\angle 2$	Corresponding angles of congruent \triangle s
(i) $m\angle 1 = m\angle 2$	Adj Supp Angles
(ii) $m\angle 1 + m\angle 2 = 180^\circ$	From (i) and (ii)
Thus $m\angle 1 = m\angle 2 = 90^\circ$	
\overline{OP} is right bisector of \overline{AB} and O lies on it.	

MATHEMATICS FOR 9TH CLASS (UNIT # 12)

Pilot Superone Mathematics 464 Class 9th

Q.2 Where will be the centre of a circle passing through three non-collinear points? And why?



Given: A, B, C are three non-collinear points and a circle is passing through these points

Required: To find the centre of the circle and to prove that O is the centre.

Construction: (i) Join B to A and C.

(ii) Take PR right bisector of \overline{AB} .

(iii) Take QT right bisector of \overline{BC}

PR and QT intersect at point O


Proof:

Statements	Reasons
PO is right bisector of \overline{AB}	
$\overline{OA} \cong \overline{OB}$ (i)	
QO is right bisector BC	
$\overline{OB} \cong \overline{OC}$ (ii)	
$\overline{OA} \cong \overline{OB} \cong \overline{OC}$	From (i) and (ii)
Hence O is centre of the circle	

MATHEMATICS FOR 9TH CLASS (UNIT # 12)

File: Superone Mathematics 465 Class 9th

Q.3 Three villages P, Q and R are not on the same line. The people of these villages want to make a Children Park at such place which is equidistant from these three villages. After fixing the place of Children Park, prove that the park is equidistant from the three villages.



Given: P, Q, R three villages are non-collinear

Required: To locate a point for a park equidistant from these villages.

Construction: (i) Join Q to P and R.

(ii) Take \overline{AT} right bisector of \overline{PQ} .

(iii) Take \overline{BU} right bisector of \overline{QR} .

\overline{AT} and \overline{BU} intersect at point O.

Result: O is the required place

Proof:


	Reasons
$\overline{OP} \cong \overline{OQ}$ (i)	Point O is on the right bisector of \overline{PQ}
$\overline{OQ} \cong \overline{OR}$ (ii)	Point O is on the right bisector of \overline{QR}
$\overline{OP} \cong \overline{OQ} \cong \overline{OR}$	From (i) + (ii)
Hence O is equidistant from P, Q and R.	

MATHEMATICS FOR 9TH CLASS (UNIT # 12)

Pilot Superone Mathematics 466 Class 9th

Theorem

The right bisectors of the sides of a triangle are concurrent.



Given: In $\triangle ABC$
 \overline{PU} , \overline{QT} are right bisectors of \overline{AB} and \overline{BC}
 \overline{PU} and \overline{QT} intersect at I .

Required: Point I lies on the right bisector of \overline{AC}
 i.e. $IC \cong IA$.

Proof:

Statements	Reasons
$BI \cong CI$ (i)	Point I is on the right bisector of BC
$BI \cong AI$ (ii)	Point I is on the right bisector of AB
$CI \cong IA$	From (i) and (ii)
Hence Point I lies on the right bisector of AC	
Therefore three right bisectors of sides of a triangle are concurrent.	


MATHEMATICS FOR 9TH CLASS (UNIT # 12)

Hol Superone Mathematics 467

Class 9th

Theorem

Any point on the bisector of an angle is equidistant from its arms.



- Given:** (i) $\angle ABC$ is an angle.
 (ii) \overline{BM} is bisector of $\angle ABC$
 (iii) $\overline{PQ} \perp \overline{BC}$
 (iv) $\overline{PK} \perp \overline{BA}$

To prove: $\overline{PQ} \cong \overline{PK}$

Proof:

Statements	Reasons
In $\triangle PBR \longleftrightarrow \triangle PQB$	
$\overline{BP} \cong \overline{BP}$	Common
$\angle 1 \cong \angle 2$	Given
$\angle PRB \cong \angle PQB$	Each 90°
$\triangle PBR \cong \triangle PQB$	S.A.A. \cong S.A.A.
$\overline{PR} \cong \overline{PQ}$	Corresponding sides of congruent triangles.


MATHEMATICS FOR 9TH CLASS (UNIT # 12)

Pilot Superone Mathematics 468

Class 9th

Converse Theorem

Any point inside an angle, equidistant from the arms, is on the bisector of it.



- Given:** (i) A point P lies inside of angle $\angle ABC$.
 (ii) $\overline{PM} \perp \overline{BC}$, $\overline{PN} \perp \overline{BA}$
 (iii) $\overline{PM} \cong \overline{PN}$

Required: Point P is on the bisector of $\angle ABC$

Construction: Join B to P and produce.

Proof:


Statements	Reasons
In $\triangle PMB \longleftrightarrow \triangle PNB$	
$\overline{BP} \cong \overline{BP}$	Common
$\angle 1 \cong \angle 2$	Each 90° (given)
$\overline{PM} \cong \overline{PN}$	Given
$\triangle PMB \cong \triangle PNB$	H.S. \cong H.S
Thus $\angle 3 \cong \angle 4$	Corresponding sides of congruent triangles
Hence \overline{BQ} is bisector of $\angle ABC$ and point P lies on it.	

MATHEMATICS FOR 9TH CLASS (UNIT # 12)

Pilot Superone Mathematics 469 Class 9th

EXERCISE 12.2

Q.1 In a quadrilateral $ABCD$, $\overline{AB} \cong \overline{BC}$ and the right bisector of \overline{AD} , \overline{CD} meet each other at point N . Prove that \overline{BN} is bisector of $\angle ABC$.



- Given:** (i) $ABCD$ is a quadrilateral
 (ii) $\overline{AB} \cong \overline{BC}$
 (iii) \overline{PN} is right bisector of \overline{AD}
 (iv) \overline{QN} is right bisector of \overline{CD}
 \overline{PN} and \overline{QN} intersect each other at N
 (v) N is joined to B

Required: \overline{BN} is bisector of angle $\angle ABC$
 i.e. $\angle 1 \cong \angle 2$

Const: Join N to A, D, C


Proof:

Statements	Reasons
$\overline{NA} \cong \overline{ND}$ (i)	N lies on right bisector of \overline{AD}
$\overline{ND} \cong \overline{NC}$ (ii)	N lies on right bisector of \overline{CD}
$\overline{NA} \cong \overline{NC}$	From (i) and (ii)
In $\triangle NAB \longleftrightarrow \triangle NCB$	Proved
$\overline{NA} \cong \overline{NC}$	Given
$\overline{AB} \cong \overline{BC}$	Common
$\overline{BN} \cong \overline{BN}$	S.S.S \cong S.S.S
$\triangle NAB \cong \triangle NCB$	
So $\angle 1 \cong \angle 2$	
Hence \overline{BN} is bisector of $\angle ABC$	

MATHEMATICS FOR 9TH CLASS (UNIT # 12)

Plot Superone Mathematics 470 Class 9th

Q.2 The bisector of $\angle A$, $\angle B$ and $\angle C$ of a quadrilateral $ABCP$ meet each other at point O . Prove that the bisector of $\angle P$ unit also pass through the point O .



- Given:** (i) $ABCP$ is a quadrilateral
 (ii) Bisectors of $\angle A$, $\angle B$ and $\angle C$ meet at point O

Required: Point O lies on the bisector of $\angle P$

Construction: (i) Join O to P

- (ii) Draw $\overline{OL} \perp \overline{AB}$, $\overline{OM} \perp \overline{BC}$ and $\overline{ON} \perp \overline{PC}$ also
 $\overline{OQ} \perp \overline{AP}$


Proof:

Statements	Reasons
$\perp \overline{OL} \cong \perp \overline{OQ}$ (i)	Point O is on the bisector of $\angle A$
$\perp \overline{OL} \cong \overline{OM}$ (ii)	Point O is on the bisector of $\angle B$
$\perp \overline{OM} \cong \overline{ON}$ (iii)	Point O is on the bisector of $\angle C$
$\perp \overline{OQ} \cong \perp \overline{ON}$	From (i), (ii) and (iii)
Thus Point O lies on the bisector of $\angle P$	
Therefore \overline{OP} is bisector of $\angle P$	

MATHEMATICS FOR 9TH CLASS (UNIT # 12)

Pilot Superone Mathematics 471 Class 9th

Q.3 Prove that the right bisector of congruent sides of an isosceles triangle and its altitude are congruent.



Given: In $\triangle ABC$

(i) $\overline{AB} \cong \overline{AC}$

(ii) \overline{PT} is right bisector of \overline{AC}

(iii) \overline{QU} is right bisector of \overline{AB}

\overline{PT} and \overline{QU} intersect each other at O

Required: Point O lies on the altitude of $\triangle ABC$

Construction: Join A to O and produce it. It cuts \overline{BC} at D.

Proof:

Statements	Reasons
$m\overline{AB} = m\overline{AC}$	Given
$\frac{1}{2} m\overline{AB} = \frac{1}{2} m\overline{AC}$	Dividing by 2
$\overline{AP} \cong \overline{AQ}$	
In $\triangle APO \longleftrightarrow \triangle AQO$	
$\overline{AP} \cong \overline{AQ}$	proved
$\overline{AO} = \overline{AO}$	common
$\angle APO \cong \angle AQO$	each 90°
$\triangle APO \cong \triangle AQO$	H.S. \cong H.S.

MATHEMATICS FOR 9TH CLASS (UNIT # 12)

Pilot Superone Mathematics

472

Class 9th

Thus $\angle 1 \cong \angle 2$

Now in $\triangle BAD \longleftrightarrow \triangle CAD$

$\overline{AB} \cong \overline{AC}$

$\angle 1 \cong \angle 2$

$\overline{AD} = \overline{AD}$

$\triangle BAD \cong \triangle CAD$

(i) $\angle 3 \cong \angle 4$

(ii) $m\angle 3 + m\angle 4 = 180^\circ$

Thus $m\angle 3 = m\angle 4 = 90^\circ$

Therefore $\overline{AD} \perp \overline{AB}$

Point O lies on altitude \overline{AD}

Given

Proved


Common

S.A.S. \cong S.A.S.

Corresponding angles
 Congruent Δ 's

Adj supp. angles

Q.4 Prove that the altitudes of a triangle are concurrent.



Given:

In $\triangle ABC$

\overline{AD} , \overline{BE} , \overline{CF} are its altitudes.

i.e. $\overline{AD} \perp \overline{BC}$, $\overline{BE} \perp \overline{AC}$, $\overline{CF} \perp \overline{AB}$

Required:

\overline{AD} , \overline{BE} and \overline{CF} are concurrent.

Construction: Passing through A, B, C take $\overline{RQ} \parallel \overline{BC}$.

$\overline{RP} \parallel \overline{AC}$ and $\overline{QP} \parallel \overline{AB}$ respectively

Forming a triangle PQR

MATHEMATICS FOR 9TH CLASS (UNIT # 12)

Pilot Superone Mathematics 473 Class 9th

Proof.

Statements	Reasons
$BC \parallel PQ$	Construction
$AB \parallel QC$	Construction
$ABCO$ is a \square	
Hence $\overline{AO} = \overline{BC}$	
Similarly $\overline{AO} = \overline{BC}$	
Hence point A is mid-point of \overline{RQ}	
And $\overline{AD} \perp \overline{BC}$	
$\overline{BC} \parallel \overline{RQ}$	
$\overline{AD} \perp \overline{RQ}$	
Thus \overline{AD} is right bisector of \overline{RQ}	
Similarly \overline{BE} is right bisector of \overline{RP} and \overline{CF} is right bisector of \overline{PQ}	
Therefore \overline{AD} , \overline{BE} , \overline{CF} are right bisector of sides of $\triangle PQR$	Right bisector of sides of \triangle are con-current
\overline{AD} , \overline{BE} and \overline{CF} are concurrent	

MATHEMATICS FOR 9TH CLASS (UNIT # 12)


Pilot Superone Mathematics

474

Class 9th

Theorem

The bisectors of the angles of a triangle are concurrent



Given:

In $\triangle ABC$ bisectors of $\angle B$, $\angle C$ meet at I

Required:

Bisector of $\angle A$ passes through I

Construction:

(i) Join A to I

(ii) Take $IF \perp AB$

(iii) $ID \perp BC$ and

(iv) $IE \perp AC$

Proof:

Statements	Reasons
(i) $\perp IF \cong \perp ID$	I lies on bisector of angle B
(ii) $\perp IE \cong \perp ID$ $\therefore \perp IE \cong \perp IF$	I lies on bisector of angle C. From (i) and (ii)
Hence I lies on bisector of $\angle A$.	
Therefore \overline{AI} is bisector of $\angle A$.	
Thus the bisectors of angles of a triangle are concurrent.	


CONCLUSION:

MATHEMATICS FOR 9TH CLASS (UNIT # 12)

Pilot Superone Mathematics 475 Class 9th

EXERCISE 12.3

Q.1 Prove that the bisectors of the angle of base of an isosceles triangle intersect each other on its altitude



- Given:** In $\triangle ABC$
 (i) $\overline{AB} \cong \overline{AC}$
 Bisectors of $\angle B$ and $\angle C$ intersect at O
 (ii) A is joined to O and produce that cuts \overline{BC} at D .
Required: \overline{AD} is altitude of $\triangle ABC$
 i.e. $\overline{AD} \perp \overline{BC}$

Proof:

Statements	Reasons
In $\triangle ABC$ $\overline{AB} \cong \overline{AC}$ $\angle ABC \cong \angle ACB$ $\therefore \frac{1}{2} m\angle ABC = \frac{1}{2} m\angle ACB$ Thus $\angle 1 \cong \angle 3$	Given Angles opposite to congruent sides

MATHEMATICS FOR 9TH CLASS (UNIT # 12)

Pilot Superone Mathematics

476

Class 9

Now in $\triangle OBC$

$$\angle 1 \cong \angle 3$$

$$\overline{BO} \cong \overline{CO}$$

Now $\triangle AOC \longleftrightarrow \triangle AOB$

$$\overline{AO} \cong \overline{AO}$$

$$\overline{AC} \cong \overline{AB}$$

$$\overline{BO} \cong \overline{CO}$$

$$\triangle AOC \cong \triangle AOB$$

Thus $\angle 5 \cong \angle 6$

Now in $\triangle ABD \longleftrightarrow \triangle ACD$

$$\overline{AD} \cong \overline{AD}$$

$$\angle 5 \cong \angle 6$$

$$\overline{AB} \cong \overline{AC}$$

$$\triangle ABD \cong \triangle ACD$$

$$(i) \quad m\angle 7 \cong m\angle 8$$

$$(ii) \text{ point } m\angle 7 + m\angle 8 = 180^\circ$$

$$\text{Thus } m\angle 7 = m\angle 8 = 90^\circ$$

$$\therefore \overline{AD} \perp \overline{BC}$$

And \overline{AD} passes through point O

Proved

Opp Sides of congruent angles

Common

Given

Proved

S.S.S. \cong S.S.S

Common

Proved

Given

S.A.S \cong S.A.S

Corresponding angles of congruent triangles


Adj supp angles

From (i) and (i)

MATHEMATICS FOR 9TH CLASS (UNIT # 12)

Pilot Superior Mathematics 477 Class 9th

Q.2 Prove that the bisectors of two exterior and third interior angle of triangle are concurrent.



Given: In $\triangle ABC$
 $\angle ABR$ and $\angle BAP$ are two exterior angles.
 Their \overline{AP} and \overline{BQ} bisectors intersect each other at O. C is joined to O.

Required: \overline{CO} is bisector of angle $\angle C$ i.e. $\angle 1 \cong \angle 2$

Construction: $\overline{OL} \perp \overline{CR}$, $\overline{ON} \perp \overline{AB}$ and $\overline{OM} \perp \overline{CP}$

Proof:

Statements	Reasons
$\angle OLM \cong \angle ONL$ (i)	\overline{BO} is bisector of $\angle ABR$
$\angle OLM \cong \angle ONL$ (ii)	\overline{AO} is bisector of $\angle BAP$
$\angle OLM \cong \angle ONL$	From (i) and (ii)
Hence \overline{CO} is bisector of $\angle C$	
$\angle 1 \cong \angle 2$	

MATHEMATICS FOR 9TH CLASS (UNIT # 12)

Pilot Superone Mathematics 478 Class 9th

Review Exercise 12

Q.1 Which of the following are true and which are false?

- (i) Bisection means to divide into two equal parts.
- (ii) Right bisection of line segment means to draw perpendicular which passes through the mid-point.
- (iii) Any point on the right bisector of a line segment is not equidistant from its end points.
- (iv) Any point equidistant from the end points of a line segment is on the right bisector of it.
- (v) The right bisectors of the sides of a triangle are not concurrent.
- (vi) The bisectors of the angles of a triangle are concurrent.
- (vii) Any point on the bisector of an angle is not equidistant from its arms.
- (viii) Any point inside an angle, equidistant from its arms, is on the bisector.


Answers:

(i)	T	(ii)	T	(iii)	F	(iv)	T
(v)	F	(vi)	T	(vii)	F	(viii)	T

Q.2 If \overline{CD} is right bisector of line segment \overline{AB} , then

(i) $m\angle O A =$ _____

(ii) $m\angle A Q =$ _____



MATHEMATICS FOR 9TH CLASS (UNIT # 12)

Prior Superone Mathematics 479 Class 9th

Answers

$$m\overline{OA} = m\overline{OB}$$

$$m\overline{AQ} = m\overline{BQ}$$


Q.3 Define the following:

(i) Bisector of a line segment

(ii) Bisector of an angle

- Answers: (i) A line is called the bisector of a line segment if it passes through its mid-point.
 (ii) If a ray divides an angle into two congruent angles then that ray is called bisector of that angle.

Q.4 The given triangle ABC is equilateral triangle and AD is bisector of angle A, then find the values of unknowns x° , y° and z° .



Given In $\triangle ABC$

$$AB \cong BC \cong CA$$

$$\text{i.e. } \angle B \cong \angle C \cong \angle BAC \text{ each } = 60^\circ$$

$$\overline{AD} \text{ is bisector of } \angle A \text{ i.e. } x^\circ = y^\circ$$

Required: To find x° , y° and z°

Solution Proof:

$$z^\circ = 60$$

given


\overline{AD} is bisector of $\angle A$

MATHEMATICS FOR 9TH CLASS (UNIT # 12)

Pilot Supercare Mathematics 480 Class 9th

Now $m\angle A = 60^\circ$
 $\frac{1}{2} m\angle A = x^\circ = y^\circ = \frac{60}{2}$
 $x^\circ = y^\circ = 30^\circ$

Q.5 In the given congruent triangles LMO and LNO, find the unknowns x and m .



Solution: In $\triangle LMO \cong \triangle LNO$
 $MO \cong NO$
 $m = 12$ (i)
 $2x + 6 = 18$
 $2x = 18 - 6$
 $2x = 12$
 $x = \frac{12}{2} = 6$


Q.6 \overline{CD} is right bisector of the line segment \overline{AB} .

- (i) If $m\overline{AB} = 6$ cm, then find the $m\overline{AD}$ and $m\overline{BD}$.
 (ii) If $m\overline{BD} = 4$ cm, then find $m\overline{AD}$.

MATHEMATICS FOR 9TH CLASS (UNIT # 12)

Pilot Superone Mathematics 481

Class 9th



In the given figure \overline{CD} is right bisector of \overline{AB}

- i.e. $\overline{AL} \cong \overline{BL}$
 $m \overline{AB} = 6 \text{ cm}$
 $m \overline{BD} = 4 \text{ cm}$
 $m \overline{AL} = ?$
 $m \overline{LB} = ?$
 $m \overline{AD} = ?$

Proof:

Statements	Reasons
$\overline{AL} \cong \overline{LB}$ (i)	\overline{CD} is right bisector of \overline{AB}
and $m \overline{AB} = 6 \text{ cm}$	
Therefore	
$m \overline{AL} = m \overline{LB} = \frac{1}{2} m \overline{AB}$	
$\frac{1}{2} \times 6 = 3 \text{ cm}$	
$m \overline{BD} = 4 \text{ cm}$ (ii)	
$m \overline{AD} = m \overline{BD}$	Every point on right bisector is equidistant from end points
Hence $m \overline{AD} = 4 \text{ cm}$	

MATHEMATICS FOR 9TH CLASS (UNIT # 12)


Pilot Superwork Mathematics 482 Class 9th

Unit
13

SIDES AND ANGLES OF A TRIANGLE

Theorem

If two sides of a triangle are unequal in length, the longer side has an angle of greater measure opposite to it.



Given: In $\triangle ABC$
 $m\angle C > m\angle B$

To prove: $m\angle A > m\angle B$

Construction: Form \overline{CB} cut $\overline{CD} \cong \overline{CA}$
 Join D to A

Proof:

Statements	Reasons
In $\triangle ACD$ $m\angle 1 = m\angle 2$ (i)	Angles opposite to congruent sides
In $\triangle ABD$ $m\angle 2 > m\angle B$ (ii)	$\angle 2$ is exterior and $\angle B$ is interior
$m\angle 1 > m\angle B$ (iii)	By (i) and (ii)
Now $m\angle A = m\angle 1 + m\angle BAD$	Addition of angles (postulate)
$m\angle A > m\angle B$ (iv)	By (iii) and (iv)
$m\angle A > m\angle 1 > m\angle B$	Transitive property of inequality of numbers
or $m\angle A > m\angle B$	


MATHEMATICS FOR 9TH CLASS (UNIT # 12)

Prior Superone Mathematics 483

Class 9th

Converse of Theorem

If two angles of a triangle are unequal in measure, the side opposite to the greater angle is longer than the side opposite to the smaller angle.



Given: In $\triangle ABC$
 $m\angle A < m\angle B$

Required $m\overline{BC} > m\overline{AC}$

Proof:


Statements	Reasons
If $m\overline{BC} > m\overline{AC}$ Then (i) $m\overline{BC} = m\overline{AC}$	Trichotomy property of numbers.
or (ii) $m\overline{BC} < m\overline{AC}$ For first condition	
If $m\overline{BC} = m\overline{AC}$ Then $m\angle A = m\angle B$ Which is not possible	Opposite to given
For recondition condition	
If $m\overline{BC} < m\overline{AC}$ Then $m\angle A < m\angle B$ This is also not possible	Contrary to given
Now $m\overline{BC} \neq m\overline{AC}$ And $m\overline{BC} \neq m\overline{AC}$ Hence $m\overline{BC} > m\overline{AC}$	Trichotomy property of numbers.

MATHEMATICS FOR 9TH CLASS (UNIT # 12)

Pilot Superguide Mathematics 484 Class 9th

Theorem

The sum of the lengths of any two sides of a triangle is greater than the length of the third side



Given: $\triangle ABC$

- To Prove: (i) $m\overline{AB} + m\overline{AC} > m\overline{BC}$
 (ii) $m\overline{AB} + m\overline{BC} > m\overline{AC}$
 (iii) $m\overline{BC} + m\overline{CA} > m\overline{AB}$

Construction:

Take a point D on \overline{EA} such that $\overline{AD} \cong \overline{AB}$. Join B to D and name the angles $\angle 1, \angle 2$ as shown in the given figure.

Proof:

Statements	Reasons
In $\triangle ABD$	
$\angle 1 \cong \angle 2$ (i)	$\overline{AD} \cong \overline{AB}$ construction
$m\angle DBC > m\angle 1$	$m\angle DBC = m\angle 1 + m\angle ABC$
(ii) $m\angle DBC > m\angle 2$	From (i) and (i)
(iii) In $\triangle ABD \longleftrightarrow \triangle ECD$	Given
$\overline{BD} \cong \overline{CD}$	Vertical angles
$\angle 1 \cong \angle 2$	Construction
$\overline{AD} \cong \overline{ED}$	S.A.S. Postulate
$\triangle ABD \cong \triangle ECD$	Corresponding sides of $\cong \triangle$'s
$\overline{AB} \cong \overline{EC}$ (i)	$\triangle ACF$ is a triangle
$m\overline{AC} + m\overline{EC} > m\overline{AE}$ (ii)	Form I and II
$m\overline{AC} + m\overline{AB} > m\overline{AE}$	$m\overline{AE} = 2m\overline{AD}$ (construction)
Hence $m\overline{AC} + m\overline{AB} > 2m\overline{AD}$	

MATHEMATICS FOR 9TH CLASS (UNIT # 12)

Pilot Superstar Mathematics 485

Class 9th

EXERCISE 13.1

Q.1 Two sides of a triangle measure 10cm and 15cm. Which of the following measure is possible for the third side?

- (a) 5 cm (b) 20 cm (c) 25 cm (d) 30cm

Solution:

Length of one side = 10 cm
 Length of second side = 15 cm
 Sum of length of two sides = 10 + 15 = 25 cm


- (a) $5 < 25$
 (b) $20 < 25$
 (c) $25 = 25$
 (d) $30 > 25$

Sum of lengths of two sides is always greater than the third side of a triangle.

Therefore c, d are not possible.
 a and b are possible.

Q.2 O is an interior point of the $\triangle ABC$. Show that

$$m\overline{OA} + m\overline{OB} + m\overline{OC} > \frac{1}{2}(m\overline{AB} + m\overline{BC} + m\overline{CA})$$



Given: Point O is in the interior of $\triangle ABC$

Required: $m\overline{OA} + m\overline{OB} + m\overline{OC} > \frac{1}{2}(m\overline{AB} + m\overline{BC} + m\overline{CA})$

www.d-

MATHEMATICS FOR 9TH CLASS (UNIT # 12)

Pilot Superone Mathematics 486 Class 9th


Proof.

Statements	Reasons
(i) In $\triangle OAC$ $m\overline{OA} + m\overline{OC} > m\overline{AC}$	Sum of lengths of two sides is in greater than the length of the third side
(ii) In $\triangle OAB$ $m\overline{OA} + m\overline{OB} > m\overline{AB}$ In $\triangle OCB$ $m\overline{OB} + m\overline{OC} > m\overline{BC}$	Sum of lengths of two sides is in greater than the length of the third side.
(iii) $2m\overline{OA} + 2m\overline{OB} + 2m\overline{OC} > m\overline{AB} + m\overline{BC} + m\overline{CA}$	Adding (i), (ii), (iii)
$2(m\overline{OA} + m\overline{OB} + m\overline{OC}) > m\overline{AB} + m\overline{BC} + m\overline{CA}$	Dividing by 2
$\frac{1}{2}(m\overline{AB} + m\overline{BC} + m\overline{CA})$	

MATHEMATICS FOR 9TH CLASS (UNIT # 12)

Pilot Superone Mathematics 487 Class 9th

Q.3 In the $\triangle ABC$ $m\angle B = 70^\circ$ and $m\angle C = 45^\circ$ which of the sides of the triangle is longest and which is the shortest?



Solution:

Given. In $\triangle ABC$
 $m\angle B = 70^\circ$, $m\angle C = 45^\circ$

Required: (i) Which is longest from among \overline{AB} , \overline{BC} , \overline{CA}
 (ii) Which is shortest from among \overline{AB} , \overline{BC} , \overline{CA}

Proof:


Statements	Reasons
In $\triangle ABC$	
$m\angle A + m\angle B + m\angle C = 180^\circ$	Sum of angles of \triangle
$m\angle A + 70^\circ + 45^\circ = 180^\circ$	$m\angle B = 70^\circ$, $m\angle C = 45^\circ$ given
$m\angle A + 120^\circ = 180^\circ$	
$m\angle A = 180^\circ - 120^\circ$	
$m\angle A = 60^\circ$	
$m\angle A = 60^\circ$	
$m\angle B = 70^\circ$	
$m\angle C = 45^\circ$	
\overline{AC} is the longest	Opposite to the greatest angle
\overline{AB} is the shortest	Opposite to the smallest angle

MATHEMATICS FOR 9TH CLASS (UNIT # 12)

Pilot Superone Mathematics 490 Class 9th
 BD > DC

Theorem

From a point outside a line, the perpendicular is the shortest distance from the point to the line



- Given:** (i) P is a point outside \overline{AB}
 (ii) $\overline{PC} \perp \overline{AB}$
 (iii) D is any point on \overline{AB}
 (iv) P is joined to D

Required $m\angle C < m\angle D$

Proof

Statements	Reasons
In $\triangle PDC$ $m\angle C = 90^\circ$	Given
Now $m\angle D + m\angle P + m\angle C = 180^\circ$	Sum of angles of $\triangle PDC$
$m\angle D + m\angle P + 90^\circ = 180^\circ$	$m\angle C = 90^\circ$ (given)
$m\angle D + m\angle P = 180^\circ - 90^\circ$	
$m\angle D + m\angle P = 90^\circ$ (i)	(from (i))
Therefore $m\angle D < 90^\circ$	(from (i))
And $m\angle P < 90^\circ$	
Thus $m\angle D < m\angle C$	
And $m\angle PC < m\angle PD$	Side opposite to smaller

MATHEMATICS FOR 9TH CLASS (UNIT # 12)

Pilot Superone Mathematics 491 Class 9th


Hence the distance from P to AB will be greater than PC	angle
--	-------

MATHEMATICS FOR 9TH CLASS (UNIT # 12)

Pilot Superone Mathematics #92 Class 9th

EXERCISE 13.2

Q.1 In the figure, P is any point and AB is a line. Which of the following is the shortest distance between the point P and the line AB.



- (a) $m\overline{PL}$ (b) $m\overline{PM}$
 (c) $m\overline{PN}$ (d) $m\overline{PO}$

Solution: $m\overline{PN}$ is the shortest distance from P to AB

Proof:

	Statements	Reasons
In	$\triangle PNM$ $m\overline{PN} < m\overline{PO}$ (i)	$m\angle N = 90^\circ$ $m\angle O = 70^\circ$ smaller angle has small side opposite to
In	$\triangle PMO$ $m\overline{PM} > m\overline{PO}$ (ii)	$m\angle O = 70^\circ$, $m\angle M = 60^\circ$
In	$\triangle PLO$ $m\overline{PL} > m\overline{PO}$ (iii)	$m\angle I = 30^\circ$ $m\angle O = 70^\circ$ small angle has smaller side opposite to it.

Hence $m\overline{PN} < m\overline{PO}$
 $m\overline{PM} > m\overline{PO}$
 $m\overline{PL} > m\overline{PO}$


Therefore $m\overline{PN}$ is the smallest of all $\overline{PO}, \overline{PM}, \overline{PL}$

MATHEMATICS FOR 9TH CLASS (UNIT # 12)

Notes: Superior Mathematics 493 Class 9th

Q.2 In the figure, P is any point lying away from the line \overline{AB} . Then \overline{PI} will be the shortest distance if

- (a) $m\angle PLA = 80^\circ$ (b) $m\angle PLB = 100^\circ$
 (c) $m\angle PLA = 90^\circ$



Answer If $m\angle PLA = 90^\circ$

Reason: The perpendicular distance from a point out-side the line is the shortest distance from that point to the given line.

Q.3 In the figure, \overline{PI} is perpendicular to the line \overline{AB} and $m\angle N > m\angle M$. Prove that $m\overline{PN} > m\overline{PM}$.



Given: In the given figure

- (i) $\overline{PI} \perp \overline{AB}$
 (ii) $m\angle N > m\angle M$

Required: $m\overline{PN} > m\overline{PM}$

MATHEMATICS FOR 9TH CLASS (UNIT # 12)

Pilot Superone Mathematics 494 Class 9th

Proof:

Statements	Reasons
In $\triangle PIM$ $m\angle PIM = 90^\circ$ $m\angle PIM < 90^\circ$ (i)	Given In a right angle triangle other two angles are acute.
$m\angle PIM + m\angle PMN = 180^\circ$ (ii)	Adj. Supp. Angles
(iii) $m\angle PMN > 90^\circ$	
In $\triangle PIN$ $m\angle I = 90^\circ$	Given
(iv) $m\angle PNI < 90^\circ$ $m\angle PMN > m\angle PNI$	
In $\triangle PMN$ $m\angle PMN > m\angle PNI$ $m\overline{PN} > m\overline{PM}$	Proved

Q 11

www.downloadclassnotes.com

MATHEMATICS FOR 9TH CLASS (UNIT # 12)

Pilot Superone Mathematics 495

Class 9th

Review Exercise 13

- Q.1** Which of the following are true and which are false?
- (i) The angle opposite to the longer side is greater
 - (ii) In a right-angled triangle greater angle is of 60°
 - (iii) In an isosceles right-angled triangle, angles other than right angle are each of 45°
 - (iv) A triangle having two congruent sides is called equilateral triangle
 - (v) A perpendicular from a point to line is shortest distance
 - (vi) Perpendicular to line form an angle of 90°
 - (vii) A point out side the line is collinear
 - (viii) Sum of two sides of triangle is greater than the third
 - (ix) The distance between a line and a point on it is zero.
 - (x) Triangle can be formed of lengths 2 cm, 3 cm and 5 cm.


Answers.

(i)	T	(ii)	F	(iii)	T	(iv)	F	(v)	T
(vi)	T	(vii)	F	(viii)	T	(ix)	T	(x)	F

- Q.2** What will be angle for shortest distance from an outside point to the line?

Answers: 90°

- Q.3** If 13 cm, 12 cm and 5 cm are the length of a triangle, then verify that difference of measures of any two sides of a triangle is less than the measure of the third



MATHEMATICS FOR 9TH CLASS (UNIT # 12)

Pilot Supreme Mathematics 496 Class 9th

Let $m\overline{AB} = 13 \text{ cm}$

$m\overline{BC} = 12 \text{ cm}$

$m\overline{CA} = 5 \text{ cm}$


Now $m\overline{AB} - m\overline{BC} = 13 - 12 = 1 \text{ cm} < m\overline{CA} (m\overline{CA} = 5 \text{ cm})$

$m\overline{BC} - m\overline{CA} = 12 - 5 = 7 \text{ cm} < m\overline{AB} (m\overline{AB} = 13 \text{ cm})$

$m\overline{AB} - m\overline{CA} = 13 - 5 = 8 \text{ cm} < m\overline{BC} (m\overline{BC} = 12 \text{ cm})$

It shows that difference of measure of any two sides of a triangle is less than the measure of the third side.

Q.4 If 10 cm, 6 cm and 8 cm are the lengths of a triangle, then verify that sum of measures of two sides of a triangle is greater than the third side.



Let $m\overline{AB} = 10 \text{ cm}$

$m\overline{BC} = 6 \text{ cm}$

$m\overline{CA} = 8 \text{ cm}$

Now $m\overline{AB} + m\overline{BC} = 10 + 6 = 16 \text{ cm} > m\overline{CA} (m\overline{CA} = 8 \text{ cm})$

$m\overline{BC} + m\overline{CA} = 6 + 8 = 14 \text{ cm} > m\overline{AB} (m\overline{AB} = 10 \text{ cm})$

$m\overline{CA} + m\overline{AB} = 8 + 10 = 18 > m\overline{BC} (m\overline{BC} = 6 \text{ cm})$

This verifies that sum of length of any two sides of a triangle is greater than the length of the third side.


www

MATHEMATICS FOR 9TH CLASS (UNIT # 12)

Pilot Superone Mathematics 407

Class 9th

Q.5 3 cm, 4 cm and 7 cm are not the lengths of the triangle. Give the reason.



Let
 $m\overline{AB} = 7\text{ cm}$
 $m\overline{BC} = 4\text{ cm}$
 $m\overline{CA} = 3\text{ cm}$


Now (i) $m\overline{AB} + m\overline{BC} = 7 + 4 = 11$ $m\overline{CA} = 3\text{ cm}$
 (ii) $m\overline{BC} + m\overline{CA} = 4 + 3 = 7$ $m\overline{AB} = 7\text{ cm}$
 (iii) $m\overline{CA} + m\overline{AB} = 3 + 7 = 10$ $m\overline{BC} = 4\text{ cm}$

Part (ii) is not possible

Therefore 7 cm, 4 cm, 3 cm are not the sides of a triangle

Q.6 If 3 cm and 4 cm are lengths of two sides of a right triangle then what should be the third length of the triangle

Solution -



Let length of hypotenuse = x cm

$$(x)^2 = (3)^2 + (4)^2$$

$$x^2 = 16$$

$$x = 2^2$$

$$x = \sqrt{2^2} \quad (\text{taking sq. root})$$

$$x = 5\text{ cm}$$

Length of hypotenuse = 5 cm

MATHEMATICS FOR 9TH CLASS (UNIT # 14)


Pilot Superior Mathematics 198 Class 9

Unit
14

RATIO AND PROPORTIONAL

Similar Triangles

Two triangles $\triangle ABC$ and $\triangle DEF$ are similar if



In $\triangle ABC \leftrightarrow \triangle DEF$

- (i) $\angle A \cong \angle D$
 $\angle B \cong \angle E$
 $\angle C \cong \angle F$

and (ii) $\frac{m\overline{AB}}{m\overline{DE}} = \frac{m\overline{BC}}{m\overline{EF}} = \frac{m\overline{CA}}{m\overline{FD}}$

Symbolically we write it as

$$\triangle ABC \sim \triangle DEF$$

Remember:

Two congruent triangles are similar. Two similar triangles may not be congruent.

Theorem


A line parallel to one side of a triangle and intersecting the other two sides divides them proportionally.

MATHEMATICS FOR 9TH CLASS (UNIT # 14)

Pilot Superstar Mathematics

499

Class 9th



Given

(i) In $\triangle ABC$, line m cuts \overline{AB} and \overline{AC} at D and E respectively

(ii) $\overline{DE} \parallel \overline{BC}$

Required:

$$\frac{m\overline{AD}}{m\overline{DB}} = \frac{m\overline{AE}}{m\overline{EC}}$$

Construction.

(i) Join C to D

(ii) Join E to B

(iii) Take $\overline{DE} \perp \overline{AC}$ and $\overline{EM} \perp \overline{AB}$

Statements	Reasons
In $\triangle ADE$ and CDE , \overline{DE} is common Altitude	
$\triangle CDE = \frac{1}{2} \times \overline{EC} \times \overline{DE}$ (i)	Area $\Delta = \frac{1}{2} \times \text{base} \times \text{Altitude}$
$\triangle ADE = \frac{1}{2} \times \overline{AE} \times \overline{DE}$ (ii)	
(iii) $\frac{\triangle CDE}{\triangle ADE} = \frac{m\overline{EC}}{m\overline{AE}}$	Dividing (i) by (ii)
Similarly	

MATHEMATICS FOR 9TH CLASS (UNIT # 14)

Pilot Superior Mathematics 500 Class 9th

$$(iv) \frac{AB}{AD} = \frac{DB}{AD}$$

$$\text{But } \triangle ACD \cong \triangle ADB$$

Base \overline{BC} is common and
 lie between \parallel lines

$$\text{So } \frac{m\overline{EC}}{m\overline{AE}} = \frac{m\overline{DB}}{m\overline{AD}}$$

$$\text{or } \frac{m\overline{AE}}{m\overline{EC}} = \frac{m\overline{AD}}{m\overline{DB}}$$

Invert do

Invertendo

$$\text{Thus } \frac{m\overline{AD}}{m\overline{DB}} = \frac{m\overline{AE}}{m\overline{EC}}$$

Theorem

If a line segment intersects the two sides of a triangle in the same ratio then it is parallel to the third side



Given:

In $\triangle ABC$, a line segment cuts \overline{AB} at D and \overline{AC} at E.

$$\text{that } \frac{m\overline{AD}}{m\overline{DB}} = \frac{m\overline{AE}}{m\overline{EC}}$$

Required:

MATHEMATICS FOR 9TH CLASS (UNIT # 14)

Prior Superone Mathematics SUI Class 9th

$$\overline{DE} \parallel \overline{BC}$$

Construction:

Suppose $\overline{DE} \neq \overline{BC}$, then take $\overline{CP} \parallel \overline{DE}$ that meets \overline{AB} produced at P

Proof:

	Statements	Reasons
In	$\triangle APC$	
	$\overline{DE} \parallel \overline{CP}$	Construction
Therefore		
	$\frac{m\overline{AD}}{m\overline{DP}} = \frac{m\overline{AE}}{m\overline{EC}}$ (i)	
But	$\frac{m\overline{AD}}{m\overline{DP}} = \frac{m\overline{AE}}{m\overline{EC}}$ (ii)	Given
	$\frac{m\overline{AD}}{m\overline{DB}} = \frac{m\overline{AD}}{m\overline{DP}}$	From (i) and (ii)
		Property of real numbers
or	$m\overline{DB} = m\overline{DP}$	
	Therefore P lies on B	
	Hence our supposition is wrong	
	Hence $\overline{DE} \parallel \overline{BC}$	


MATHEMATICS FOR 9TH CLASS (UNIT # 14)

Pilot Superone Mathematics _____ 502

Class 9th

Exercise 14.1

Q.1. In $\triangle ABC$, $\overline{DE} \parallel \overline{BC}$



In $\triangle ABC$, $\overline{DE} \parallel \overline{BC}$

- (i) If $\overline{AD} = 1.5$ cm, $\overline{BD} = 3$ cm, $\overline{AE} = 1.3$ cm, then find \overline{CE}
- (ii) If $\overline{AD} = 2.4$ cm, $\overline{AE} = 3.2$ cm, $\overline{EC} = 4.8$ cm, find \overline{AB}
- (iii) If $\frac{\overline{AD}}{\overline{DB}} = \frac{3}{5}$ cm, $\overline{AC} = 4.8$, find \overline{AE}
- (iv) If $\overline{AD} = 2.4$ cm, $\overline{AE} = 3.2$ cm, $\overline{DE} = 2$ cm, $\overline{BC} = 5$ cm, find \overline{AB} , \overline{DB} , \overline{AC} , \overline{CE}
- (v) If $\overline{AD} = 4x - 3$, $\overline{AE} = 8x - 7$, $\overline{BD} = 3x - 1$ and $\overline{CE} = 5x - 3$, find the value of x

Solutions:

In $\triangle ABC$, $\overline{DE} \parallel \overline{BC}$

- (i) $\overline{AD} = 1.5$ cm
 $\overline{BD} = 3$ cm
 $\overline{AE} = 1.3$ cm
 $\overline{CE} = ?$

In $\triangle ABC$, $\overline{DE} \parallel \overline{BC}$

MATHEMATICS FOR 9TH CLASS (UNIT # 14)

Not a store Mathematics 503 Class 9

Let $m\angle C = x$ cm

$\frac{AD}{DB} = \frac{AE}{EC}$

Putting values of \overline{AD} , \overline{BD} and \overline{AE}

$\frac{1.5}{3} = \frac{1.3}{x}$

$$x = \frac{1.3 \times 3}{1.5} = \frac{3.9}{1.5} = \frac{39}{15} = 2.6 \text{ cm}$$

(ii)

$\overline{AD} = 2.4 \text{ cm}$
 $\overline{AE} = 3.2 \text{ cm}$
 $\overline{EC} = 4.8 \text{ cm}$
 $\overline{AB} = 9$

In $\triangle ABC$ $\overline{DE} \parallel \overline{BC}$
 Now $m\angle B = m\angle D + m\angle B$

We find $m\angle B$

Let $m\angle B = x$ cm

Now $\frac{AD}{DB} = \frac{AE}{EC}$

Putting values of \overline{AD} , \overline{AE} , \overline{EC}

$\frac{2.4}{3} = \frac{3.2}{4.8}$

$$x = \frac{4.8 \times 2.4}{3.2} = \frac{48 \times 24}{10 \times 10} = \frac{10}{32}$$

MATHEMATICS FOR 9TH CLASS (UNIT # 14)

Pilot Superstore Mathematics

(iii)

$$\frac{mAD}{mDB} = \frac{3}{5}$$

AB

$$mAD + mDB = 24 + 36 = 60 \text{ cm}$$

MAC

MAI

$$48 \text{ cm}$$

In ABC

AD

DB

and

AD

DB

Putting values of AD, DB, AC

$$\frac{3}{5}$$

$$\frac{48 - x}{x}$$

or

$$3x = 48 - x$$

$$3x + x = 48$$

$$4x = 48$$

or

$$x = 12$$

Now mAE

AC

CF


mAE

$$48 - 3 = 45 \text{ cm}$$

MATHEMATICS FOR 9TH CLASS (UNIT # 14)

Pilot Superone Mathematics 505 Class 9th

- (iv) $AD = 2.4$ cm (i)
 $AF = 3.2$ cm (ii)
 $DE = 2$ cm (iii)
 $BC = 5$ cm
 $AB = ?$, $DB = ?$, $AC = ?$, $CE = ?$



Construction:

Take $\overline{DF} \parallel \overline{AC}$
 Now $DF \parallel AC$ is given

$$m\overline{DF} = m\overline{DE} = 2 \text{ cm}$$

$$\text{Thus } m\overline{BF} = 5 - 2 = 3 \text{ cm}$$

Now in $\triangle ABC$ $\overline{DF} \parallel \overline{AC}$

$$\frac{CE}{EA} = \frac{CF}{FB}$$

Putting values of EA, CF, FB

$$\frac{CE}{3.2} = \frac{3}{2}$$

$$\overline{CE} = \frac{3 \times 3.2}{2}$$

$$\overline{CE} = 3 \times 1.6 = 4.8 \text{ cm} \quad (iv)$$

Now in $\triangle ABC$ $DE \parallel BC$

MATHEMATICS FOR 9TH CLASS (UNIT # 14)

Pilot Supercone Mathematics 506 Class 9th

$$\frac{\overline{BD}}{\overline{AD}} = \frac{\overline{CE}}{\overline{AE}}$$

$$\frac{2.4}{\overline{BD}} = \frac{4.8}{3.2}$$

(Putting the values of \overline{AD} , \overline{CE} , \overline{AE})

$$\overline{BD} = \frac{2.4 \times 4.8}{3.2} \quad (v)$$

$$= 3.6 \text{ cm}$$

Now $\overline{AB} = \overline{AD} + \overline{DB}$

$$= 2.4 + 3.6 \quad (i), (v)$$

$$= 6.0 \text{ cm}$$

Hence $\overline{DB} = 3.6 \text{ cm} \quad (v)$

$$\overline{AC} = \overline{AE} + \overline{EC}$$

$$= 3.2 + 4.8 \quad \text{From (ii), (iv)}$$

$$= 8.0 \text{ cm}$$

$$\overline{CE} = 4.8 \text{ cm} \quad \text{From (iv)}$$

(v)



$$\begin{aligned} \overline{AD} &= 4x \quad 3 \\ \overline{AE} &= 8x \quad 7 \\ \overline{BD} &= 3x \quad 1 \\ \overline{CE} &= 5x \quad 3 \\ x &= ? \end{aligned}$$

Now in $\triangle ABC$ $\overline{DE} \parallel \overline{BC}$

$$\frac{\overline{AD}}{\overline{BD}} = \frac{\overline{AE}}{\overline{CE}}$$

MATHEMATICS FOR 9TH CLASS (UNIT # 14)

Pilot Superone Mathematics 507 Class 9th

Putting values

$$\frac{4x-3}{3x-1} = \frac{8x-7}{5x-3}$$

Now $(4x-3)(5x-3) = (8x-7)(3x-1)$

$$20x^2 - 27x + 9 = 24x^2 - 29x + 7$$

or $20x^2 - 24x^2 - 27x + 29x + 9 - 7 = 0$

$$-4x^2 + 2x + 2 = 0$$

Dividing by 2

or $2x^2 - x - 1 = 0$

$$2x^2 - 2x + x - 1 = 0$$

$$2x(x-1) + 1(x-1) = 0$$

$$(x-1)(2x+1) = 0$$

or $x-1 = 0$

$$x = 1$$

or $2x+1 = 0$

$$2x = -1$$

$$x = -\frac{1}{2}$$

$$x = -\frac{1}{2}$$

$$x = -\frac{1}{2}$$

(impossible)

$$x = 1$$

Q.2. If $\triangle ABC$ is an isosceles triangle, $\angle A$ is vertex angle and \overline{DE} intersects the sides \overline{AB} and \overline{AC} as shown in the figure so that


$$m\angle D = m\angle B \quad m\angle E = m\angle C$$

prove that $\triangle ADE$ is also an isosceles triangle



MATHEMATICS FOR 9TH CLASS (UNIT # 14)

Pink Superone Mathematics SOB Class 9th



Given:

In $\triangle ABC$, $\angle A$ is its vertex

$$\overline{AB} = \overline{AC}$$

$$\frac{m\overline{AD}}{m\overline{DB}} = \frac{m\overline{AE}}{m\overline{EC}}$$

$$\frac{m\overline{AD}}{m\overline{DB}} = \frac{m\overline{AE}}{m\overline{EC}}$$

Required:

$$\overline{AD} = \overline{AE}$$

Proof.

$$\text{Now } \frac{m\overline{DB}}{m\overline{AD}} = \frac{m\overline{EC}}{m\overline{AE}} \quad \text{given}$$

$$\text{or } \frac{m\overline{DB} + m\overline{AD}}{m\overline{AD}} = \frac{m\overline{EC} + m\overline{AE}}{m\overline{AE}}$$

$$\frac{m\overline{AB}}{m\overline{AD}} = \frac{m\overline{AC}}{m\overline{AE}}$$

$$\frac{m\overline{AB}}{m\overline{AD}} = \frac{m\overline{AC}}{m\overline{AE}}$$

$$\frac{m\overline{AB}}{m\overline{AD}} = \frac{m\overline{AC}}{m\overline{AE}}$$

$$m\overline{AB} = m\overline{AC} \quad (\text{given})$$

$\triangle ADE$ is also an isosceles triangle


Q.3. In an equilateral triangle ABC shown in the figure.

$$m\overline{AE} : m\overline{EC} = m\overline{AD} : m\overline{DB}$$

MATHEMATICS FOR 9TH CLASS (UNIT # 14)

Pilot Superior Mathematics 509 Class 9

Find all the three angles of $\triangle ADE$ and name it



Given:

$\triangle ABC$ is an equilateral triangle i.e. $AB \cong BC \cong AC$ also

$$m\angle A = m\angle B = m\angle C = 60^\circ \text{ and } \frac{m\overline{AE}}{m\overline{AC}} = \frac{m\overline{AD}}{m\overline{AB}}$$

Required:

To find measures of angles of $\triangle ADE$


Solution:

Statements	Reasons
$\frac{m\overline{AE}}{m\overline{AC}} = \frac{m\overline{AD}}{m\overline{AB}}$	Given
$DE \parallel BC$	
$m\angle A = m\angle B = m\angle C = 60^\circ$	Given
Now $\overline{DE} \parallel \overline{BC}$	Corresponding angles
$\therefore m\angle 1 = m\angle B$	
60°	Corresponding angles
and $m\angle 2 = m\angle C$	
60°	Corresponding angles
$m\angle A = 60^\circ$	
$\triangle ADE$ is an isosceles triangle	Given

MATHEMATICS FOR 9TH CLASS (UNIT # 14)

Pict Supering Mathematics 510 Class 9th

Q.4. Prove that the line segment drawn through the mid-point of one side of a triangle and parallel to another side bisects the third side.



Given:
 In $\triangle ABC$

$$m\overline{AD} = m\overline{DB} \quad (i)$$

$$DE \parallel BC$$

Required.

$$m\overline{AE} = m\overline{EC}$$

Proof:

Statements	Reasons
In $\triangle ABC$	
$\overline{DE} \parallel \overline{BC}$	Given
(i) $\frac{m\overline{AD}}{m\overline{DB}} = \frac{m\overline{AE}}{m\overline{EC}}$	Theorem
Now $m\overline{AD} = m\overline{DB}$	Given
$\frac{m\overline{DB}}{m\overline{DB}} = \frac{m\overline{AE}}{m\overline{EC}}$	Putting $m\overline{AD} = m\overline{DB}$ in (i)
$1 = \frac{m\overline{AE}}{m\overline{EC}}$	

MATHEMATICS FOR 9TH CLASS (UNIT # 14)

Notes: Supreme Mathematics

511


Class 9th

$$\frac{m\overline{AE}}{m\overline{EC}}$$

or $m\overline{AE} = m\overline{EC}$

Hence, \overline{DE} bisects \overline{AC}

Q.5. Prove that the line segment joining the mid-points of any two sides of a triangle is parallel to the third side.



Given:

In $\triangle ABC$

$$m\overline{AD} = m\overline{DB}$$

$$m\overline{AE} = m\overline{EC}$$

i.e. $\frac{m\overline{AD}}{m\overline{DB}} = \frac{m\overline{AE}}{m\overline{EC}}$

Required:

$$\overline{DE} \parallel \overline{BC}$$

Proof:

Statements	Reasons
If $\overline{DE} \nparallel \overline{BC}$ then take $\overline{BK} \parallel \overline{DE}$ which cuts produced \overline{AC} at k	"

MATHEMATICS FOR 9TH CLASS (UNIT # 14)

Pilot Superone Mathematics 512 Class 9th

Now

$$(i) \frac{m\overline{AD}}{m\overline{DB}} = \frac{m\overline{AE}}{m\overline{EC}}$$

$$(ii) \text{ Now } \frac{m\overline{AD}}{m\overline{DB}} = \frac{m\overline{AE}}{m\overline{EC}}$$

Given

$$\frac{m\overline{AE}}{m\overline{EC}} = \frac{m\overline{AD}}{m\overline{DB}}$$

(From i. and ii)


$$\text{i.e. } m\overline{AE} = m\overline{AD}$$

This is possible only when K lies on C

$$\text{Hence, } \overline{AD} = \overline{BC}$$

Theorem

The internal bisector of an angle of a triangle divides the side opposite to it in the ratio of the length of the sides containing the angle.



Given:

In $\triangle ABC$ internal bisector of $\angle A$ meets \overline{BC} at D

MATHEMATICS FOR 9TH CLASS (UNIT # 14)

Page Sandstone Mathematics 513 Class 9th

To prove:

$$\frac{mAC}{mAB} = \frac{mCD}{mBD}$$

Construction:

Draw $BK \parallel DA$ which meets CA produced at K

Proof:


Statements	Reasons
$AD \parallel BK$ and CK cuts them $m\angle 1 = m\angle 2$	Corresponding angles
$AD \parallel KB$ and AB cuts them $m\angle 3 = m\angle 4$	Alternate angles
But $m\angle 1 = m\angle 3$ $m\angle 2 = m\angle 4$	Given
and $AK \cong AB$ or $AK = AB$	
Now $AD \parallel KB$	Construction
$\frac{mBD}{mDC} = \frac{mKA}{mAC}$	Theorem
or $\frac{mBD}{mDC} = \frac{mAB}{mAC}$	
or $\frac{mAC}{mAB} = \frac{mDC}{mBD}$	Proved

MATHEMATICS FOR 9TH CLASS (UNIT # 14)

Pilot Superior Mathematics 514 Class 9th

Theorem

If two triangles are similar, then the measures of their corresponding sides are proportional.



Given:

$\triangle ABC \sim \triangle DEF$

i.e. $\angle A \cong \angle D$, $\angle B \cong \angle E$ and $\angle C \cong \angle F$

To Prove:

$$\frac{m\overline{AB}}{m\overline{DE}} = \frac{m\overline{AC}}{m\overline{DF}} = \frac{m\overline{BC}}{m\overline{EF}}$$

Construction:

(i) Suppose that $m\overline{AB} > m\overline{DE}$

(ii) $m\overline{AB} \leq m\overline{DE}$

On \overline{AB} take a point L such that $m\overline{AL} = m\overline{DE}$

On \overline{AC} take a point M such that $m\overline{AM} = m\overline{DF}$ Join L and M by the line segment LM.

Proof:

Statements	Reasons
(i) In $\triangle ALM \leftrightarrow \triangle DEF$	
$\angle A \cong \angle D$	Given
$\overline{AL} \cong \overline{DE}$	Construction
$\overline{AM} \cong \overline{DF}$	Construction
Thus $\triangle ALM \cong \triangle DEF$	S.A.S. postulate
$\angle L \cong \angle E$	Corresponding angles

MATHEMATICS FOR 9TH CLASS (UNIT # 14)

Pilot Superone Mathematics 515

Class 9th

$\angle M \cong \angle F$
 and $\angle I \cong \angle F$, $\angle M \cong \angle F$

Now $\angle E \cong \angle B$ and $\angle F \cong \angle C$
 $\angle L \cong \angle B$, $\angle M \cong \angle C$

Thus $\overline{MI} \parallel \overline{BC}$

Hence, $\frac{m\overline{AI}}{m\overline{AB}} = \frac{m\overline{DI}}{m\overline{AC}}$

or $\frac{m\overline{DE}}{m\overline{AB}} = \frac{m\overline{DF}}{m\overline{AC}}$ (i)

Similarly by intercepting segments on \overline{BA} and \overline{BC} , we can prove that

$$\frac{m\overline{DE}}{m\overline{AB}} = \frac{m\overline{EF}}{m\overline{BC}} \quad (ii)$$

Thus $\frac{m\overline{DE}}{m\overline{AB}} = \frac{m\overline{DF}}{m\overline{AC}} = \frac{m\overline{EF}}{m\overline{BC}}$

$$\frac{m\overline{AB}}{m\overline{DE}} = \frac{m\overline{AC}}{m\overline{DF}} = \frac{m\overline{BC}}{m\overline{EF}}$$

(II) If $m\overline{AB} < m\overline{DE}$, it can similarly be proved by taking intercepts on the sides of $\triangle DEF$

If $m\overline{AB} = m\overline{DE}$, then

of congruent Δ
 (Corresponding angles of congruent triangles)
 Given
 Transitivity of congruence

Corresponding angles are equal
 by Theorem 14.11

$m\overline{AI} = m\overline{DI}$ and
 $m\overline{AM} = m\overline{DF}$
 (Construction)

by (i) and (ii)


by taking reciprocals

MATHEMATICS FOR 9TH CLASS (UNIT # 14)

Pilot Superstar Mathematics 516 Class 9th


In $\triangle ABC \cong \triangle DEF$ Given
 $\angle A \cong \angle D$ Given
 $\angle B \cong \angle E$
 and $AB \cong DE$ A.S.A \cong A.S.A
 So $\triangle ABC \cong \triangle DEF$
 Thus

$\frac{m\overline{AB}}{m\overline{DE}} = \frac{m\overline{AC}}{m\overline{DF}} = \frac{m\overline{BC}}{m\overline{EF}}$ $\therefore \overline{AC} \cong \overline{DF}, \overline{BC} \cong \overline{EF}$
 Hence the result is true for all the cases



Exercise 14.2

- Q.1. In $\triangle ABC$ as shown in the figure, \overline{CD} bisects $\angle C$
 and meets \overline{AB} at D. $m\overline{BD}$ is equal to
 (a) 5 (b) 16 (c) 10 (d) 18



Solution:

$$\begin{aligned} \frac{m\overline{BD}}{m\overline{DA}} &= \frac{m\overline{BC}}{m\overline{CA}} \\ \frac{m\overline{BD}}{6} &= \frac{10}{12} \\ m\overline{BD} &= \frac{10}{12} \times 6 \\ m\overline{BD} &= 5 \end{aligned}$$


MATHEMATICS FOR 9TH CLASS (UNIT # 14)

Pilot Superone Mathematics

517

(Class 9)

Q.2. $\triangle ABC$ shown in the figure. \overline{CD} bisects $\angle C$. If $m\overline{AC} = 3$, $m\overline{CB} = 6$ and $m\overline{AB} = 7$, then find $m\overline{AD}$ and $m\overline{DB}$



Solution:


Let $m\overline{AD} = x$
 $m\overline{DB} = 7 - x$
 $\frac{m\overline{AD}}{m\overline{DB}} = \frac{m\overline{AC}}{m\overline{CB}}$
 Putting values
 $\frac{x}{7 - x} = \frac{3}{6}$
 or $\frac{x}{7 - x} = \frac{1}{2}$
 or $\frac{2(x)}{2(7 - x)} = \frac{1}{2}$
 $\frac{2x}{7 - x} = \frac{1}{2}$
 $2x \cdot 2 = 1(7 - x)$
 $4x = 7 - x$
 $4x + x = 7$
 $5x = 7$
 or $x = \frac{7}{5}$
 or $m\overline{AD} = \frac{7}{5}$
 and $m\overline{DB} = 7 - \frac{7}{5}$
 $= \frac{35}{5} - \frac{7}{5}$
 $= \frac{28}{5}$

MATHEMATICS FOR 9TH CLASS (UNIT # 14)

Pilot Superone Mathematics 518 Class 9th

$$m\angle B = \frac{21}{3} = \frac{7}{1} = \frac{14}{3}$$

Q.3. Show that in any correspondence of two triangles, if two angles of one triangle are congruent to the corresponding angles of the other, then the triangles are similar.



Given.

$\triangle ABC$ and $\triangle DEF$

$m\angle B = m\angle E$

$m\angle C = m\angle F$

Required:

$\triangle ABC \sim \triangle DEF$


Proof.

Statements	Reasons
$m\angle B + m\angle C + m\angle A = 180^\circ$	Angles of $\triangle ABC$
$m\angle E + m\angle F + m\angle D = 180^\circ$	Angles of $\triangle DEF$
$m\angle B = m\angle E$ & $m\angle C = m\angle F$	As $m\angle B = m\angle E$, $m\angle C = m\angle F$
$m\angle A = m\angle D$	Subtracting (ii) from (i)
or $m\angle A = m\angle D$	
Now in $\triangle ABC$, $\triangle DEF$	
$m\angle A = m\angle D$	
$m\angle B = m\angle E$	
$m\angle C = m\angle F$	
Hence $\triangle ABC \sim \triangle DEF$	

MATHEMATICS FOR 9TH CLASS (UNIT # 14)

Pilot Superline Mathematics 519 Class 9th

Q.4 If line segments AB and CD are intersecting at point X and $\frac{m\overline{AX}}{m\overline{XB}} = \frac{m\overline{CX}}{m\overline{XD}}$. Then show that $\triangle AXC$ and $\triangle BXD$ are similar.



Given:

AB and CD intersect at X that $\frac{m\overline{AX}}{m\overline{XB}} = \frac{m\overline{CX}}{m\overline{XD}}$

Required:

$\triangle AXC \sim \triangle BXD$

Proof:

Statements	Reasons
In $\triangle AXC$ and $\triangle BXD$	
$\angle 1 \cong \angle 2$	Vertical opp. Angles
$\frac{m\overline{AX}}{m\overline{XB}} = \frac{m\overline{CX}}{m\overline{XD}}$	Given
$AC \parallel BD$	Corresponding angles (corresponding angles)
$\angle 4 \cong \angle 3$	
$\angle 6 \cong \angle 5$	
Thus	
$\frac{m\overline{AX}}{m\overline{XB}} = \frac{m\overline{CX}}{m\overline{XD}} = \frac{m\overline{AC}}{m\overline{BD}}$	
Hence $\triangle AXC \sim \triangle BXD$	

MATHEMATICS FOR 9TH CLASS (UNIT # 14)

Pilot Superone Mathematics 520 Class 9

Review Exercise 14

Q.1. Which of the following are true and which are false?

- (i) Congruent triangles are of same size and shape
- (ii) Similar triangles are of same shape but different sizes
- (iii) Symbol used for congruent is ' \cong '
- (iv) Symbol used for similarity is ' \sim '
- (v) Congruent triangles are similar
- (vi) Similar triangles are congruent
- (vii) A line segment has only one mid point
- (viii) One and only one line can be draw through two points
- (ix) Proportion is non-equality of two ratios.
- (x) Ratio has no unit

Answers:

- | | | | | |
|------|-------|--------|------|-----|
| (i) | (ii) | (iii) | (iv) | (v) |
| (vi) | (vii) | (viii) | (ix) | (x) |

Q.2. Define the following:

- | | |
|---------------------------|------------------------|
| (i) Ratio | (ii) Proportion |
| (iii) Congruent Triangles | (iv) Similar Triangles |

Answers:

(i) Ratios

We define ratio $a : b = \frac{a}{b}$ as the comparison of two like

quantities 'a' and 'b' called the elements of a ratio. 'a' is the first element and 'b' is called the second element of the ratio.

(ii) Proportion:

The relation of equality of two ratios is called proportion. If $a : b = c : d$, then a, b, c and d are said to be in proportion.

(iii) Congruent Triangles:

MATHEMATICS FOR 9TH CLASS (UNIT # 14)

Pilot Superone Mathematics 531 Class 9th

If three sides of a triangle are equal in length to the corresponding three sides of another triangles then the triangles are called congruent triangles

(iv) **Similar Triangles**

If three angles of a triangle are congruent to three corresponding angles of another triangle and then corresponding sides are proportional then these triangles are called similar triangles

Remember:

Congruent triangles are also similar triangles but all similar triangles may or may not be congruent triangles

Q.3 In ΔMN shown in the figure, $\overline{MN} \parallel \overline{PQ}$

(i) If $m\angle M = 5\text{cm}$, $m\angle P = 2.5\text{cm}$, $m\angle Q = 2.3\text{cm}$, then find $m\angle N$

(ii) If $m\angle M = 6\text{cm}$, $m\angle Q = 2.5\text{cm}$, $m\angle N = 5\text{cm}$, then find $m\angle P$



30) Given:
 In ΔMN

$$\overline{MN} \parallel \overline{PQ}$$

$$m\angle M = 5\text{cm}, m\angle P = 2.5\text{cm}$$

$$\text{and } m\angle Q = 2.3\text{cm}$$

Required:

$$m\angle N = ?$$

MATHEMATICS FOR 9TH CLASS (UNIT # 14)

Pilot Superone Mathematics 522 Class 9th

Calculations:

$$\begin{aligned} \text{Now } \frac{m\angle N}{m\angle Q} &= \frac{m\angle M}{m\angle P} && \text{as } \overline{PQ} \parallel \overline{MN} \\ \frac{m\angle N}{23} &= \frac{5}{25} && \text{putting the values} \\ \therefore \angle N &= \frac{5 \times 23}{25} \\ &= \frac{5 \times 23}{25} = 4.6 \text{ cm} \end{aligned}$$

3(ii) Given:
In ΔMN

$$\overline{MN} \parallel \overline{PQ}$$

$$m\angle QN = 5 \text{ cm. and } m\angle Q = 25 \text{ cm, } m\angle M = 6 \text{ cm}$$

Required:

$$m\angle P$$

Calculations:

$$\begin{aligned} \frac{m\angle P}{m\angle M} &= \frac{m\angle Q}{m\angle N} && \text{as } \overline{PQ} \parallel \overline{MN} \\ \frac{m\angle P}{m\angle M} &= \frac{m\angle Q}{m\angle Q + m\angle QN} \end{aligned}$$

Putting values


$$\begin{aligned} \frac{m\angle P}{6} &= \frac{25}{25 + 5} \\ m\angle P &= \frac{25}{75} \times 6 \\ &= \frac{1}{3} \times 6 \\ \angle P &= 2 \text{ cm} \end{aligned}$$

MATHEMATICS FOR 9TH CLASS (UNIT # 14)

Prior Superone Mathematics 523

Class 9th

Q.4 In the shown figure, let $m\overline{PA} = 8x - 7$, $m\overline{PB} = 4x - 3$, $m\overline{AQ} = 5x - 3$, $m\overline{BR} = 3x - 1$. Find the value of x if $\overline{AB} \parallel \overline{QR}$.



Given.

In $\triangle PQR$, $\overline{AB} \parallel \overline{QR}$

and $m\overline{PA} = 8x - 7$

$m\overline{PB} = 4x - 3$

$m\overline{AQ} = 5x - 3$

$m\overline{BR} = 3x - 1$

$\overline{AB} \parallel \overline{QR}$

Required:

Calculations:

Statements	Reasons
In $\triangle PQR$	
$m\overline{PA} = m\overline{PB}$	
$m\overline{AQ} = m\overline{BR}$	$\overline{AB} \parallel \overline{QR}$
Putting values	
$8x - 7 = 4x - 3$	
$5x - 3 = 3x - 1$	
$(8x - 7) - (4x - 3) = (5x - 3) - (3x - 1)$	
$24x - 29x + 7 = 20x - 27x + 9$	

MATHEMATICS FOR 9TH CLASS (UNIT # 14)

Pilot Superone Mathematics

9th Class

Class 9

$$24x - 20x - 29x + 27x + 7 \neq 0$$

$$4x - 2x - 2 = 0$$

$$2x^2 - x - 1 = 0$$

$$2x^2 - 2x + x - 1 = 0$$

$$2x(x - 1) + 1(x - 1) = 0$$

$$(x - 1)(2x + 1) = 0$$

$$x - 1 = 0$$

$$2x + 1 = 0$$

$$2x = -1$$

$$x = -\frac{1}{2}$$

$$\text{or } x - 1 = 0$$

$$x = 1$$

$$x = \frac{1}{2}$$

is not possible

Q.5. In $\triangle LMN$ shown in the figure, \overline{LT} bisects $\angle L$. If $m\overline{LN} = 4$, $m\overline{MT} = 6$, $m\overline{TN} = 8$, then find $m\overline{MT}$ and $m\overline{TN}$.



Given:

In $\triangle LMN$, \overline{LT} bisector of $\angle L$
 $m\overline{LN} = 4$ cm, $m\overline{MT} = 6$ cm, $m\overline{TN} = 8$ cm

Required:

$m\overline{MA} = ?$ and

$m\overline{AN} = ?$

MATHEMATICS FOR 9TH CLASS (UNIT # 14)

Pilot Superiore Mathematics

Class 9th

Calculations:

Let $\overline{MA} = x \text{ cm}$

$\overline{MA} = 8 - x \text{ cm}$

Now $\frac{\overline{MA}}{\overline{AN}} = \frac{\overline{MA}}{\overline{AN}}$ (\overline{AK} is bisector of $\angle A$)

Putting the values

$\frac{8-x}{x} = \frac{6}{4}$

$4(8-x) = 6x$

$32 - 4x = 6x$

$32 = 6x + 4x$


$10x = 32$

$x = 3.2$

$\overline{AN} = 3.2 \text{ cm}$

and $\overline{MA} = 8 - x$
 $= 8 - 3.2 = 4.8 \text{ cm}$

Q.6. In isosceles $\triangle PQR$ shown in the figure find the value of x and y .



MATHEMATICS FOR 9TH CLASS (UNIT # 14)

Pilot Superone Mathematics 576 Class 9th

Given:

$$\overline{PQ} \parallel \overline{PR}$$

In $\triangle PQR$

$$\overline{PL} \perp \overline{QR}$$

Required:

$$x = ? \quad y = ?$$

Proof:

Statements	Reasons
In $\triangle PRL$ and $\triangle PQL$	
$m\overline{PQ} = m\overline{PR}$	(i) Each 90°
$\angle PLQ = \angle PLR$	Common
$m\overline{PL} = m\overline{PL}$	HS \cong H.S
$\triangle PQL \cong \triangle PRL$	
$m\overline{QL} = m\overline{RL}$	Corresponding sides of congruent triangles.
$6 = x$	
$y = 6 \text{ cm}$	
$x = 10 \text{ cm}$	

MATHEMATICS FOR 9TH CLASS (UNIT # 15)

Pilot Superior Mathematics 527

Class 9th

Unit
15

PYTHAGORAS THEOREM

In a right angled triangle, the square of the length of hypotenuse is equal to the sum of the squares of the lengths of the other two sides.



Given

$\triangle ACB$ is right angled triangle in which $m\angle C = 90^\circ$ and
 $m\overline{BC} = a$, $m\overline{AC} = b$ and $m\overline{AB} = c$

To prove

$$c^2 = a^2 + b^2$$

Construction

Draw \overline{CD} perpendicular from C on \overline{AB} .

Let $m\overline{CD} = h$, $m\overline{AD} = x$ and $m\overline{BD} = y$. Line segment \overline{CD} splits $\triangle ABC$ into two $\triangle s$ ADC and BDC which are separately shown in the figures (ii) -a and (ii) -b respectively

Proof (Using similar $\triangle s$)

Statements	Reasons
In $\triangle ADC \sim \triangle ACB$	Refer to figure (ii) -a and (i)
$\angle A \cong \angle A$	common self congruent
$\angle ADC \cong \angle ACB$	Construction given, each angle = 90°
$\angle C \cong \angle B$	$\angle C$ and $\angle B$, complements of $\angle A$
$\triangle ADC \sim \triangle ACB$	Congruency of three angles

MATHEMATICS FOR 9TH CLASS (UNIT # 15)

Pilot Superone Mathematics 528 Class 9


$\frac{x}{b} = \frac{b}{c}$ or $x = \frac{b^2}{c}$(I)	(Measures of corresponding sides of similar triangles are proportional)
Again in $\triangle BDC \sim \triangle BCA$ $\angle B \cong \angle B$ $\angle BDC \cong \angle BCA$ $\angle C \cong \angle C$ $\triangle BDC \sim \triangle BCA$ $\frac{y}{a} = \frac{a}{c}$ or $y = \frac{a^2}{c}$(II)	Refer to figure (ii)- b and (i) Common-angle congruent Construction given, each angle = 90° $\angle C$ and $\angle A$, complementary $\angle B$ Congruency of three angles. (Corresponding sides of similar triangles are proportional).
But $y + x = c$ $\frac{a^2}{c} + \frac{b^2}{c} = c$ or $a^2 + b^2 = c^2$ i.e., $c^2 = a^2 + b^2$	Supposition from (I) and (II) Multiplying both sides by c.

Converse of Pythagoras' Theorem

If the square of length one side of a triangle is equal to the sum of the squares of the lengths of the other two sides then the triangle is a right angled triangle.

MATHEMATICS FOR 9TH CLASS (UNIT # 15)

Pilot Superone Mathematics 529 Class 9th



Given: In a $\triangle ABC$, $m\overline{AB} = c$, $m\overline{BC} = a$ and $m\overline{AC} = b$ such that
 $a^2 + b^2 = c^2$

To prove: $\triangle ABC$ is a right angled triangle.

Construction: (i) Draw \overline{CD} perpendicular to \overline{BC} such that
 $\overline{CD} \cong \overline{CA}$

(ii) Join B to D.

Proof

Statements	Reasons
$\triangle DCB$ is a right angled triangle.	Construction
$(m\overline{BD})^2 = a^2 + b^2$ (i)	Pythagoras theorem
But $a^2 + b^2 = c^2$ (ii)	Given
$(m\overline{BD})^2 = c^2$	From i. ii
or $(m\overline{BD}) = c$	
Now in $\triangle DCB \leftrightarrow \triangle ACB$	
$\overline{CD} \cong \overline{CA}$	Construction
$\overline{BC} \cong \overline{BC}$	Common
$\overline{DB} \cong \overline{AB}$	each = c
$\triangle DCB \cong \triangle ACB$	S.S.S \cong S.S.S
$\angle DCB \cong \angle ACB$	Corresponding angle of congruent triangles.
But $m\angle DCB = 90^\circ$	Construction
$m\angle ACB = 90^\circ$	
Hence $\triangle ACB$ is a right angled triangle	

MATHEMATICS FOR 9TH CLASS (UNIT # 15)

Pilot Superior Mathematics 530 Class 9th

EXERCISE 15

Q.1. Verify that the Δ s having the following measures of sides are right angled.

- (i) $a = 5\text{cm}$, $b = 12\text{cm}$, $c = 13\text{cm}$
 (ii) $a = 1.5\text{cm}$, $b = 2\text{cm}$, $c = 2.5\text{cm}$
 (iii) $a = 9\text{cm}$, $b = 12\text{cm}$, $c = 15\text{cm}$
 (iv) $a = 16\text{cm}$, $b = 30\text{cm}$, $c = 34\text{cm}$

Solutions: 1(i)

$$\left. \begin{array}{l} a = 5\text{cm} \\ b = 12\text{cm} \\ c = 13\text{cm} \end{array} \right\} \begin{array}{l} a^2 = (5)^2 = 25 \\ b^2 = (12)^2 = 144 \\ c^2 = (13)^2 = 169 \end{array}$$

Now we see $c^2 = a^2 + b^2$
 $169 = 25 + 144 = 169$

The given lengths are the sides of a right angled triangle

1 (ii)

$$\left. \begin{array}{l} a = 1.5\text{cm} \\ b = 2\text{cm} \\ c = 2.5\text{cm} \end{array} \right\} \text{and} \begin{array}{l} a^2 = (1.5)^2 = 2.25 \\ b^2 = (2)^2 = 4 \\ c^2 = (2.5)^2 = 6.25 \end{array}$$

Now $c^2 = a^2 + b^2$
 $6.25 = 2.25 + 4 = 6.25$

The given lengths are the lengths of the sides of a right angled triangle

1 (iii)

$$\left. \begin{array}{l} a = 9\text{cm} \\ b = 12\text{cm} \\ c = 15\text{cm} \end{array} \right\} \text{and} \begin{array}{l} a^2 = (9)^2 = 81 \\ b^2 = (12)^2 = 144 \\ c^2 = (15)^2 = 225 \end{array}$$

Now we see $c^2 = a^2 + b^2$
 $225 = 81 + 144 = 225$

MATHEMATICS FOR 9TH CLASS (UNIT # 15)

Pilot Superone Mathematics 531 Class 9th

The given lengths are the lengths of the sides of a right angled triangle

1 (iv)

$$\left. \begin{array}{l} a = 16 \text{ cm} \\ b = 30 \text{ cm} \\ c = 34 \text{ cm} \end{array} \right\} \text{ and } \left. \begin{array}{l} a^2 = (16)^2 = 256 \\ b^2 = (30)^2 = 900 \\ c^2 = (34)^2 = 1156 \end{array} \right\}$$

$$c^2 = a^2 + b^2$$

Now we see $1156 = 256 + 900 = 1156$

The given lengths are the lengths of the sides of a right angled triangle

Q.2. Verify that $a^2 + b^2$, $a^2 - b^2$ and $2ab$ are the measures of the sides of a right angled triangle where a and b are any two real numbers ($a > b$).

Solution:

$$\left. \begin{array}{l} a^2 + b^2 \\ a^2 - b^2 \\ 2ab \end{array} \right\} \left\{ \begin{array}{l} (a^2 + b^2)^2 = a^4 + b^4 + 2a^2b^2 \quad (i) \\ (a^2 - b^2)^2 = a^4 + b^4 - 2a^2b^2 \quad (ii) \\ (2ab)^2 = 4a^2b^2 \quad (iii) \end{array} \right.$$

Now we see if

$$(a^2 + b^2)^2 = (a^2 - b^2)^2 + (2ab)^2$$

From i, ii, iii

$$\begin{aligned} a^4 + b^4 + 2a^2b^2 &= (a^4 + b^4 - 2a^2b^2) + 4a^2b^2 \\ &= a^4 + b^4 - 2a^2b^2 + 4a^2b^2 \end{aligned}$$

$$a^4 + b^4 + 2a^2b^2 = a^4 + b^4 + 2a^2b^2$$

The given lengths are the lengths of the side of a right angled triangle

Q.3. The three sides of a triangle are of measure 8, x and 17 respectively. For what value of x will it become base of a right angled triangle?

Solution: Lengths of the sides of a triangle

$$a = 8$$

$$c = 17$$

MATHEMATICS FOR 9TH CLASS (UNIT # 15)

Pilot Superone Mathematics 532 Class 9th

$$b = x$$

$$c = 17$$

In case of right angled triangle

$$c^2 = a^2 + b^2$$

$$(17)^2 = (8)^2 + (x)^2$$

$$289 = 64 + x^2$$

$$x^2 + 64 = 289$$

$$x^2 = 289 - 64$$

$$x^2 = 225$$


Thus $x = 15$ (Taking sq. root)

Q.4. In an isosceles Δ , the base $\overline{BC} = 28$ cm,

and $\overline{AB} = \overline{AC} = 50$ cm.

If $\overline{AD} \perp \overline{BC}$, then find

- (i) Length of \overline{AD} (ii) area of ΔABC



Given in ΔABC

$$m\overline{AC} = m\overline{AB} = 50 \text{ cm}$$

$$m\overline{BC} = 28 \text{ cm}$$

$$\overline{AD} \perp \overline{BC}$$

Required: $m\overline{AD} = ?$

Area of ΔABC


Statements	Reasons
In $\Delta ACD \rightarrow \Delta ABD$	

MATHEMATICS FOR 9TH CLASS (UNIT # 15)

Pilot Superone Mathematics	533	Class 9 th
$m\angle ADC = m\angle ADB$ $\overline{mAC} = \overline{mAB}$	each 90° given (Hyp)	
$\overline{mAD} = \overline{mAD}$ $\Delta ABD = \Delta ACD$	common Hyp side = Hyp side	
$\overline{mBD} = \overline{mCD} = \frac{28}{2}$ $= 14 \text{ cm}$		
Now in right angled ΔADC $\overline{mCD} = 14 \text{ cm}$ $\overline{AC} = 50$	given (Hyp)	
Now $(\overline{mAD})^2 = (\overline{mAC})^2 - (\overline{mCD})^2$ $(50)^2 - (14)^2$ $= 2500 - 196$ $= 2304$		
Hence $\overline{mAD} = 18$	Taking sq root	
Part (i) Area $\Delta ABC = \frac{\text{Base} \times \text{Alt}}{2}$ $\frac{28 \times 48}{2}$ $= 14 \times 48 = 672 \text{ Sq.cm}$		

Q.5. In a quadrilateral ABCD, the diagonals \overline{AC} and \overline{BD} are perpendicular to each other.

Prove that $\overline{mAB}^2 + \overline{mCD}^2 = \overline{mAD}^2 + \overline{mBC}^2$



MATHEMATICS FOR 9TH CLASS (UNIT # 15)

Pilot Superone Mathematics 534 Class 9th

Given ABCD is a quadrilateral in which its diagonals \overline{AC} and \overline{BD} are perpendicular to each other

Required: $(mAB)^2 + (mCD)^2 = (mAD)^2 + (mBC)^2$


Proof

Statements	
Let $\overline{OC} = l$ $\overline{OD} = m$ $\overline{OA} = n$ $\overline{OB} = p$	
In right angled $\triangle OCD$ $(mCD)^2 = l^2 + m^2$ (i)	
In right angled $\triangle ABO$ $(mAB)^2 = n^2 + p^2$	
Now $(mAB)^2 + (mCD)^2 = l^2 + m^2 + n^2 + p^2$	from (i) + (ii) (L)
In right angled $\triangle AOD$ $(mAD)^2 = n^2 + m^2$ (iii)	
In right angled $\triangle BOC$ $(mBC)^2 = p^2 + l^2$ (iv)	
Now $(mAD)^2 + (mBC)^2 = n^2 + m^2 + p^2 + l^2$	From iii + iv (M)
Thus $(mAB)^2 + (mCD)^2 = (mAD)^2 + (mBC)^2$	from L and M

Q.4. (i) In the $\triangle ABC$ as shown in the figure,

$m\angle ACB = 90^\circ$ and $CD \perp AB$. Find the lengths a , b and

h if $mBD = 5$ units and $mAD = 7$ units



Given: In $\triangle ABC$

(i) $m\angle C = 90^\circ$

MATHEMATICS FOR 9TH CLASS (UNIT # 15)

Pilot Supercode Mathematics 535

(ii) $\overline{CD} \perp \overline{AB}$
 $m\overline{BD} = 5, m\overline{AD} = 7$

Required: To find measures of a, b and h
 Calculations: In right angled triangle ABC

$$(m\overline{AB})^2 = a^2 + b^2$$

$$(7+5)^2 = a^2 + b^2$$

$$(12)^2 = a^2 + b^2$$

$$a^2 + b^2 = 144 \quad \dots (i)$$

or

In right angled triangle CDB

$$a^2 = (h)^2 + (5)^2$$

$$a^2 - h^2 = 25 \quad \dots (ii)$$

or

In right-angled triangle ACD

$$b^2 = h^2 + (7)^2$$

$$b^2 - h^2 = 49 \quad \dots (iii)$$

or

$$a^2 - b^2 = -24 \quad \dots (iv)$$

$$2a^2 = 144 - 24$$

$$2a^2 = 120$$

$$a^2 = 60$$

$$a = \sqrt{60}$$

$$= 2\sqrt{15} \quad \dots (v)$$

$$a^2 + b^2 = 144$$

$$b^2 = 144 -$$

$$= 144$$

$$b^2 =$$

Now

Now

www.dow

MATHEMATICS FOR 9TH CLASS (UNIT # 15)

Superior Mathematics

536

Class 9th

or $h^2 = 35$
 $h = \sqrt{35}$
 6 (ii) Find the value of x in the shown figure.



Given. In $\triangle ADC$ angle $m\angle ADC = 90^\circ$
 $(mAD)^2 = (13)^2 - (5)^2$ (Pythagoras theorem)
 $= 169 - 25$
 $= 144$ (i)
 $mAD = \sqrt{144}$
 $= 12 \text{ cm}$

Now in right angle triangle ABD
 angle ADB is 90°

$$x^2 = (15)^2 - (mAD)^2$$

$$x^2 = (15)^2 - 144$$

$$= 225 - 144$$

$$81$$


$$\sqrt{81}$$

$$9$$

A height of 300 m and is 500 m away
 from as shown in the figure. How much
 travel to land at the airport?

MATHEMATICS FOR 9TH CLASS (UNIT # 15)

Pilot Superone Mathematics 537 Class 9th



Solution: ABC is a right-angled triangle at C.

$$(\overline{AB})^2 = (\overline{AC})^2 + (\overline{BC})^2 \text{ (Pythagorean Theorem)}$$


$$\begin{aligned} & (300)^2 + (500)^2 \\ &= 90000 + 250000 \end{aligned}$$

$$(\overline{AB})^2 = 340000$$

$$\begin{aligned} \overline{AB} &= \sqrt{340000} \\ &= \sqrt{34 \times 100 \times 100} \end{aligned}$$

$$\overline{AB} = 100\sqrt{34} \text{ m}$$

- B. A ladder 17m long rests against a vertical wall. The foot of the ladder is 8 m away from the base of the wall. How high up the wall will the ladder reach?



Solution. Let \overline{AB} be the ladder

Now ABC is a right-angled triangle at C

MATHEMATICS FOR 9TH CLASS (UNIT # 15)

Pilot Superone Mathematics 538 Class 9th

$$(mAC')^2 = (mAB)^2 + (mBC')^2 \text{ (Pythagoras Theorem)}$$


$$\begin{aligned} \text{Now } (mAC')^2 &= (17)^2 + (8)^2 \\ &= 289 + 64 \\ &= 353 \end{aligned}$$

$$mAC' = \sqrt{353} \text{ Taking sq root}$$

$$mAC' = 18.8 \text{ m}$$

The ladder reaches upto 18.8m high.

9. A student travels to his school by the route as shown in the figure. Find mAD , the direct distance from his house to school.



Solution:

$$\begin{aligned} \text{Let } mBP &= x \text{ km} \\ mPC &= (13 - x) \text{ km} \end{aligned}$$

In $\triangle PCD \sim \triangle PBA$

$$\begin{aligned} \frac{x}{13-x} &= \frac{12}{5} \\ 5x &= 12(13-x) \\ 5x &= 156 - 12x \end{aligned}$$

$$x = \frac{12}{17} = mBP$$

MATHEMATICS FOR 9TH CLASS (UNIT # 15)

Pilot Superone Mathematics 539 Class 9th

$$m\overline{PC} = 6 - \frac{12}{5} = \frac{18}{5}$$

$$\text{In } \triangle PCD, m\overline{PD}^2 = m\overline{PC}^2 + m\overline{CD}^2 \text{ in}$$

$$= \left(\frac{18}{5}\right)^2 + (3)^2$$

$$= \frac{324}{25} + 9 = \frac{324 + 225}{25}$$

$$= \frac{549}{25}$$

$$m\overline{PD} = \sqrt{\frac{549}{25}} = \sqrt{\frac{3 \times 3 \times 61}{5 \times 5}}$$

$$= \frac{3}{5} \sqrt{61} \text{ km}$$

REVIEW EXERCISE 15

Which of the following are true and which are false?

- (i) In a right angled triangle greater angle is of 90°
- (ii) In a right angled triangle, right angle is of 60°
- (iii) In a right triangle hypotenuse is a side opposite to right angle.
- (iv) If a, b, c are sides of right angled triangle with c as longer side then $c^2 = a^2 + b^2$
- (v) If 3 cm and 4 cm are two sides of a right angled triangle, then hypotenuse is 5 cm.
- (vi) If hypotenuse of an isosceles right triangle is $\sqrt{2}$ cm then each of the other side is of length 2 cm

Answers:

- (i) T (ii) F (iii) T
- (iv) T (v) T (vi) F





2. Find the unknown value in each of the following figures.

MATHEMATICS FOR 9TH CLASS (UNIT # 15)

Pilot Superone Mathematics

540

Class 9th



Solutions: In right angled $\triangle PQR$

2(i) $mPR^2 = mPQ^2 + mQR^2$ (Pythagoras Theorem)

$x^2 = (4)^2 + (3)^2$ Putting values

$= 16 + 9$

$x^2 = 25$

$x = \sqrt{25} = 5 \text{ cm}$

2(ii) In right angled triangle ABC (Pythagoras Theory)

$mAC^2 = mAB^2 + mBC^2$ (Pythagoras Theorem)

$(10)^2 = (6)^2 + x^2$ Putting values

$100 = 36 + x^2$

$x^2 = 100 - 36 = 64$

$x = \sqrt{64} = 8 \text{ cm}$

MATHEMATICS FOR 9TH CLASS (UNIT # 15)

Pilot Superone Mathematics 541 Class 9th

2(iii) In right angled ΔABC

$$(mAC)^2 = (mAB)^2 + (mBC)^2 \text{ (Pythagoras Theorem)}$$

$$(13)^2 = x^2 + (5)^2 \text{ Putting values}$$

$$169 = x^2 + 25$$

$$x^2 = 169 - 25 = 144$$

$$x = \sqrt{144} = 12 \text{ cm (Taking sq. root)}$$

2(iv) In right angled ΔPQR

$$(mPR)^2 = (mPQ)^2 + (mQR)^2 \text{ (Pythagoras Theorem)}$$

$$(\sqrt{2})^2 = x^2 + (1)^2 \text{ Putting values}$$

$$2 = x^2 + 1$$

$$x^2 = 2 - 1 = 1$$

$$x = \sqrt{1} \text{ cm (Taking sq. root)}$$

$$x = 1 \text{ cm}$$

MATHEMATICS FOR 9TH CLASS (UNIT # 16)

Past Superone Mathematics 242 Class 9th

**Unit
16**


THEOREMS RELATED WITH AREA

Area of a Figure

The region enclosed by the bounding lines of a closed figure is called the Area of the figure.

Triangular Region

The interior of a triangle is the part of the plane enclosed by the triangle




A triangular region is the union of a triangle and its interior

Area of a triangle means the area of its triangular region.

Rectangular Region

The interior of a rectangle is the part of the plane enclosed by the rectangle



A rectangular region is the union of a rectangle and its interior

Area of a rectangular region is the product of the length and width of the region.

MATHEMATICS FOR 9TH CLASS (UNIT # 16)

Pilot Superone Mathematics 543 Class 9th

Altitude of a parallelogram

If one side of a parallelogram is taken as its base, the perpendicular distance between that side and the side parallel to it is called the Altitude or Height of the parallelogram.

Altitude of a Triangle


If one side of a triangle is taken as the base, the perpendicular to that side from the opposite vertex is called the Altitude or Height of the triangle.

Theorem

Parallelograms on the same base and between the same parallel lines (or of the same altitude) are equal in area.

Given

Two parallelograms ABCD and ABEF having the same base AB and between the same parallel lines AB and DE.



To prove

area of parallelogram ABCD = area of parallelogram ABEF

Proof

Statement	Reason
In $\triangle EBC \leftrightarrow \triangle FAD$	
$\overline{CB} \cong \overline{DF}$	opp Sides of \parallel^m ABCD
$\overline{BE} \cong \overline{AF}$	opp Sides of \parallel^m ABEF
$\angle 1 \cong \angle 2$	$\overline{BC} \parallel \overline{AD}, \overline{BE} \parallel \overline{AF}$
$\triangle EBC \cong \triangle FAD$	
Area of $\triangle EBC$ = Area of $\triangle FAD$	


MATHEMATICS FOR 9TH CLASS (UNIT # 16)

Prior Superone Mathematics 544 Class 9th

Also area of quad ABED = Area of quad ABED	(same quad) (ii)
Adding (i) and (ii)	
$\Delta EBC + \text{area quad ABED} = \Delta FAD + \text{area quad ABED}$	
Hence $\text{Area } ABCD = \text{Area } EFGH$	
Q.E.D	

Theorem

Parallelograms on equal bases and having the same (or equal) altitude are equal in area.



Given

Parallelograms ABCD, EFGH are on the equal bases \overline{BC} , \overline{FG} , having equal altitudes.

To Prove

area of (parallelogram ABCD) = area of (parallelogram EFGH)

Construction

Place the parallelograms ABCD and EFGH so that their equal bases \overline{BC} , \overline{FG} are in the straight line BCFG. Join

\overline{AC} and \overline{EH}

Proof


Statements	Reasons
The given $\text{Area } ABCD$ and $EFGH$ are between the same parallels	Their altitudes are equal (given)
Hence ADEH is a straight line	
\parallel to \overline{BC}	
$m\angle ACB = m\angle EFG$	Given
$= m\angle EHF$	EFGH is a parallelogram

MATHEMATICS FOR 9TH CLASS (UNIT # 16)

Pilot Superone Mathematics	545	Class 9 th
Now $m\angle BCL = m\angle EHL$ and they are \perp		
BL and CH are both equal and \parallel		
Hence $EBCH$ is a parallelogram		A quadrilateral with two opposite sides congruent and parallel is a parallelogram
Now $\text{area}(\square ABCD) = \text{area}(\square EBCH)$ (i)		Being on the same base BC and between the same parallels
But $\text{area}(\square EBCH) = \text{area}(\square EFGH)$ (ii)		Being on the same base EH and between the same parallels
Hence $\text{area}(\square ABCD) = \text{area}(\square EFGH)$		From (i) and (ii)

EXERCISE 16.1

Q.1 Show that the line segment joining the mid-points of opposite sides of a parallelogram, divides it into two equal parallelograms.



Given:

- (i) $ABCD$ is a \square
- (ii) P is mid point of \overline{DC} i.e. $\overline{DP} \cong \overline{PC}$
- (iii) Q is mid point of \overline{AB} i.e. $\overline{AQ} \cong \overline{QB}$
- (iv) P is joined to Q .

Required:

$\square AQPQ \cong \square QBCP$

MATHEMATICS FOR 9TH CLASS (UNIT # 16)

Pilot Superone Mathematics

546

Class 9th

Construction:

Join Q to C.


Proof:

Statements	Reasons
$m\overline{AB} = m\overline{DC}$	Opp. sides of \parallel^m ABCD
$\frac{1}{2}m\overline{AB} = \frac{1}{2}m\overline{DC}$	Dividing by 2
or $m\overline{QB} = m\overline{PC}$	
Now in $\triangle PQC$ $\triangle QBC$	Common
$\overline{QC} \cong \overline{QC}$	Proved
$\overline{QB} \cong \overline{PC}$	Alternate angles $\overline{AB} \parallel \overline{DC}$
$\angle 3 \cong \angle 4$	S.A.S \cong S.A.S.
$\triangle QBC \cong \triangle PQC$	Corresponding sides of cong. triangles
(i) $\overline{PQ} \cong \overline{CB}$	Opp. side of \parallel^m
But (ii) $\overline{AD} \cong \overline{CB}$	Corresponding angles of cong. Triangles
Thus $\overline{PQ} \cong \overline{AD} \cong \overline{BC}$	
And $\angle 1 \cong \angle 2$	
or $m\angle 1 + m\angle 3 = m\angle 2 + m\angle 4$	
or $\angle PQB \cong m\angle PCB$	Given
and $\angle A \cong m\angle PCB$	Proved
or $\angle A \cong m\angle PQB$	Proved
Now in $\square QBCP = \square PQDA$	
$\overline{AQ} \cong \overline{QB}$	
$\overline{AD} \cong \overline{PQ}$	
$\angle A \cong \angle PQB$	
Therefore $\square AQPD \cong \square QBCP$	

MATHEMATICS FOR 9TH CLASS (UNIT # 16)

Pilot Supertone Mathematics 547 Class 9th

Q.2 In a parallelogram $ABCD$, $m\overline{AB} = 10\text{cm}$. The altitudes corresponding to sides AB and AD are respectively 7 cm and 8 cm. Find $m\overline{AD}$



Given:

$ABCD$ is a \parallel^m in which.

(i) $m\overline{AB} = 10\text{ cm}$

(ii) Alt \overline{DP}

Required:

To find $m\overline{AD}$

Solution:

In $\parallel^m ABCD$ -

Base = 10 cm

Alt = 7 cm

$$\begin{aligned}\text{Area of } \parallel^m ABCD &= 10 \times 7 \\ &= 70 \text{ sq cm} \quad (i)\end{aligned}$$

In second case:

Base = \overline{AD}

Alt $\overline{BQ} = 8\text{ cm}$.

$$\text{Area of } \parallel^m ABCD = (\text{Base}) \times (m\overline{AD})$$

Now


$$70 = 8 \times (m\overline{AD}) \quad \text{from (i)}$$

$$m\overline{AD} = \frac{70}{8} = \frac{35}{4} \text{ cm}$$

MATHEMATICS FOR 9TH CLASS (UNIT # 16)

Pilot Superone Mathematics 548 Class 9th

Q.3 If two parallelograms of equal areas have the same or equal bases, their altitudes are equal.



Given:

In \parallel^m ABCD and PQRS

Area \parallel^m ABCD = Area \parallel^m PQRS

And Base AB = Base PQ

Alt. of ABCD is \overline{DM}

Alt. of PQRS is \overline{SN}

Required:

$m\overline{DM} = m\overline{SN}$

Proof:


Statements	Reasons
Area ABCD \parallel^m = Area PQRS \parallel^m Now $(mAB)(mDM) = (mPQ)(mSN)$ $(mAB)(mDM) = (mAB)(mSN)$ $m\overline{DM} = m\overline{SN}$	Area of \parallel^m = Base \times Alt. $mAB = mPQ$ (given)

MATHEMATICS FOR 9TH CLASS (UNIT # 16)

Pilot Superone Mathematics 549 Class 9th

Theorem

Triangles on the same base and of the same (i.e. equal) altitudes are equal in area.



Given. ΔABC , ΔDBC on the same base \overline{BC} , and having equal altitudes.

To Prove: area of (ΔABC) = area of (ΔDBC)

Construction: Draw $BM \parallel$ to CA , $CN \parallel$ to BD AD meeting produced in M , N .

Proof:


Statements	Reasons
ΔABC and ΔDBC are between the same \parallel^s	Their altitudes are equal
Hence $MADN$ is parallel to \overline{BC}	
Area (\parallel^m $BCAM$) = Area (\parallel^m $BCND$) (i)	These \parallel^m are on the same base \overline{BC} and between the same \parallel^s
But $\Delta ABC = \frac{1}{2} (\parallel^m \text{ } BCAM)$ (ii)	Each diagonal of a \parallel^m bisects it into two congruent triangles
And $\Delta DBC = \frac{1}{2} (\parallel^m \text{ } BCND)$ (iii)	
Hence Area (ΔABC) = Area (ΔDBC)	From (i), (ii) and (iii)

MATHEMATICS FOR 9TH CLASS (UNIT # 16)

Pilot Superone Mathematics 550 Class 9th

Theorem

Triangle on equal bases and of equal altitudes are equal in area.



Given: Δ s ABC, DEF on the same base \overline{BC} , \overline{EF} and having altitudes equal.

To Prove: Area of (Δ ABC) = Area of (Δ DEF)

Construction:

Place the Δ ABC and DEF so that their equal bases \overline{BC} and \overline{EF} are in the same straight line BCEF and their vertices on the same side of it. Draw $BX \parallel$ to CA and $FY \parallel$ to ED meeting AD produced in X, Y respectively.

Proof:

Statements	Reasons
Δ ABC and Δ DEF are between the same parallels	Their altitudes are equal (given)
$XADY$ is \parallel to BCEF	
Area (\parallel^m BCAX) = Area (\parallel^m EFYD) (i)	These \parallel^m are on equal bases and between the same parallels
But $\Delta ABC = \frac{1}{2}$ (\parallel^m BCAX) (ii)	Diagonal of a \parallel^m bisects it
And $\Delta DEF = \frac{1}{2}$ (\parallel^m EFYD) (iii)	
Hence Area (Δ ABC) = Area (Δ DEF)	From (i), (ii) and (iii)


MATHEMATICS FOR 9TH CLASS (UNIT # 16)

Pilot Superone Mathematics 551

Class 9th

EXERCISE 16.2

Q.1 Show that a median of a triangle divides it into two triangles of equal area.



Given: In $\triangle ABC$: \overline{AM} is a median i.e. $m\overline{BM} = m\overline{MC}$

Required: Area $\triangle AMC$ = Area $\triangle ABM$

Proof:


Statements	Reasons
Area of $\triangle ABM = \frac{m\overline{AP} \times m\overline{BM}}{2}$ (i)	
Area of $\triangle AMC = \frac{m\overline{AP} \times m\overline{MC}}{2}$	
$= \frac{m\overline{AP} \times m\overline{BM}}{2}$ (ii)	$m\overline{BM} = m\overline{MC}$ (given)
Area of $\triangle ABM$ = Area of $\triangle AMC$	
Similarly it can be proved that every median of a triangle divides it into two triangles equal in area.	



MATHEMATICS FOR 9TH CLASS (UNIT # 16)

Pilot Superone Mathematics SS2 Class 9th

Q.2 Prove that a parallelogram is divided by its diagonals into four triangles of equal area.



- Given:** (i) ABCD is a \parallel^m
 (ii) \overline{AC} and \overline{BD} are its diagonals that intersect each other at O.

Required: $\Delta OBC = \Delta OCD = \Delta OAD = \Delta OAB$

Construction:

- (i) Drop perpendicular from C to \overline{DB} i.e. \overline{CN}
 (ii) Drop perpendicular from A to \overline{DB} i.e. \overline{AM}


Proof:

Statements	Reasons
$m\overline{OB} = m\overline{OD}$ $m\overline{OA} = m\overline{OC}$ $\Delta OBC = \frac{m\overline{OB} \times m\overline{CN}}{2}$ (i)	Diagonals of a \parallel^m bisect each other
And $\Delta OAD = \frac{m\overline{OD} \times m\overline{CN}}{2}$ $= \frac{m\overline{OB} \times m\overline{CN}}{2}$ (ii)	
$\Delta OBC = \Delta OAD$	
Similarly $\Delta OAD = \Delta OAB$ $\Delta OBC = \Delta OCD = \Delta OAD = \Delta OAB$	
	From (i) and (ii)

MATHEMATICS FOR 9TH CLASS (UNIT # 16)

Pilot Superone Mathematics 553 Class 9th

Q.3 Divide a triangle into six equal triangular parts.



Given: $\triangle ABC$ is a triangle

Required: To divide $\triangle ABC$ into six equal triangular parts

Construction:

- (i) Take \overline{BP} any ray making an acute angle with \overline{BC}
- (ii) Draw six arcs of the same radius on \overline{BP} such that $mBd = mde = mef = mfg = mgh = mhc$
- (iii) Join c To C and parallel lines are drawn through d, e, f, g, h to BC such that $cC \parallel dH \parallel eG \parallel fF \parallel gE \parallel hC$
- (iv) Join A to O, E, F, G, H

Proof:

Base \overline{BC} of $\triangle ABC$ has been divided to equal parts six in number

We get six triangles having equal base and same

Altitude

Their area is equal

Hence $\triangle BOA = \triangle OEA = \triangle EFA = \triangle FGA = \triangle GHA = \triangle HCA$

Visit

MATHEMATICS FOR 9TH CLASS (UNIT # 16)

Pilot Superone Mathematics 554 Class 9th

Review Exercise 16

Q.1 Which of the following are true and which are false?

- (i) Area of a figure means region enclosed by bounding lines of closed figure.
- (ii) Similar figures have same area.
- (iii) Congruent figures have same area.
- (iv) A diagonal of a parallelogram divides it into two non-congruent triangles.
- (v) Altitude of a triangle means perpendicular from vertex to the opposite side (base).
- (vi) Area of a parallelogram is equal to the product of base and height.

Answers:

(i)	T	(ii)	F	(iii)	T	(iv)	F	(v)	T
(vi)	T								

Q.2 Find the area of the following figures.


(i)




(ii)



(iii)



(iv)



Answers:

2(i)

Length of the rectangle = 6 cm
 Width of the rectangle = 3 cm
 Area of the rectangle = 6×3

MATHEMATICS FOR 9TH CLASS (UNIT # 16)

Pilot Superons Mathematics 555 Class 9th

- 2(ii) Length of side of square = 18 sq. cm
 = 4 cm
 Area = (side)²
 = (4)²
 = 16 sq. cm
- 2(iii) Length of side of a ||^m = 8 cm
 Act of the ||^m = 4
 Area = 8 × 4
 = 32 sq. cm
- 2(iv) Base of the triangle = 16 cm
 Alt = 10 cm
 Area of the Δ = $\frac{16 \times 10}{2}$
 = 80 sq. cm

Q.3 Define the following

- (i) Area of a figure (ii) Triangular Region
 (iii) Rectangular Region
 (iv) Altitude or Height of a triangle

Solution:

3(i) Area of a figure

Area of a figure means region enclosed by the boundary lines of a closed figure.

3(ii) Triangular Region

A triangular region means the union of triangle and its interior

3(iii) Rectangular Region

A rectangular region is the union of a rectangle and its interior

3(iv) Altitude or Height of a triangle

Altitude or height of a triangle means perpendicular distance to base to base from its opposite vertex.

W11



MATHEMATICS FOR 9TH CLASS (UNIT # 17)

Pilot Superone Mathematics 556 Class 9th

Unit
17

PRACTICAL GEOMETRY


EXERCISE 17.1

Q.1 Construct a $\triangle ABC$ in which

- (i) $m\overline{AB} = 3.2 \text{ cm}$, $m\overline{BC} = 4.2 \text{ cm}$, $m\overline{CA} = 5.2 \text{ cm}$
- (ii) $m\overline{AB} = 4.2 \text{ cm}$, $m\overline{BC} = 3.9 \text{ cm}$, $m\overline{CA} = 3.6 \text{ cm}$
- (iii) $m\overline{AB} = 4.8 \text{ cm}$, $m\overline{BC} = 3.7 \text{ cm}$, $m\angle B = 60^\circ$
- (iv) $m\overline{AB} = 3 \text{ cm}$, $m\overline{AC} = 3.2 \text{ cm}$, $m\angle A = 45^\circ$
- (v) $m\overline{BC} = 4.2 \text{ cm}$, $m\overline{CA} = 3.5 \text{ cm}$, $m\angle C = 75^\circ$
- (vi) $m\overline{AB} = 2.5 \text{ cm}$, $m\angle A = 30^\circ$, $m\angle B = 105^\circ$
- (vii) $m\overline{AB} = 3.6 \text{ cm}$, $m\angle A = 75^\circ$, $m\angle B = 45^\circ$

1(i) Construct a $\triangle ABC$ in which

$m\overline{AB} = 3.2 \text{ cm}$, $m\overline{BC} = 4.2 \text{ cm}$, $m\overline{CA} = 5.2 \text{ cm}$



Given.

$m\overline{AB} = 3.2 \text{ cm}$

$m\overline{BC} = 4.2 \text{ cm}$

$m\overline{CA} = 5.2 \text{ cm}$

Required: Construct a $\triangle ABC$

Step of Construction:

- (i) Take a line segment $\overline{CA} = 5.2 \text{ cm}$

MATHEMATICS FOR 9TH CLASS (UNIT # 17)

=====

www.downloadclassnotes.com

MATHEMATICS FOR 9TH CLASS (UNIT # 17)


Pilot Superone Mathematics 557 Class 9th

- (ii) Taking C as centre and draw an arc of radius 4.2 cm.
- (iii) Taking A as centre and draw another arc of radius 3.2 cm. It intersect the first arc at B
- (iv) Join C to A and B.

Result. ABC is the required triangle

- 1(ii) Construct a $\triangle ABC$ in which**

$$m\overline{AB} = 4.2 \text{ cm}, m\overline{BC} = 3.9 \text{ cm}, m\overline{CA} = 3.6 \text{ cm}$$



Given. $m\overline{AB} = 4.2 \text{ cm}$
 $m\overline{BC} = 3.9 \text{ cm}$
 $m\overline{CA} = 3.6 \text{ cm}$

Required: Construct a $\triangle ABC$

Steps of Construction:

- (i) Take a line segment \overline{AB} of length 4.2 cm.
- (ii) Take A as centre and draw an arc of 3.6 cm radius.
- (iii) Take B as centre and draw an arc of 3.9 cm radius. This cuts the first arc at C.
- (iv) Join C to A, B


Result: ABC is the required triangle.

MATHEMATICS FOR 9TH CLASS (UNIT # 17)

Pilot Superone Mathematics 558 Class 9th

1(iii) Construct a $\triangle ABC$ in which

$$m\overline{AB} = 4.8 \text{ cm}, m\overline{BC} = 3.7 \text{ cm}, m\angle B = 60^\circ$$



Given: $m\overline{AB} = 4.8 \text{ cm}$
 $m\overline{BC} = 3.7 \text{ cm}$
 $m\angle B = 60^\circ$

Required: Construct triangle ABC

Steps of Construction:

- (i) Take a line segment $\overline{AB} = 4.8 \text{ cm}$.
- (ii) Make an angle of 60° at B
- (iii) Cut off $\overline{BC} = 3.7 \text{ cm}$ from \overline{BP}
- (iv) Join C to A.


Result. $\triangle ABC$ is the required triangle.

MATHEMATICS FOR 9TH CLASS (UNIT # 17)

Pilot Superone Mathematics 55 ■ Class 9th

1(iv) Construct a triangle ABC in which

$$m\overline{AB} = 3 \text{ cm}, m\overline{AC} = 3.2 \text{ cm}, m\angle A = 45^\circ$$



Given: $m\overline{AB} = 3 \text{ cm}$

$$m\overline{AC} = 3.2 \text{ cm}$$

$$m\angle A = 45^\circ$$

Required: Construct a triangle ABC

Steps of Construction.

- (i) Take a line segment $\overline{AB} = 3 \text{ cm}$.
- (ii) Make an angle of 45° at A.
- (iii) Take A as centre and cut off $\overline{AC} = 3.2 \text{ cm}$ on \overline{AR} .
- (iv) Join C to B.

Result: ABC is the required triangle.

1(v) Construct a $\triangle ABC$ in which

$$m\overline{BC} = 4.2 \text{ cm}, m\overline{CA} = 3.5 \text{ cm}, m\angle C = 75^\circ$$

MATHEMATICS FOR 9TH CLASS (UNIT # 17)

Pilot Super one Mathematics 560 Class 9th



Given: $m\widehat{BC} = 4.2 \text{ cm}$
 $m\widehat{CA} = 3.5 \text{ cm}$
 $m\angle C = 75^\circ$

Required: Construct a $\triangle ABC$


Steps of Construction.

- Take a line segment $\widehat{CA} = 3.5 \text{ cm}$.
- Make an angle of 75° at C.
- Cut off $\widehat{CB} = 4.2 \text{ cm}$ from \widehat{CP}
- Join B to A

Result: $\triangle ABC$ is the required triangle.

1(vi) Construct a $\triangle ABC$ in which

$m\widehat{AB} = 2.5 \text{ cm}$, $m\angle A = 30^\circ$, $m\angle B = 105^\circ$



Given: $m\widehat{AB} = 2.5 \text{ cm}$

MATHEMATICS FOR 9TH CLASS (UNIT # 17)

Pilot Superiore Mathematics 561 Class 9th

$$m\angle A = 30^\circ$$

$$m\angle B = 105^\circ$$

Required: Construct a triangle ABC


Steps of Construction:

- (i) Take a line segment $\overline{AB} = 2.5$ cm
- (ii) Make an angle of 45° at A.
- (iii) Make an angle of 105° at B
- (iv) \overline{AP} , \overline{BQ} intersect at C

Result. ABC is the required triangle

1(vii) Construct a $\triangle ABC$ in which

$$m\overline{AB} = 3.6 \text{ cm}, m\angle A = 75^\circ, m\angle B = 45^\circ$$



Given: $m\overline{AB} = 3.6$ cm

$$m\angle A = 75^\circ$$

$$m\angle B = 45^\circ$$

Required: Construct a triangle ABC

Steps of Construction:

- (i) Take a line segment $\overline{AB} = 3.6$ cm
- (ii) Make an angle of 75° at A.
- (iii) Make an angle of 45° at B
- (iv) \overline{AR} and \overline{BP} intersect at C

MATHEMATICS FOR 9TH CLASS (UNIT # 12)

Plot Superone Mathematics 562 Class 9


Result. ABC is the required triangle

Q.2 Construct a ΔXYZ in which.

- (i) $mYZ = 7.6$ cm, $mXY = 6.1$ cm, $m\angle X = 90^\circ$
- (ii) $mZX = 6.4$ cm, $mYZ = 2.4$ cm, $m\angle Y = 90^\circ$
- (iii) $mXY = 5.5$ cm, $m\angle X = 4.5$ cm, $m\angle Z = 90^\circ$

2(i) Construct a ΔXYZ in which.

$mYZ = 7.6$ cm, $mXY = 6.1$ cm, $m\angle X = 90^\circ$



Given: $mYZ = 7.6$ cm
 $mXY = 6.1$ cm
 $m\angle X = 90^\circ$

Required: Construct a triangle XYZ

Steps of Construction.

- (i) Take a line segment $XY = 6.1$ cm
- (ii) Make an angle of XPY of 90° at X
- (iii) Take Y as centre and draw an arc of radius of 7.6 cm. This arc intersect XP at Z
- (iv) Join Z to Y

Result. XYZ is the required triangle


MATHEMATICS FOR 9TH CLASS (UNIT # 17)

Pilot Superone Mathematics 563

Class 9th

2(ii) Construct a $\triangle XYZ$ in which,

$$m\angle X = 64^\circ, m\angle Y = 24^\circ, m\angle Z = 90^\circ$$



Given $m\angle X = 64^\circ$
 $m\angle Y = 24^\circ$
 $m\angle Z = 90^\circ$

Required: To construct $\triangle XYZ$

Steps of Construction:

- Take a line segment $YZ = 2.4$ cm
- Make an angle of 90° at Y .
- Take Z as centre and draw an arc of radius 6.4 cm which cuts $Y\bar{P}$ at X .
- Join X to Z .


Result: $\triangle XYZ$ is the required triangle

MATHEMATICS FOR 9TH CLASS (UNIT # 17)

Pilot Superone Mathematics 564 Class 9

2(iii) Construct a $\triangle XYZ$ in which

$$m\overline{XY} = 5.5 \text{ cm}, m\overline{ZX} = 4.5 \text{ cm}, m\angle Z = 90^\circ$$



Given. $m\overline{XY} = 5.5 \text{ cm}$
 $m\overline{ZX} = 4.5 \text{ cm}$
 $m\angle Z = 90^\circ$

Required: Construct a $\triangle XYZ$

Steps of Construction:

- (i) Take a line segment $\overline{ZX} = 4.5 \text{ cm}$
- (ii) Draw an angle $\angle XZP$ of Z of 90°
- (iii) Take X as centre and draw an arc of 5.5 cm radius, which cuts \overline{ZP} at Y
- (iv) Join Y to X .


Result: $\triangle XYZ$ is the required triangle

✓

MATHEMATICS FOR 9TH CLASS (UNIT # 17)

Plot Superone Mathematics 565 Class 9th

Q.3 Construct a right angled triangle measure of whose hypotenuse is 5 cm and one side is 3.2 cm.



Given: Length of hypotenuse = 5 cm
One side = 3.2 cm

Required: To construct a right angled triangle.

Steps of Construction:

- (i) Take a line segment of 3.2 cm.
- (ii) Make an angle $\angle BAP = 90^\circ$ at A.
- (ii) Take B as centre and draw an arc of radius 5 cm which cuts \overrightarrow{AP} cuts at C
- (iii) Join C to A.


Result: ABC is the required triangle

Q.4 Construct a right angled isosceles triangle whose hypotenuse is

- | | |
|--------------|-------------------|
| (i) 5.2 cm | (ii) 4.8 cm |
| (iii) 6.2 cm | (iv) 5.4 cm long. |

MATHEMATICS FOR 9TH CLASS (UNIT # 17)

Expt. Superior Mathematics Page 566 Class 9th
 Q(i) 5.2 cm



Given: Length of hypotenuse = 5.2 cm

Required: To construct a right angled isosceles triangle

Steps of Construction.

- (i) Take a line segment AB = 5.2 cm
- (ii) Take D as midpoint by drawing \overleftrightarrow{PQ} as perpendicular bisector of AB
- (iii) Take O as centre and draw a semi circle of radius OA or OB which cuts \overleftrightarrow{PQ} at C
- (iv) Join C to A and B

Result: $\triangle ABC$ is the required isosceles right angled triangle at C

Q(ii) 4.8 cm



MATHEMATICS FOR 9TH CLASS (UNIT # 17)

Pilot Superone Mathematics 567 Class 9th

Given: Length of hypotenuse = 4.8 cm


Required: To construct a right angled isosceles triangle

Steps of Construction:

- (i) Take a line segment $AB = 4.8$ cm
- (ii) Draw PQ right bisector of AB which cuts AB at O
- (iii) Take O as centre and draw a semi-circle of radius \overline{OA} or \overline{OB} which cuts PQ at C
- (iv) Join C to A and B

Result: $\triangle ABC$ is the required triangle which is right angled at C and $m\angle C = m\angle B$.

4(ii) 6.2 cm



Given: Length of hypotenuse = 6.2 cm

Required: To construct a right angled isosceles triangle with hypotenuse = 6.2 cm

Steps of Construction:

- (i) Take a line segment $AB = 6.2$ cm
- (ii) Draw \hat{PQ} right bisector of AB which cuts AB at O

MATHEMATICS FOR 9TH CLASS (UNIT # 17)


Pilot Superone Mathematics 568 Class 9th

(iii) Take O as centre and draw a semi-circle of radius \overline{OA} or \overline{OB} which cuts \overline{PQ} at C

(iv) Join C to A and B

Result: ABC is the required triangle which right-angled at C and $m\widehat{AC} = m\widehat{BC}$

4(iv) 5.4 cm



Given: Length of hypotenuse = 5.4 cm

Required: To construct a right angled isosceles triangle

Steps of Construction:

- (i) Take a line segment $\overline{AB} = 5.4$ cm.
- (ii) Draw \overline{PQ} right bisector of \overline{AB} which cuts it at point O
- (iii) Take O as centre and draw a semi-circle of radius \overline{OA} or \overline{OB} which cuts \overline{PQ} at C
- (iv) Join C to A and B

Result: ABC is the required triangle which is right-angled at C and $m\widehat{AC} = m\widehat{BC}$

Q.5 (Ambiguous Case) Construct a ΔABC in which


- (i) $m\widehat{AC} = 4.2$ cm, $m\widehat{AB} = 5.2$ cm, $m\angle B = 45^\circ$
- (ii) $m\widehat{BC} = 2.5$ cm, $m\widehat{AB} = 5.0$ cm, $m\angle A = 30^\circ$

MATHEMATICS FOR 9TH CLASS (UNIT # 17)

Pilot Superone Mathematics 569 Class 9th

(iii) $m\overline{BC} = 5 \text{ cm}$, $m\overline{AC} = 3.5 \text{ cm}$, $m\angle B = 60^\circ$

5(i) $m\overline{AC} = 4.2 \text{ cm}$, $m\overline{AB} = 5.2 \text{ cm}$, $m\angle B = 45^\circ$



Given: $m\overline{AC} = 4.2 \text{ cm}$

$m\overline{AB} = 5.2 \text{ cm}$

$m\angle B = 45^\circ$

Required: To construct triangle ABC.

Steps of Construction:

- (i) Take a line segment $\overline{AB} = 5.2 \text{ cm}$
- (ii) Make an angle $\angle ABP = 45^\circ$ at point B
- (iii) Take point A as centre and draw an arc of 4.2 cm radius. It cuts \overline{BP} at two points C, C'
- (iv) Join C, C' to A

Result: Two Δ^s ABC, ABC' fulfill the requirements

5(ii) $m\overline{BC} = 2.5 \text{ cm}$, $m\overline{AB} = 5.0 \text{ cm}$, $m\angle A = 30^\circ$



Given: $m\overline{BC} = 2.5 \text{ cm}$

$m\overline{AB} = 5.0 \text{ cm}$

MATHEMATICS FOR 9TH CLASS (UNIT # 17)

Pilot Superone Mathematics 570 Class 9th

$m\angle A = 30^\circ$

Required: Construct a $\triangle ABC$


Steps of Construction:

- (i) Take a line segment $AB = 5$ cm.
- (ii) Make an angle $\angle BAP = 30^\circ$ at A.
- (iii) Take B as centre and draw an arc of 2.5 cm radius. This arc touches AP at C.
- (iv) Join C to A.

Result: $\triangle ABC$ is the required triangle.

5(ii) Construct a triangle ABC in which

$m\angle C = 5$ cm, $m\angle A = 3.5$ cm, $m\angle B = 60^\circ$



Given: $m\angle C = 5$ cm

$m\angle A = 3.5$ cm

$m\angle B = 60^\circ$

Required: Construct a triangle ABC

Steps of Construction:

- (i) Take a line segment $BC = 5$ cm
- (ii) Make an angle $\angle CBR$ of 60° at B
- (iii) Take C as centre and draw an arc of 3.5 cm radius. This arc does not cut and cut BR at any point

MATHEMATICS FOR 9TH CLASS (UNIT # 17)

Plot Superio in Mathematics 571 Class 9

Result: Triangle ABC cannot be constructed with the given sides and angle.


Remember:

- (i) The internal bisectors of the angles of a triangle meet at a point called the incentre of the triangle.
- (ii) The point of concurrency of the three perpendicular bisectors of the sides of a Δ is called the circumcentre of the Δ .
- (iii) The point of concurrency of the three altitudes of a Δ is called its orthocenter.
- (iv) The point where the three medians of a Δ meet is called the centroid of the triangle.

EXERCISE 17.2

Q 1 Construct the following ΔABC . Draw bisectors of their angles and verify their concurrency.

- (i) $mAB = 4.5 \text{ cm}$, $mBC = 3.1 \text{ cm}$ and $m\angle A = 52^\circ$
- (ii) $mAB = 4.2 \text{ cm}$, $mBC = 6 \text{ cm}$ and $m\angle A = 52^\circ$
- (iii) $mAB = 3.6 \text{ cm}$, $mBC = 4.2 \text{ cm}$ and $m\angle B = 75^\circ$
- (iv) $mAB = 4.5 \text{ cm}$, $mBC = 3.1 \text{ cm}$ and $m\angle A = 52^\circ$



Given: $mAB = 4.5 \text{ cm}$

MATHEMATICS FOR 9TH CLASS (UNIT # 17)

Pilot Superone Mathematics 572 Class 9th

$$m\widehat{BC} = 3.1\text{cm}$$

$$m\widehat{CA} = 5.2\text{cm}$$

Required: Construct $\triangle ABC$ and see the concurrency of the bisectors of its angles


Steps of Construct:

- (i) Take a line segment $\overline{AB} = 4.5\text{ cm}$
- (ii) Take A and B as centers and draw arcs of 5.2, 3.1cm radii respectively. These arcs intersect each other at point C
- (iii) Join C to A and B. $\triangle ABC$ is the required triangle.
- (iv) Take \overline{AP} , \overline{BQ} , \overline{CR} bisectors of angles A, B, C respectively.

Result:

We see that bisectors \overline{AP} , \overline{BQ} , \overline{CR} are concurrent at point ..

1(ii) $m\widehat{AB} = 4.2\text{ cm}$, $m\widehat{BC} = 6\text{ cm}$ and $m\widehat{CA} = 5.2\text{cm}$



Given:

$$m\widehat{AB} = 4.2\text{ cm}$$

$$m\widehat{BC} = 6\text{ cm}$$

$$m\widehat{CA} = 5.2\text{ cm}$$

MATHEMATICS FOR 9TH CLASS (UNIT # 17)

Pilot SuperOne Mathematics 573 Class 9th

Required: Construct $\triangle ABC$ and verify the concurrency of the bisectors of its angles.


Steps of Construction:

- (i) Take a line segment $\overline{BC} = 6$ cm
- (ii) Take B as center and draw an arc of 4.2 cm radius
- (iii) Take C as centre and draw an arc of 5.2 cm radius that cuts the first arc at point A.
- (iv) Join A to B and C.
 $\triangle ABC$ is the required triangle
- (v) Take \overline{AP} , \overline{BQ} and \overline{CR} bisectors of angle A, B and C respectively

Result:

\overline{AP} , \overline{BQ} , \overline{CR} are concurrent at point I

1(iii) $m\overline{AB} = 3.6$ cm, $m\overline{BC} = 4.2$ cm and $m\angle B = 75^\circ$



Given: $m\overline{AB} = 3.6$ cm
 $m\overline{BC} = 4.2$ cm
 $m\angle B = 75^\circ$

Required: Construct $\triangle ABC$ and see that bisectors of angles are concurrent.

W!



MATHEMATICS FOR 9TH CLASS (UNIT # 17)

Plot Supertone Mathematics 574 _____ Class 9th

Steps of Construct.


- (i) Take a line segment $\overline{BC} = 4.2$ cm
- (ii) Make an angle $\angle CBF = 75^\circ$
- (iii) Take B as centre and draw an arc of radius 3.6 cm which cuts BF at A
- (iv) Join A to C
ABC is the required triangle
- (v) Take \overline{AP} , \overline{BQ} and \overline{CR} bisectors of angles.

Result.

\overline{AP} , \overline{BQ} and \overline{CR} are concurrent at point I.

Q.2 Construct the following Δ 'PQR', draw their altitudes and show that they are concurrent.

- (i) $mPQ = 6$ cm, $mQR = 4.5$ cm, $mPR = 5.5$ cm
- (ii) $mPQ = 4.5$ cm, $mQR = 3.9$ cm, $m\angle R = 45^\circ$
- (iii) $mRP = 3.6$ cm, $m\angle Q = 30^\circ$, $m\angle P = 105^\circ$
- 2(ii) $mPQ = 6$ cm, $mQR = 4.5$ cm, $mPR = 5.5$ cm



Given: $mPQ = 6$ cm

MATHEMATICS FOR 9TH CLASS (UNIT # 17)

Pilot Superone Mathematics 575 Class 9th

$m\angle Q = 45^\circ$

$m\angle R = 55^\circ$

Required: Construct $\triangle PQR$ and take their altitudes.


Steps of Construction:

- (i) Take a line segment $PQ = 6$ cm
- (ii) Take P as centre and draw an arc of 5.5 radius
- (iii) Take Q as centre and draw an arc of 4.5 radius.
- (iv) Join R to P and Q.
PQR is the required triangle
- (v) Drop $RA \perp PB$ and $QC \perp PB$ perpendiculars to PQ ,
QR, RP respectively

Result:

$RA \perp PB$ and $QC \perp PB$ are concurrent at M

2(ii) $m\angle P = 45^\circ$, $m\angle Q = 39^\circ$, $m\angle R = 45^\circ$



Given: $m\angle P = 45^\circ$

$m\angle Q = 39^\circ$

$m\angle R = 45^\circ$

Required: Construct $\triangle PQR$ and draw its altitude

MATHEMATICS FOR 9TH CLASS (UNIT # 17)

Pilot Super one Mathematics 576 Class 9th

Steps of Construction:

- (i) Take $RQ = 3.9$ cm.
- (ii) Make an angle $\angle QRA = 45^\circ$
- (iii) Take Q as centre and draw an arc of 4.5 cm radius, which cuts \overline{RA} at P
- (iv) Join P to Q.
PQR is the required triangle
- (v) Drop \overline{QM} , \overline{RN} and \overline{PW} perpendiculars on \overline{RP} , \overline{PQ} and \overline{RQ} respectively

Result:

\overline{QM} , \overline{RN} and \overline{PW} are concurrent at O.

2.(iii) $m\overline{RP} = 3.6$ cm, $m\angle R = 30^\circ$, $m\angle P = 105^\circ$



Given: $m\overline{RP} = 3.6$ cm
 $m\angle R = 30^\circ$
 $m\angle P = 105^\circ$

Required: Construct ΔPQR and verify that its altitudes are concurrent

Steps of Construction:

MATHEMATICS FOR 9TH CLASS (UNIT # 17)

Pilot Superone Mathematics 577 Class 9th


- (i) Take a line segment $\overline{RP} = 3.6$ cm
- (ii) Draw an angle $\angle PRQ = 30^\circ$
- (iii) Draw an angle $\angle RPQ = 105^\circ$
 \overline{RM} and \overline{PW} intersect at point Q
 PQR in the required triangle
- (iv) Drop \overline{RB} , \overline{PA} and \overline{QC} perpendicular on \overline{QP} , \overline{PA} ,
 \overline{RP} respectively

Result:

\overline{RB} , \overline{PA} , \overline{QC} are concurrent at O.

Q.3 Construct the following triangles ABC. Draw the perpendicular bisector of their sides and verify their concurrency. Do they meet inside the triangle.

- (i) $m\overline{AB} = 5.3$ cm, $m\angle A = 45^\circ$, $m\angle B = 30^\circ$
- (ii) $m\overline{BC} = 2.9$ cm, $m\angle A = 30^\circ$, $m\angle B = 60^\circ$
- (iii) $m\overline{AB} = 2.4$ cm, $m\overline{AC} = 3.2$ cm, $m\angle A = 120^\circ$
- 3(i) $m\overline{AB} = 5.3$ cm, $m\angle A = 45^\circ$, $m\angle B = 30^\circ$



Given: $m\overline{AB} = 5.3$ cm
 $m\angle A = 45^\circ$

MATHEMATICS FOR 9TH CLASS (UNIT # 17)

Pilot Superone Mathematics 578 Class 9th

$$m\angle B = 30^\circ$$

Required: Construct $\triangle ABC$ Draw perpendicular bisectors of its sides and verify their concurrency

Steps of Construction:

- (i) Take a line segment $\overline{AB} = 5.3$ cm
- (ii) Make an angle $\angle BAW = 45^\circ$.
- (iii) Make an angle $\angle ABJ = 30^\circ$
 \overline{AW} , \overline{BJ} intersect at point C
 $\triangle ABC$ is the required triangle.
- (iv) Take \overline{PQ} , \overline{TS} , \overline{UV} right bisectors of \overline{AB} , \overline{BC} , \overline{CA} respectively

Result:

\overline{PQ} , \overline{TS} , \overline{UV} are concurrent at point M.

- 3(ii) $m\overline{BC} = 2.9$ cm, $m\angle A = 30^\circ$, $m\angle B = 60^\circ$



Given: $m\overline{BC} = 2.9$ cm
 $m\angle A = 30^\circ$
 $m\angle B = 60^\circ$

MATHEMATICS FOR 9TH CLASS (UNIT # 17)

Pilot Superone Mathematics 579 Class 9th

Required: Construct triangle ABC Draw perpendicular bisector of its sides and verify their concurrency


Steps of Construction:

- (i) Take a line segment $\overline{BC} = 2.9 \text{ cm}$
- (ii) Draw an angle $\angle CBW = 60^\circ$,
 Place of A is not known, we find $m\angle C$
 $m\angle C = 180 - m\angle B - m\angle A$
 $m\angle C = 180 - 60 - 30 = 90^\circ$
- (iii) Draw an angle $\angle BCN = 90^\circ$
 \overline{PW} , \overline{CN} intersect at point A
 ABC is the required triangle.
- (iv) Take \overline{PQ} , \overline{RS} , \overline{UV} the right bisectors of \overline{BC} , \overline{CA} , \overline{AB} sides.

Result:

\overline{PQ} , \overline{RS} , \overline{UV} are concurrent at point M.

3(iii) $m\overline{AB} = 2.4 \text{ cm}$, $m\overline{AC} = 3.2 \text{ cm}$, $m\angle A = 120^\circ$



Given: $m\overline{AB} = 2.4 \text{ cm}$
 $m\overline{AC} = 3.2 \text{ cm}$
 $m\angle A = 120^\circ$

MATHEMATICS FOR 9TH CLASS (UNIT # 17)

Pilot Superone Mathematics 580 Class 9th

Required: Construct a $\triangle ABC$ and draw perpendiculars of its sides and verify their concurrency

Steps of Construction:


- (i) Take a line segment $\overline{AB} = 2.4$ cm.
- (ii) Make an angle $\angle BAW = 120^\circ$ at A.
- (iii) Cut off $m\overline{AB} = 3.2$ cm from \overline{AW}
- (iv) Join C to B
ABC is the required triangle
- (v) Take \overline{PQ} , \overline{RS} , \overline{UV} right bisectors of \overline{AB} , \overline{BC} , \overline{CA} sides

Result:

\overline{PQ} , \overline{RS} , \overline{UV} are concurrent at point M.
 (out side the \triangle)

Q.4 Construct the following $\triangle XYZ$. Draw their three medians and show that they are concurrent.

- (i) $m\overline{YZ} = 4.1$ cm, $m\angle Y = 60^\circ$ and $m\angle X = 75^\circ$
- (ii) $m\overline{XY} = 4.5$ cm, $m\overline{YZ} = 3.4$ cm, $m\overline{ZX} = 5.6$ cm
- (iii) $m\overline{ZX} = 4.3$ cm, $m\angle X = 75^\circ$, $m\angle Y = 45^\circ$
- 4(i) $m\overline{YZ} = 4.1$ cm, $m\angle Y = 60^\circ$ and $m\angle X = 75^\circ$



MATHEMATICS FOR 9TH CLASS (UNIT # 17)

Pilot Superone Mathematics 581 Class 9th


Given: $m\overline{YZ} = 4.1 \text{ cm}$
 $m\angle Y = 60^\circ$
 $m\angle X = 75^\circ$

Required: Construct $\triangle XYZ$. Draw medians and verify their concurrency

Steps of Construct:

- (i) Take a line segment $\overline{YZ} = 4.1 \text{ cm}$
- (i.) Make an angle $\angle ZYW = 60^\circ$
- (ii) Make an angle $\angle YZS = 45^\circ$
 \overline{YW} and \overline{ZS} intersect at X.
 $\triangle XYZ$ is the required triangle.
- (iv) Find mid point of \overline{YZ} , \overline{ZX} , \overline{XY} as Q, P, R.
- (v) Join X to Q, Y to P, Z to R.
 \overline{XQ} , \overline{YP} and \overline{ZR} are medians. These are concurrent at O.

4(ii) $m\overline{XY} = 4.5 \text{ cm}$, $m\overline{YZ} = 3.4 \text{ cm}$, $m\angle X = 5.6 \text{ cm}$



Given: $m\overline{XY} = 4.5 \text{ cm}$
 $m\overline{YZ} = 3.4 \text{ cm}$
 $m\angle X = 5.6 \text{ cm}$

Required: Construct triangle XYZ. Draw medians and verify their concurrency

MATHEMATICS FOR 9TH CLASS (UNIT # 17)

Pilot Superone Mathematics 582 Class 9th

Steps of Construction:

- (i) Take a line segment $\overline{XY} = 4.5$ cm.
- (ii) Take X as centre and draw an arc of radius 5.6 cm
- (iii) Take Y as centre and draw an arcs of radius 3.4 cm
These arcs intersect at point Z
- (iv) Join Z to X and Y
 $\triangle XYZ$ is the required triangle
- (v) Find mid points P, Q, R of \overline{XY} , \overline{YZ} , \overline{ZX}
- (vi) Draw \overline{PZ} , \overline{QX} , \overline{RY} medians.

Result:

\overline{PZ} , \overline{QX} , \overline{RY} are concurrent at O.

4(iii) $m\angle X = 4.3$ cm, $m\angle X = 75^\circ$, $m\angle Y = 45^\circ$



Given: $m\angle X = 4.3$ cm
 $m\angle X = 75^\circ$
 $m\angle Y = 45^\circ$

Required: Construct $\triangle XYZ$ and draw its medians and verify their concurrency

MATHEMATICS FOR 9TH CLASS (UNIT # 17)

Pilot Superone Mathematics 584 Class 9th

Given: $m\overline{AB} = m\overline{AC} = 5.3 \text{ cm}$
 $m\overline{BC} = m\overline{CD} = 3.8 \text{ cm}$
 $m\overline{AD} = 2.8 \text{ cm}$

Required. Construct a quadrilateral ABCD

(ii) Construct a D equal in area to quad. ABCD.

Steps of Construction:

- (i) Take a line segment $\overline{BC} = 3.8 \text{ cm}$
- (ii) Take B, C as centre and draw two arcs of 5.3 cm radius. These intersect each other at A. Join A to B and C.
- (iii) Take A as centre and draw an arc of radius 2.8 cm.
- (iv) Take C as centre and draw an arc of radius 3.8 cm. This cuts the first arc at D.
- (v) Join D to A and C. ABCD is the required quadrilateral.


Part II

- (vi) Join B to D
- (vii) Take $\overline{AP} \parallel \overline{BD}$ which meets extended \overline{CB} at P
- (viii) Join P to D.
 PCD is the required triangle which is equal in area to quadrilateral ABCD

Q.2 Construct a Δ equal in area to the quadrilateral PQRS, having $m\overline{QR} = 7\text{cm}$, $m\overline{RS} = 6\text{cm}$, $m\overline{SP} = 2.75$, $m\angle QRS = 60^\circ$ and $m\angle RSP = 90^\circ$

MATHEMATICS FOR 9TH CLASS (UNIT # 17)

Pilot Superone Mathematics S85 Class 9th



Given: $m\overline{QR} = 7\text{cm}$ $m\overline{RS} = 6\text{cm}$, $m\overline{SP} = 2.75$,
 $m\angle QRS = 60^\circ$ and $m\angle RSP = 90^\circ$

Required: Construct a triangle equal in area to quadrilateral PQRS of which measurements are given.

Steps of Construction:

- (i) Take a line segment $\overline{RS} = 6\text{cm}$
- (ii) Make an angle of 60° at R.
- (iii) Make an angle of 90° at S.
- (iv) Cut off $\overline{SP} = 2.75\text{ cm}$.
- (v) Cut off $\overline{RQ} = 7\text{ cm}$

PQRS is the required quadrilateral

Part II

- (vi) Join P to R. \overline{PR} is diagonal of the quadrilateral
- (vii) Take $\overline{QA} \parallel \overline{RP}$ which meets extended SR at A.
- (viii) Join A to P


Result:

ASP is the required Δ of which area is equal to quadrilateral PQRS

MATHEMATICS FOR 9TH CLASS (UNIT # 17)

Pilot Superone Mathematics 586 Class 9th

Q.3 Construct a Δ equal in area to the quadrilateral ABCD having $m\overline{AB} = 6$ cm, $m\overline{BC} = 4$ cm, $m\overline{AC} = 7.2$ cm, $m\angle BAD = 105^\circ$ and $\overline{BD} = 8$ cm.



Given: $m\overline{AB} = 6$ cm, $m\overline{BC} = 4$ cm, $m\overline{AC} = 7.2$ cm,
 $m\angle BAD = 105^\circ$ and $\overline{BD} = 8$ cm

Required: Construct a quadrilateral ABCD and a Δ equal in area of ABCD.

Steps of Construction.

- (i) Take a line segment $\overline{AB} = 6$ cm
- (ii) Make an angle $\angle BAW = 105^\circ$ at A.
- (iii) Take B as centre and draw an arc of radius 8 cm which cuts \overline{AW} at D.
- (iv) Take A and B as centre and draw arcs of radius 7.2 and 4 cm respectively. These intersect at C.
- (v) Join C to B and D.
- (vi) ABCD is the required quadrilateral

Part II

- (vi) Join C to A.
- (vii) Take $\overline{CP} \parallel \overline{CA}$ which meets BA extended at P


Result:

MATHEMATICS FOR 9TH CLASS (UNIT # 17)

Pilot Superone Mathematics 587 Class 9th

CPB is the required Δ which is equal in area to quadrilateral ABCD

Q.4 Construct a right angled triangle equal in area to a given square.



Given: ABCD is a given square

Required: Construct a right angled triangle equal in area to square ABCD.

Steps of Construction:

- (i) Take \overline{CA} the diagonal of the square ABCD
- (ii) Extend \overline{BA} towards A
- (iii) Take $\overline{DP} \parallel \overline{CA}$ which meets extended BA at P

Result:


PBC is the required Δ equal in area to square ABCD

EXERCISE 17.4

Q.1 Construct a Δ with sides 4 cm, 5 cm and 6 cm and construct a rectangle having its area equal to that of the Δ . Measure its diagonals. Are they equal?

MATHEMATICS FOR 9TH CLASS (UNIT # 17)

Pilot Superone Mathematics 588 Class 9th



Given: $mBC = 6\text{cm}$
 $mAB = 5\text{cm}$
 $mAC = 4\text{cm}$

Required: To construct the triangle and a rectangle equal in area to the Δ

Steps of Construction:

- (i) Take line segment $\overline{BC} = 6\text{ cm}$.
- (ii) Take B and C as centre and draw arc of radii 4, 5 cm respectively. These arcs intersect at A.
- (iii) Join A to B, C
 ΔABC is the required triangle
- (iv) Draw $\overline{NAP} \parallel$ to \overline{BC} passing through A.
- (v) Take right bisector WS which cuts \overline{BC} at D and \overline{NP} at Q.
- (vi) Take Q as centre and draw an arc of radius \overline{DQ}
- (vii) Take C as centre and draw an arc of radius \overline{DQ} , which cuts the first arc at P
- (viii) Join P to Q and C.


MATHEMATICS FOR 9TH CLASS (UNIT # 17)

Prior Superone Mathematics 589 Class 9th

Result:

DCPQ is the required rectangle which is equal in area to $\triangle ABC$ $mDP = mQC = 4.5$ cm

Q.2 Transform an isosceles \triangle into a rect angle.



Given: ABC is an isosceles \triangle in which $\overline{AB} \cong \overline{AC}$

Required: To construct a rectangle equal in area to the \triangle .

Steps of Construction:

- (i) Take $\overline{SAT} \parallel \overline{BC}$ passing through A.
- (ii) Take right bisector of \overline{BC} which cuts \overline{BC} at D and passes through A.
- (iii) Take A as centre and draw an arc of radius \overline{AD} which cuts \overline{ST} at P.
- (iv) Join P to C


Result:

ADCP is the required rectangle which is equal in area to $\triangle ABC$.

MATHEMATICS FOR 9TH CLASS (UNIT # 17)

Plus Superong Mathematics 590 Class 9th

Q.3 Construct a ΔABC such that $m\overline{AB} = 3$ cm, $m\overline{BC} = 3.8$ cm, $m\overline{AC} = 4.8$ cm. Construct a rectangle equal in area to the ΔABC , and measure its sides.



Given: ABC is a triangle that
 $m\overline{AB} = 3$ cm
 $m\overline{BC} = 3.8$ cm
 $m\overline{AC} = 4.8$ cm

Required: Construct the ΔABC and a rectangle equal in area to ΔABC

Steps of Construction:

- Take a line segment $\overline{CA} = 4.8$ cm.
- Take C and A as centre and draw arcs of 3.8 cm, 3 cm radius respectively. These cut each other at B .
- Join B to C and A .
 ABC is the required Δ .
- Take $\overline{NB} \parallel \overline{CA}$ which passes through B
- Take right bisector of \overline{CA} which cuts \overline{CA} at D and \overline{NB} at Q .
- Cut off $OP = mOD$ and P to A

MATHEMATICS FOR 9TH CLASS (UNIT # 17)

Plot Super One Mathematics 591 Class 9th


Result:

QDAP is the required rectangle equal in area to $\triangle ABC$

$$m\overline{DA} = 2.4 \text{ cm}, m\overline{DQ} = 2.4 \text{ cm}$$

EXERCISE 17.5

Q.1 Construct a rectangle whose adjacent sides are 2.5 cm and 5 cm respectively. Construct a square having area equal to the given rectangle.



Given: Sides of a rectangle are 2.5 cm, 5 cm

Required: (i) Construct rectangle

(ii) Construct a square equal in area to the rectangle

Steps of Construction:

- (i) Take a line segment $\overline{AB} = 5 \text{ cm}$
 - (ii) Make an angle of 90° at A.
 - (iii) Cut off 2.5 cm AD and \overline{AW}
 - (iv) Draw arcs with centre at D, B respectively of radius 5 cm and 2.5 cm. These cut each other at C.
 - (v) Join C to D, B.
- ABCD is the required rectangle


MATHEMATICS FOR 9TH CLASS (UNIT # 17)

Pilot Supreme Mathematics 592 Class 9th

Part II

- (vi) Extend DC towards C
 - (vii) Cut off $CE = \overline{BC}$ i.e 2.5 cm on \overline{DN}
 - (viii) Find midpoint O of \overline{DE}
 - (ix) Take O as centre and a semicircle on \overline{DE}
 - (x) Extend side \overline{BC} towards C which meets the semicircle at R.
 - (xi) \overline{mRC} is the length of the required square
 - (xii) Cut off $\overline{CP} = \overline{CR}$ on \overline{DE}
 - (xiii) Take R and P as centre and draw two arcs that cut each other at Q.
 - (xiv) Join Q to R and P.
- \overline{CPQR} is the required square whose area is equal to the area of rectangle ABCD

Q.2 Construct a square equal in area to a rectangle whose adjacent sides are 4.5 cm and 2.2 cm respectively. Measure the sides of the square and find its area and compare with the area of the rectangle.



Given: Sides of a rectangle are 2.5 cm, 5 cm.

MATHEMATICS FOR 9TH CLASS (UNIT # 17)

Pink Superone Mathematics 593 Class 9th

Required: Construct the rectangle and construct a square equal in area to the rectangle

Steps of Construction:

- (i) Take a line segment $AB = 4.5$ cm
- (ii) Make an angle $\angle BAW = 90^\circ$ at A.
- (iii) Cut off 2.2 cm from \overline{AW} as \overline{AD} .
- (iv) Take D and B as centre and draw of radius 4.5 and 2.2 cm respectively. These arcs cut each other at C.
- (v) Join C to D and B.
ABCD is the required rectangle.

Part II

- (vi) Extend \overline{DC} towards C.
- (vii) Cut off $\overline{CE} = \overline{CB} = 2.2$ cm on \overline{DN}
- (viii) Find midpoint O of \overline{DE} .
- (ix) Take O as centre and draw a semi-circle on \overline{DE} taking OD as radius
- (x) Extend \overline{BC} towards C which meets the semicircle at R. $m\overline{RC}$ is the length of side of the required square
- (xi) Cut off $\overline{CP} = m\overline{RC}$ on \overline{DN} .
- (xii) Take R and P as centre and draw two arcs of radius \overline{CR} . These cut each other at Q
- (xiii) Join Q to R and P

Result:

CPQR is the required square whose area is equal to area of rectangle ABCD.



MATHEMATICS FOR 9TH CLASS (UNIT # 17)

Pilot Superone Mathematics 594 Class 9th

Part II

$$\begin{aligned}\text{Area of the rectangle} &= 4.5 \times 2.2 \\ &= 9.9 \text{ sq. cm}\end{aligned}$$

$$\text{Side of the square} = 3.1 \text{ cm}$$

$$\begin{aligned}\text{Area of the square} &= 3.1 \times 3.1 \\ &= 9.6 \text{ sq. cm}\end{aligned}$$

Area of the square is approximately equal to the area of the rectangle.

Q.3 In Q.2 above verify by measurement that the perimeter of the square is less than that of the rectangle.

Solution:

$$\begin{aligned}\text{Perimeter of the rectangle} &= 2(4.5 + 2.2) \\ &= 2(6.7) \\ &= 13.4 \text{ cm}\end{aligned}$$

$$\text{Side of the square} = 3.1 \text{ cm}$$

$$\begin{aligned}\text{Perimeter of the square} &= 4(3.1) \\ &= 12.4 \text{ cm}\end{aligned}$$

$$\text{Now } 12.4 < 13.4$$

Therefore:


Perimeter of the square is less than the perimeter of the rectangle.

WIKI

MATHEMATICS FOR 9TH CLASS (UNIT # 17)

Pilot Superone Mathematics 595 Class 9th

Q.4 Construct a square equal in area to the sum of two squares having sides 3 cm and 4 cm respectively.



Given: Two square having 3 cm and 4 cm sides are given.

Required: To construction a square equal in area to the sum of the area of two squares.

Steps of construction:

- (i) Take a line segment $\overline{AB} = 4 \text{ cm}$
- (ii) Complete square ADCB on \overline{AB}
- (iii) Extend \overline{DA} towards A and cut off $\overline{AP} = 3 \text{ cm}$.
OR Make an angle of 90° at A and cut off $\overline{AP} = 3 \text{ cm}$.
- (iv) Construct a square ARQP on \overline{AP}
- (v) Join B to P
ABP is a right angled triangle of which one side is 4 cm and the other side is 3 cm.

MATHEMATICS FOR 9TH CLASS (UNIT # 17)

Pilot Superone Mathematics 596 Class 9th


(vi) Construct a square on hypotenuse \overline{BP}

Result:

$$(mPB)^2 = (mAB)^2 + m(AP)^2$$

(Pythagorean theorem)

- Q.5** Construct a Δ having base 3.5 cm and other two sides equal to 3.4 cm and 3.8 cm respectively. Transform it into an equal square.



Given: ABC is a triangle in which (suppose)

$$m\overline{AB} = 3.5 \text{ cm}$$

$$m\overline{AC} = 3.8 \text{ cm}$$

$$m\overline{BC} = 3.4 \text{ cm}$$

Required: To construct a square equal in area to the triangle ABC

Steps of Construction:

Part I

- (i) Take a line segment $\overline{AB} = 3.5 \text{ cm}$
- (ii) Take A and B as centre and draw two arcs of 3.8 cm and 3.4 cm respectively. These arcs intersect at C.

MATHEMATICS FOR 9TH CLASS (UNIT # 17)

Pilot Superone Mathematics 597 Class 9th

(iii) Join C to A and B.

ABC is the required Δ .

Part II

(i) Take $\overline{MN} \parallel AB$ passing through C

(ii) Take right bisector GH of AB which bisects \overline{AB} at O, and \overline{MN} at I

(iii) Sides line segments OI and OB are the sides of a rectangle

(iv) Complete rectangle OBYI. Its area is equal to the area of ΔABC

Part II

(i) Extend YB towards B.

(ii) Cut off $BK = mBC$ on \overline{YW}

(iii) Find midpoint V of \overline{BK}

(iv) Take V as centre and draw a semicircle on \overline{YK}

(v) Extend \overline{OB} towards B which meets the semicircle at S.

(vi) Complete square on BS square BQRS is equal in area of rectangle OBYI

From Part I, II, III


Square BQRS is equal in area to triangle ABC.



MATHEMATICS FOR 9TH CLASS (UNIT # 17)

Pilot Superone Mathematics SUB Class 9

Q.6 Construct a Δ having base 5 cm and other sides equal to 5 cm and 6 cm. Construct square equal in area to given Δ .



Given: Let ABC be the Δ in which $m\overline{AC} = 5$ cm
 $m\overline{AB} = 6$ cm
 $m\overline{BC} = 5$ cm

Required. Construct a square equal in area to ΔABC .

Steps of construction:

Part I

- (i) Take a line segment $\overline{AB} = 5$ cm
- (ii) Take B and C as centre and draw two arcs of radius 6 cm, 5 cm, respectively. These cut each other at A.
- (iii) Join A to B and C
 ABC is the required Δ .

Part II

- (i) Take $\overline{MN} \parallel \overline{BC}$ passing through A.

MATHEMATICS FOR 9TH CLASS (UNIT # 17)

Pilot Superone Mathematics 599 Class 9th

- (ii) Take WW' right bisector of \overline{BC} at O and cuts MN at E.
- (iii) Complete the rectangle EOCD which is equal in area to $\triangle ABC$

Ex III

- (i) Extend DC side of rectangle towards C and take $mCK = mCO$
- (ii) Find F the midpoint of \overline{DK}
- (iii) Take F as centre and complete a semicircle on \overline{DK}
- (iv) Extend \overline{OC} towards C, this cut the semicircle on R.
- (v) Complete square CRQP and its area is equal to the area of rectangle EOCD

Result

Area of square CRQP is equal to the area of rectangle EOCD which is equal to the area of the triangle ABC.

16



MATHEMATICS FOR 9TH CLASS (UNIT # 17)

Pilot Superone Mathematics 600 Class 9th

Review Exercise 17

1. Fill in the following blanks to make the statement true:

- (i) The side of a right angled triangle opposite to 90° is called _____
- (ii) The line segment joining a vertex of a triangle to the mid-point of its opposite side is called a _____
- (iii) A line drawn from a vertex of a triangle which is _____ to its opposite side is called an altitude of the triangle.
- (iv) The bisectors of the three angles of a triangle are _____
- (v) The point of concurrency of the right bisectors of the three sides of the triangle is _____ from its vertices.
- (vi) Two or more triangles are said to be similar if they are equiangular and measures of their corresponding sides are _____
- (vii) The altitudes of a right triangle are concurrent at the _____ of the right angle.

Answers:

- | | |
|---------------------|-------------------|
| (i) hypotenuse | (ii) median |
| (iii) perpendicular | (iv) concurrent |
| (v) equidistant | (vi) proportional |
| (vii) vertex | |

Q.2 Multiple Choice Questions. Choose the correct answer.

- (i) A triangle having two sides congruent is called:
(a) scalene (b) right angled
(c) equilateral (d) isosceles
- (ii) A quadrilateral having each angle equal to 90° is called:

MATHEMATICS FOR 9TH CLASS (UNIT # 17)

Pilot Superone Mathematics 601 Class 9th

- (a) parallelogram (b) rectangle
(c) trapezium (d) rhombus
- (iii) The right bisectors of the three sides of a triangle are:
(a) congruent (b) collinear
(c) concurrent (d) parallel
- (iv) The altitudes of an isosceles triangle are congruent:
(a) two (b) three
(c) four (d) none
- (v) A point equidistant from the end points of a line segment is on its:
(a) bisector (b) right-bisector
(c) perpendicular (d) median
- (vi) congruent triangles can be made by joining the mid-point of the sides of a triangle:
(a) three (b) four
(c) five (d) two
- (vii) The diagonals of a parallelogram each other:
(a) bisect (b) trisect
(c) bisect at right angle (d) none of these
- (viii) The medians of a triangle cut each other in the ratio:
(a) 4 : 1 (b) 3 : 1
(c) 2 : 1 (d) 1 : 1
- (ix) One angle on the base of an isosceles triangle is 30° .
What is the measure of its vertical angle:
(a) 30° (b) 60°
(c) 90° (d) 120°
- (x) If the three altitudes of a triangle are congruent, then the triangle is:
(a) equilateral (b) right angled
(c) isosceles (d) acute angled

MATHEMATICS FOR 9TH CLASS (UNIT # 17)

Pilot Superone Mathematics 602 Class 9th

(xi) If two medians of a triangle are congruent then the triangle will be:

- (a) isosceles (b) equilateral
 (c) right angled (d) acute angled

Answers:

(i) d	(ii) b	(iii) c	(iv) a
(v) b	(vi) b	(vii) a	(viii) c
(ix) d	(x) a	(xi) a	

Q.3. Define the following

- (i) In-centre (ii) Circum-centre
 (iii) Ortho-centre (iv) Centroid
 (v) Point of concurrency
 (i) In-centre: The point of concurrency of the bisectors of interior angles of a triangle is called its In-centre.
 (ii) Circum-centre: The point of concurrency of the three right bisectors of sides of a triangle is called its circum-centre.
 (iii) Ortho-centre: The point of concurrency of the three altitudes of a triangle is called its Ortho-centre.
 (iv) Centroid: The point of concurrency of three medians of a triangle is called its Centroid.
 (v) Point of concurrency: If three or more than three straight lines, rays, line-segments, intersect each other at a single point then that point is a point of concurrency of these straight lines, rays or line-segments.

W11

